

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification. 7 : <b>C12N 15/12, 15/82, 5/10, A01H 5/00</b>		<b>A1</b>	(11) International Publication Number: <b>WO 00/26371</b>
			(43) International Publication Date: <b>11 May 2000 (11.05.00)</b>
(21) International Application Number: <b>PCT/US99/26086</b>		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: <b>4 November 1999 (04.11.99)</b>			
(30) Priority Data: <b>09/186,002</b> <b>4 November 1998 (04.11.98)</b> <b>US</b>			
(63) Related by Continuation (CON) or Continuation-in-Part (CIP) to Earlier Application <b>US</b> <b>09/186,002 (CON)</b> <b>Filed on</b> <b>4 November 1998 (04.11.98)</b>			
(71) Applicant (for all designated States except US): <b>MONSANTO CO. [US/US]; 800 N. Lindbergh Boulevard, St. Louis, MO 63167 (US).</b>		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(72) Inventors; and			
(75) Inventors/Applicants (for US only): <b>CORBIN, David, R. [US/US]; 14453 Britannia Drive, Chesterfield, MO 63017 (US). ROMANO, Charles, P. [US/US]; 38 Charlesdale Road, Medfield, MA (US).</b>			
(74) Agent: <b>KAMMERER, Patricia, A.; Arnold White &amp; Durkee, 750 Bering Drive, Houston, TX 77057-2198 (US).</b>			
(54) Title: <b>METHODS FOR TRANSFORMING PLANTS TO EXPRESS BACILLUS THURINGIENSIS DELTA-ENDOTOXINS</b>			
(57) Abstract <p>Disclosed is a means of controlling plant pests by a novel method of expressing Cry2A <i>B. thuringiensis</i> <math>\delta</math>-endotoxins in plants. The invention comprises novel nucleic acid segments encoding proteins comprising Cry2A <i>B. thuringiensis</i> <math>\delta</math>-endotoxins. The nucleic acid segments are disclosed, as are transformation vectors containing the nucleic acid segments, plants transformed with the claimed segments, methods for transforming plants, and methods of controlling plant infestation by pests.</p>			

*FOR THE PURPOSES OF INFORMATION ONLY*

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

## METHODS FOR TRANSFORMING PLANTS TO EXPRESS BACILLUS THURINGIENSIS DELTA-ENDOTOXINS

**1.0 Background of the Invention****1.1 Field of the Invention**

The present invention relates generally to transgenic plants having insecticidal capabilities, and to DNA constructs utilized to transfer genes conferring insect resistance into plant genomes. More specifically, the present invention relates to a method of expressing insecticidal proteins in plants transformed with a *B. thuringiensis*  $\delta$ -endotoxin encoding gene, resulting in effective control of susceptible target pests.

**1.2 Description of Related Art****1.2.1 Methods of Controlling Insect Infestation in Plants**

The Gram-positive soil bacterium *B. thuringiensis* is well known for its production of proteinaceous parasporal crystals, or  $\delta$ -endotoxins, that are toxic to a variety of Lepidopteran, Coleopteran, and Dipteran larvae. *B. thuringiensis* produces crystal proteins during sporulation which are specifically toxic to certain species of insects. Many different strains of *B. thuringiensis* have been shown to produce insecticidal crystal proteins. Compositions comprising *B. thuringiensis* strains which produce proteins having insecticidal activity have been used commercially as environmentally-acceptable topical insecticides because of their toxicity to the specific target insect, and non-toxicity to plants and other non-targeted organisms.

$\delta$ -endotoxin crystals are toxic to insect larvae by ingestion. Solubilization of the crystal in the midgut of the insect releases the protoxin form of the  $\delta$ -endotoxin which, in most instances, is subsequently processed to an active toxin by midgut protease. The activated toxins recognize and bind to the brush-border of the insect midgut epithelium through receptor proteins. Several putative crystal protein receptors have been isolated from certain insect larvae (Knight *et al.*, 1995; Gill *et al.*, 1995; Masson *et al.*, 1995). The binding of active toxins is followed by intercalation and aggregation of toxin molecules to form pores within the midgut epithelium. This process leads to osmotic imbalance, swelling, lysis of the cells lining the midgut epithelium, and eventual larvae mortality.

**1.2.2 Transgenic *B. thuringiensis*  $\delta$ -Endotoxins as Biopesticides**

Plant resistance and biological control are central tactics of control in the majority of insecticide improvement programs applied to the most diverse crops. With the advent of molecular genetic techniques, various  $\delta$ -endotoxin genes have been isolated and their DNA sequences determined. These genes have been used to construct certain genetically engineered *B. thuringiensis* products that have been approved for commercial use. Recent developments have seen new  $\delta$ -endotoxin delivery systems developed, including plants that contain and express genetically engineered  $\delta$ -endotoxin genes.

Expression of *B. thuringiensis*  $\delta$ -endotoxins in plants holds the potential for effective management of plant pests so long as certain problems can be overcome. These problems include the development of insect resistance to the particular Cry protein expressed in the plant, and development of morphologically abnormal plants because of the presence of the transgene.

5        Expression of *B. thuringiensis*  $\delta$ -endotoxins in transgenic cotton, corn, and potatoes has proven to be an effective means of controlling agriculturally important insect pests (Perlak *et al.*, 1990; Koziel *et al.*, 1993; Perlak *et al.*, 1993). Transgenic crops expressing *B. thuringiensis*  $\delta$ -endotoxins enable growers to significantly reduce the application of costly, toxic, and sometimes ineffective topical chemical insecticides. Use of transgenes encoding *B. thuringiensis*  $\delta$ -endotoxins is particularly advantageous when  
10       insertion of the transgene has no negative effect on the yield of desired product from the transformed plants. Yields from crop plants expressing certain *B. thuringiensis*  $\delta$ -endotoxins such as Cry1A or Cry3A have been observed to be equivalent or better than otherwise similar non-transgenic commercial plant varieties. This indicates that expression of some *B. thuringiensis*  $\delta$ -endotoxins does not have a significant negative impact on plant growth or development. This is not the case, however, for all *B.*  
15       *thuringiensis*  $\delta$ -endotoxins that may be used to transform plants.

      The use of topical *B. thuringiensis*-derived insecticides may also result in the development of insect strains resistant to the insecticides. Resistance to Cry1A *B. thuringiensis*  $\delta$ -endotoxins applied as foliar sprays has evolved in at least one well documented instance (Shelton *et al.*, 1993). It is expected that insects may similarly evolve resistance to *B. thuringiensis*  $\delta$ -endotoxins expressed in transgenic  
20       plants. Such resistance, should it become widespread, would clearly limit the commercial value of corn, cotton, potato, and other germplasm containing genes encoding *B. thuringiensis*  $\delta$ -endotoxins. One possible way to both increase the effectiveness of the insecticide against target pests and to reduce the development of insecticide-resistant pests would be to ensure that transgenic crops express high levels of  
      *B. thuringiensis*  $\delta$ -endotoxins (McGaughey and Whalon, 1993; Roush, 1994).

25       In addition to producing a transgenic plant which expresses *B. thuringiensis*  $\delta$ -endotoxins at high levels, commercially viable *B. thuringiensis* genes must satisfy several additional criteria. For instance, expression of these genes in transgenic crop plants must not reduce the vigor, viability or fertility of the plants, nor may it affect the normal morphology of the plants. Such detrimental effects have two undesired results: they may interfere with the recovery and propagation of transgenic plants; they may  
30       also impede the development of mature plants, or confer unacceptable agronomic characteristics.

      There remains a need for compositions and methods useful in producing transgenic plants which express *B. thuringiensis*  $\delta$ -endotoxins at levels high enough to effectively control target plant insect pests as well as prevent the development of insecticide-resistant pest strains. A method resulting in higher levels of expression of the *B. thuringiensis*  $\delta$ -endotoxins will also provide the advantages of more

frequent attainment of commercially viable transformed plant lines and more effective protection from infestation for the entire growing season.

There also remains a need for a method of increasing the level of expression of *B. thuringiensis*  $\delta$ -endotoxins which does not simultaneously result in plant morphological changes that interfere with optimal growth and development of desired plant tissues. For example, the method of potentiating expression of the *B. thuringiensis*  $\delta$ -endotoxins in corn should not result in a corn plant which cannot optimally develop for cultivation. Achievement of these goals such as high expression levels as well as recovery of morphologically normal plants has been elusive, and their pursuit has been ongoing and an important aspect of the long term value of insecticidal plant products.

## 2.0 Summary of the Invention

Described are novel methods for expressing Cry2A *B. thuringiensis*  $\delta$ -endotoxins which lack significant Dipteran inhibiting activity in transformed plants. This method advantageously results in both increased levels of expression of *B. thuringiensis*  $\delta$ -endotoxins as well as a higher rate of recovery of morphologically-normal plants.

By achieving high rates of expression, the present invention addresses another limitation of the prior art: development of insect resistance. Specifically, the instant invention provides a superior strategy for the delay or elimination of the development of resistance to Cry1A  $\delta$ -endotoxins, the *B. thuringiensis* proteins most commonly expressed by transgenic lines. The disclosed methods involve expression of the Cry2A class of *B. thuringiensis*  $\delta$ -endotoxins and particularly those that lack Dipteran-inhibiting activity. *B. thuringiensis*  $\delta$ -endotoxins of the Cry2A group have no significant homology to Cry1A-type  $\delta$ -endotoxins and display distinct binding and pore-forming characteristics (English *et al.*, 1994), and as such are expected to control insects that become resistant to, or that are not affected by, Cry1A  $\delta$ -endotoxins (Hofte and Whiteley, 1989).

In preferred embodiments, the present invention provides an isolated and purified DNA construct comprising a Cry2A  $\delta$ -endotoxin-encoding region localized to a plastid or chloroplast, or localized to a plant cell nuclear genome and operably linked to a region encoding a plastid transit peptide (PTP). Preferred DNA constructs of the present invention include those constructs that encode Cry2A  $\delta$ -endotoxins lacking Dipteran-inhibitory activity, though complete inactivity towards Dipterans is not required. In an illustrative embodiment, DNA constructs of the present invention encode a Cry2Ab  $\delta$ -endotoxin operably linked to a DNA segment (or sequence) encoding a plastid transit peptide, which is one means of enabling localization of a Cry2Ab  $\delta$ -endotoxin to a plastid or chloroplast. In certain embodiments, the Cry2Ab  $\delta$ -endotoxin comprises the sequence of SEQ ID NO:2. The inventors contemplate, however, that any Cry2A  $\delta$ -endotoxin lacking Dipteran-inhibitory activity may be utilized according to the present invention, with those bearing substantial homologies to Cry2Ab being particularly preferred.

In another embodiment, the DNA constructs of the present invention exploit nucleic acid segments encoding PTPs to potentiate expression of the  $\delta$ -endotoxin. The use of one type of PTP, a chloroplast targeting peptide (CTP), in conjunction with a *cryIA B. thuringiensis* transgene to promote expression of the transgene in the transformed plant is disclosed in U. S. Patent 5,500,365 (specifically incorporated herein by reference in its entirety). Where increased expression was observed, however, it was ascribed in part to the use of a new 5' untranslated leader sequence in the expression vector.

In contrast to the prior art, the present invention discloses a structural DNA sequence that causes the production of an RNA sequence which encodes a targeted fusion protein comprising an amino-terminal plastid transit peptide with a Cry2Ab  $\delta$ -endotoxin; and a 3' non-translated DNA sequence which functions in plant cells to cause transcriptional termination and the addition of polyadenylated nucleotides to the 3' end of the RNA sequence. Surprisingly, this DNA construct results in increased levels of expression of the Cry2A  $\delta$ -endotoxin. The targeted fusion protein is non-active to all species, but is produced as a means for localizing the mature, insecticidally active  $\delta$ -endotoxin protein to the chloroplast, yielding surprising and unexpected beneficial agronomic effects.

One embodiment conceived of in the present invention is the introduction of a gene encoding a Cry2A  $\delta$ -endotoxin lacking Dipteran activity into the chloroplast or plastid genome. Alternatively, a gene encoding a Cry2A  $\delta$ -endotoxin lacking Dipteran activity could be expressed from an autonomously replicating episomal element located within the chloroplast or plastid.

In another preferred embodiment, the invention provides for transgenic plants which have been transformed with an isolated and purified DNA construct that is translated and expressed at high levels by the plant. Both monocot and dicot plants may be transformed according to the methods and with the DNA constructs disclosed herein. The plant transformed by the instant invention may be prepared, in a further preferred embodiment, by a process including obtainment of the isolated and purified DNA construct, and then transforming the plant with the construct so that the plant expresses the proteins for which the construct encodes. The inventors have observed that transformation of plants by the disclosed methods results in increased frequency of transformants which express the transgene, as well as the generation of more morphologically normal plants from initial transformants.

It is contemplated that the increased expression levels observed in the disclosed invention will allow for reduced development of insect resistance to Bt  $\delta$ -endotoxins. This may be achieved by transforming a plant with the preferred DNA construct to achieve high rates of Cry2A expression alone, or by simultaneously exposing target insects to Cry1A and non-Dipteran active Cry2A  $\delta$ -endotoxins expressed in susceptible plants. Such insects include *Ostrina* spp., *Diatraea* spp., *Helicoverpa* spp., and *Spodoptera* spp., in *Zea mays*; *Heliothis virescens*, *Helicoverpa* spp., *Pectinophora* spp., in *Gossypium hirsutum*; *Anticarsia* spp., *Pseudoplusia* spp., *Epinotia* spp., in *Glycine max*; and *Scirpophaga incertulas* in *Oryza sativa*.

It is therefore contemplated that the method disclosed by the present invention will provide many advantages over the prior art including those specifically outlined above. These advantages include: obtaining improved control of susceptible insects; minimizing the development of insecticide-resistant insect strains; obtaining a greater number of commercially viable insect resistant plant lines; achieving season long protection from insect pathogens; and increasing the incidence of morphologically-normal transformed plants. An additional advantage of the present invention is that reduced numbers of transgenic lines would need to be produced in order to identify a transgenic event with normal growth characteristics.

It is therefore contemplated that the method disclosed by the present invention will provide many advantages over the prior art including those specifically outlined above. These advantages include: obtaining improved control of susceptible insects; minimizing the development of insecticide-resistant insect strains; obtaining a greater number of commercially viable insect resistant plant lines; achieving season long protection from insect pathogens; and increasing the incidence of morphologically-normal transformed plants. An additional advantage of the present invention is that reduced numbers of transgenic lines would need to be produced in order to identify a transgenic event with normal growth characteristics.

## 2.1 Nucleic Acid Compositions

In one important embodiment, the invention provides an isolated and purified nucleic acid construct comprising a Cry2A coding region and a PTP coding region. These DNA constructs, when transferred into a plant, undergo cellular processes resulting in increased expression of  $\delta$ -endotoxins in the transgenic plant. The Cry2A endotoxins of the instant invention are preferably not effective against Dipteran species, though some adverse effects on Dipterans may be tolerated. In certain embodiments, the DNA construct encodes a Dipteran-inactive Cry2Ab  $\delta$ -endotoxin, and in more preferred embodiments, the Cry2Ab  $\delta$ -endotoxin has the polypeptide sequence of SEQ ID NO:2, or one substantially homologous to the polypeptide sequence of SEQ ID NO:2. Such nucleotide homologues may be greater than approximately 88% homologous, greater than about 90% homologous, greater than about 95% homologous, and even greater than about 99% homologous with the Cry2Ab  $\delta$ -endotoxin disclosed in SEQ ID NO:2. Exemplary peptides include those that are about 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98 or even 99 or greater percent homologous to the Cry2Ab  $\delta$ -endotoxin disclosed in SEQ ID NO:2.

In even more preferred embodiments, the DNA construct of the present invention comprises a Cry2Ab  $\delta$ -endotoxin-encoding region with the nucleic acid sequence of SEQ ID NO:1, or a sequence substantially homologous to that of SEQ ID NO:1. Also envisioned as within the scope of this invention are those DNA constructs having segments with substantial homologies to the nucleic acid sequence disclosed in SEQ ID NO:1, such as those which may be about 90% homologous, or about 95 %

homologous, or even about 99% homologous. More specifically, homologous nucleic acid sequences included in the present invention include those that are about 90, 91, 92, 93, 94, 95, 96, 97, 98, and 99 percent homologous to the nucleic acid sequence of SEQ ID NO:1.

The DNA constructs provided herein also include a PTP coding region positioned upstream of the *cry2A*  $\delta$ -endotoxin coding region and downstream of a promoter. These plastid transit peptide coding regions may encode any plant functional PTP, and may operate to target encoded proteins to certain plastids within the plant cell, or to increase the expression of the  $\delta$ -endotoxin for which the DNA construct encodes. In preferred embodiments, the present invention may include a PTP selected from the group including zmSSU, PTPI, PTPIA, and PTP2, or any other plant functional PTPs. More preferably, the plastid transit peptide coding region encodes a plastid transit peptide having the amino acid sequence of SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, or SEQ ID NO:10, or any polypeptide sequence substantially homologous to these. Even more preferably, the instant invention comprises a plastid transit peptide coding region having the nucleic acid sequence of SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, or SEQ ID NO:9, or a nucleic acid sequence which is substantially homologous to these.

Also, the inventors contemplate that the present invention would further achieve the goals of increased pathogenicity to pests, and result in decreased development of pesticide-resistant insects, if the DNA constructs provided herein were co-expressed along with other pesticidal compositions such as other proteins. Accordingly, the invention provides for use of the disclosed DNA constructs which further comprise plant-expressible coding regions for other Cry proteins. Included in these would be coding regions for CryI proteins such as CryIA, CryIAb, CryIBb, or CryI chimeras (see co-pending US applications Serial No.'s 08/754,490 and 08/922,505, and co-pending PCT Application PCT/US97/17507 based on US Application Serial No. 08/721,259, each specifically incorporated herein by reference in its entirety).

In certain preferred embodiments, the DNA construct is an expression cassette which can be excised and isolated from said plasmid.

## 2.2 Additional Nucleic acid Composition Elements

The polynucleotide compositions of the present invention are useful in transforming both monocotyledonous and dicotyledonous plants. Accordingly, the DNA construct of the present invention may further comprise other various regulatory elements to aid in protein expression and to further facilitate introduction of the DNA construct into the plant. One example of this is the inclusion, in the DNA construct, of an intron positioned in the untranslated leader, upstream relative to the plastid transit peptide coding region. One useful leader sequence is the petunia heat shock protein. In various alternative embodiments, the intron may be any of the following: Adh intron 1, sucrose synthase intron, TMV omega element, maize heat shock protein (hsp) 70, or the rice *Act1* intron. In preferred embodiments, the intron is either maize heat shock protein 70 or petunia heat shock protein 70.



Provided in another preferred embodiment of the present invention is a polynucleotide sequence comprising a substantially Dipteran inactive *cry2A*  $\delta$ -endotoxin coding region and a PTP coding region positioned under the control of a plant operable promoter. The use of a promoter is required for driving cellular processes so that expression of the gene is maximized. Preferred promoters include the following: CaMV 35S, histone, CaMV 19S, *nos*, OCS, *Adh*, sucrose synthase,  $\alpha$ -tubulin, actin, cab, PEPCase, ssRUBISCO, *Act1*, *Famv*, enhanced FMV, or R-gene complex associated promoters. In more preferred embodiments, the promoter is the enhanced or duplicated CaMV 35S promoter (Kay *et al.*, 1987). In additional preferred embodiments, the promoter is the FMV35S promoter. Plant chloroplast or plastid functional promoters are also within the scope of the present invention.

The present invention further contemplates the inclusion of a terminator region in the DNA construct to aid cellular processes involved with protein expression. In various embodiments, this terminator may be any of the following: the *Agrobacterium tumefaciens* nopaline synthase gene terminator, the *Agrobacterium tumefaciens* octopine synthase gene terminator, and the 3' end of the protease inhibitor I or II genes from potato or tomato. In an especially preferred embodiment, the terminator is the *Agrobacterium tumefaciens* nopaline synthase gene terminator.

### 2.3 Transformation Vectors

Because the DNA construct of the present invention is primarily, though not exclusively, intended for use in the transformation of plants, it is in certain preferred embodiments, contained within an expression vector. Such expression vectors may contain a variety of regulatory and other elements intended to allow for optimal expression of the desired proteins for which the expression vector encodes. These additional elements may include promoters, terminators, and introns as outlined above in section 2.2. The vector containing the DNA construct and any regulatory or other elements may be selected from the group consisting of a yeast artificial chromosome, bacterial artificial chromosome, a plasmid, or a cosmid.

Further, the expression vectors themselves may be of a variety of forms. These forms may differ for various reasons, and will likely be comprised of varying components depending upon whether they are intended to transform a monocotyledonous plant or a dicotyledonous plant. For example, FIG. 1 illustrates one possible embodiment, where the monocotyledonous expression vector contains the *cry2Ab* gene in the plasmid designated as (SEQ ID NO:16). It is further contemplated that other expression vectors containing the expression cassettes embodied in these plasmid vectors, as well as expression cassettes containing substantial homologues, will also be useful transformation constructs. Accordingly, any transformation vector containing the nucleic acid sequence of from nucleic acid 1781 to 5869 of SEQ ID NO:16.

FIG. 2 illustrates one possible dicotyledonous expression vector. It contains the *cry2Ab* gene embodied in the plasmids designated as pMON33827 (SEQ ID NO:13), pMON33828 (SEQ ID NO:14), and pMON33829 (SEQ ID NO:15). As with the illustrative monocotyledonous transformation vectors,

the inventors further contemplate that other expression vectors containing the expression cassettes embodied in these plasmid vectors, or substantial homologues to those expression cassettes, will be useful as dicotyledonous transformation constructs. Preferred dicotyledonous expression cassettes include those embodied by nucleic acids 17 to 3182 of SEQ ID NO:13; nucleic acids 17 to 3092 of SEQ ID NO:14; and  
5 nucleic acids 17 to 3155 of SEQ ID NO:15. Illustrative embodiments of vectors containing such expression cassettes are disclosed in the sequences designated herein as SEQ ID NO:13, SEQ ID NO:14, and SEQ ID NO:15.

Vectors further envisioned to be within the scope of the present invention include those vectors capable of containing both the Dipteran-inactive *cry2A* nucleic acid compositions disclosed in section 2.1  
10 above, as well as any other DNA constructs which further comprise plant-expressible coding regions for other Cry proteins such as a Cry1 protein. Vectors capable of containing both of these constructs may further comprise an internal ribosome entry site between the DNA construct; they may also contain a variety of different cistrons, rendering them polycistronic or multicistronic

#### 2.4 Transformed Host Cells

15 Another preferred embodiment of the present invention encompasses cells transformed with the DNA constructs disclosed herein in sections 2.1 and 2.2, and by use of the transformation vectors disclosed in section 2.3. Transformed cells contemplated in the present invention include both prokaryotic and eukaryotic cells which express the proteins encoded-for by the novel DNA constructs of the present invention. The process of producing transgenic cells is well-known in the art. In general, the  
20 method comprises transforming a suitable host cell with a DNA segment which contains a promoter operatively linked to a coding region that encodes a *B. thuringiensis*  $\delta$ -endotoxin. Such a coding region is generally operatively linked to a transcription-terminating region, whereby the promoter is capable of driving the transcription of the coding region in the cell, and hence providing the cell the ability to produce the  $\delta$ -endotoxin *in vivo*. Alternatively, in instances where it is desirable to control, regulate, or  
25 decrease the amount of a particular  $\delta$ -endotoxin or endotoxins expressed in a particular transgenic cell, the invention also provides for the expression of  $\delta$ -endotoxin antisense mRNA; intron antisense mRNA; PTP antisense mRNA; or UTR antisense mRNA. The use of antisense mRNA as a means of controlling or decreasing the amount of a given protein of interest in a cell is well-known in the art.

In a preferred embodiment, the invention encompasses a plant cell which has been transformed  
30 with a nucleic acid segment or DNA construct of the invention, and which expresses a gene or gene segment encoding one or more of the Dipteran-inactive Cry2A *B. thuringiensis*  $\delta$ -endotoxins as disclosed herein. As used herein, the term "transgenic plant cell" is intended to refer to a plant cell that has incorporated DNA sequences, including but not limited to genes which are perhaps not normally present, DNA sequences not normally transcribed into RNA or translated into a protein ("expressed"), or any  
35 other genes or DNA sequences which one desires to introduce into the non-transformed plant, such as

genes which may normally be present in the non-transformed plant but which one desires to either genetically engineer or to have altered expression.

It is contemplated that in some instances the genome of a transgenic plant of the present invention will have been augmented through the stable introduction of a Dipteran-inactive Cry2A *B. thuringiensis*  $\delta$ -endotoxin-encoding DNA constructs as disclosed in sections 2.1 and 2.2 above. In some instances, more than one transgene will be incorporated into the nuclear genome, or into the chloroplast or plastid genome of the transformed host plant cell. Such is the case when more than one crystal protein-encoding DNA segment is incorporated into the genome of such a plant. In certain situations, it may be desirable to have one, two, three, four, or even more *B. thuringiensis* crystal protein-encoding polynucleotides (either native or recombinantly-engineered) incorporated and stably expressed in the transformed transgenic plant.

In preferred embodiments, the introduction of the transgene into the genome of the plant cell results in a stable integration wherein the offspring of such plants also contain a copy of the transgene in their genome. The heritability of this genetic element by the progeny of the plant into which the gene was originally introduced is a preferred aspect of this invention. A preferred gene which may be introduced includes, for example a *B. thuringiensis*  $\delta$ -endotoxin, and particularly one or more of those described herein.

Means for transforming a plant cell and the preparation of a transgenic cell line are well-known in the art (as exemplified in U. S. Patents 5,550,318; 5,508,468; 5,482,852; 5,384,253; 5,276,269; and 5,225,341, all specifically incorporated herein by reference in their entirety), and are briefly discussed herein. Vectors, plasmids, cosmids, YACs (yeast artificial chromosomes) and DNA segments for use in transforming such cells will, of course, generally comprise either the operons, genes, or gene-derived sequences of the present invention, either native, or synthetically-derived, and particularly those encoding the disclosed crystal proteins. These DNA constructs can further include structures such as promoters, enhancers, polylinkers, or even gene sequences which have positively- or negatively-regulating activity upon the particular genes of interest as desired. The DNA segment or gene may encode either a native or modified crystal protein, which will be expressed in the resultant recombinant cells, and/or which will impart an improved phenotype to the regenerated plant.

Transgenic cells specifically contemplated in the present invention include transgenic plant cells. Particularly preferred plant cells include those cells obtained from corn, wheat, soybean, turf grasses, ornamental plant, fruit tree, shrubs, vegetables, grains, legumes, and the like, or any plant into which introduction of a Dipteran-inactive *B. thuringiensis*  $\delta$ -endotoxin transgene is desired.

## 2.5 Transformed Plants

In another aspect, plants transformed with any DNA construct of the present invention that express the proteins for which the construct encodes, are contemplated as being a part of this invention. Accordingly, the invention further provides transgenic plants which have been transformed with a DNA

construct, as disclosed herein in sections 2.1 and 2.2, and transformed by use of transformation vectors as disclosed in section 2.3. Agronomic, horticultural, ornamental, and other economically or commercially useful plants can be made in accordance with the methods described herein, to express *B. thuringiensis*  $\delta$ -endotoxins at levels high enough to confer resistance to insect pathogens while remaining morphologically normal.

Such plants may co-express the  $\delta$ -endotoxin polypeptide along with other antifungal, antibacterial, or antiviral pathogenesis-related peptides, polypeptides, or proteins; insecticidal proteins; proteins conferring herbicide resistance; and proteins involved in improving the quality or quantity of plant products or agronomic performance of plants. Simultaneous co-expression of multiple proteins in plants is advantageous in that it exploits more than one mode of action to control plant pathogenic damage. This can minimize the possibility of developing resistant pathogen strains, broaden the scope of resistance, and potentially result in a synergistic insecticidal effect, thereby enhancing a plant's ability to resist insect infestation (Intl. Patent Appl. Publ. No. WO 92/17591, 15 October 1992, specifically incorporated herein by reference in its entirety).

The transformed plant of the current invention may be either a monocotyledonous plant or a dicotyledonous plant. Where the plant is a monocotyledonous plant, it may be any one of a variety of species. Preferred monocotyledonous species encompassed by the present invention may include maize, rice, wheat, barley, oats, rye, millet, sorghum, sugarcane, asparagus, turfgrass, or any of a number of other grains or cereal plants. In preferred embodiments, the monocot is a maize plant.

The present invention also contemplates a variety of dicotyledonous plants such as cotton, soybean, tomato, potato, citrus, tobacco, sugar beet, alfalfa, fava bean, pea, bean, apple, cherry, pear, strawberry, raspberry, or any other legume, tuber, or fruit plant. In preferred embodiments, the dicot is a soybean plant, a tobacco plant, or a cotton plant.

Many of the plants intended to be transformed according to the disclosed invention are commercial crop plants. The commercial form of these plants may be the original plants, or their offspring which have inherited desired transgenes. Accordingly, plants further contemplated within the ambit of the present invention include any offspring of plants transformed with any of the permutations of the DNA construct which are noted in this application. Specifically, the offspring may be defined as an  $R_0$  transgenic plant. Other progeny of the transformed plant are also included within the scope of the present invention, including any progeny plant of any generation of the transformed plant, wherein the progeny plant has inherited the DNA construct from any  $R_0$  plant.

Upon transformation with a specific DNA construct, the nucleic acid or polynucleotide segments of the construct may be incorporated in various portions into a chromosome of the transformant. Therefore, in another embodiment, the present invention encompasses any transgenic plant or plant cell prepared by the use of a DNA construct disclosed herein. Such a plant or cell encompassed by the present invention includes those prepared by a process which has the following steps: (1) obtaining a DNA

construct including a Dipteran-inactive Cry2A *B. thuringiensis*  $\delta$ -endotoxin coding region positioned in frame and under the control of a promoter operable in the plant, and a plastid transit peptide coding region positioned upstream of the Cry2A *B. thuringiensis*  $\delta$ -endotoxin coding region and downstream of the promoter; and (2) transforming the plant with the obtained DNA construct, so that the plant expresses the Cry2A *B. thuringiensis*  $\delta$ -endotoxin. The plant may also have been transformed so that it further incorporates into its genome and expresses other Cry  $\delta$ -endotoxins.

In a related aspect, the present invention also encompasses a seed produced by the transformed plant, a progeny from such seed, and a seed produced by the progeny of the original transgenic plant, produced in accordance with the above process. Such progeny and seeds will have a Dipteran-inactive *B. thuringiensis*  $\delta$ -endotoxin transgene stably incorporated into its genome, and such progeny plants will inherit the traits afforded by the introduction of a stable transgene in Mendelian fashion. All such transgenic plants having incorporated into their genome transgenic DNA segments encoding any DNA construct disclosed herein, particularly those disclosed in sections 2.1 and 2.2 are aspects of this invention.

Recombinant plants, cells, seeds, and other tissues could also be produced in which only the mitochondrial or chloroplast DNA has been altered to incorporate the molecules envisioned in this application. Promoters which function in chloroplasts have been known in the art (Hanley-Bowden et al., Trends in Biochemical Sciences 12:67-70, 1987). Methods and compositions for obtaining cells containing chloroplasts into which heterologous DNA has been inserted has been described by Daniell et al., U.S. Pat. No. 5,693,507 (1997).

## 2.6 Plant Transformation Methods

### 2.6.1 Method of Expressing a Cry2A $\delta$ -Endotoxin in a Plant

In another preferred embodiment, the present invention provides a method for expressing Dipteran-inactive Cry2A *B. thuringiensis*  $\delta$ -endotoxins at high levels in transgenic plants. The disclosed methods may exploit any of the DNA constructs disclosed in sections 2.1 and 2.2 above, as well as any of the transformation vectors disclosed, for example, in section 2.3 above. The contemplated methods enable Cry2A  $\delta$ -endotoxins, an alternative to Cry1A *B. thuringiensis*  $\delta$ -endotoxins for the control of several insect pests, to be expressed in plants without negatively affecting the recovery of agronomic qualities of transgenic plants. The invention described herein also enables expression of Cry2A  $\delta$ -endotoxins at levels up to 25 times higher than that achieved by current methods.

The method described here thus enables plants expressing Cry2A to be used as either an alternative or supplement to plants expressing Cry1A-type *B. thuringiensis*  $\delta$ -endotoxins for both control and resistance management of key insect pests, including *Ostrina* sp., *Diatraea* sp., *Helicoverpa* sp., *Spodoptera* sp. in *Zea mays*; *Heliothis virescens*, *Helicoverpa* sp., *Pectinophora* sp. in *Gossypium hirsutum*; and *Anticarsia* sp., *Pseudoplusia* sp., *Epinotia* sp. in *Glycine max*. It is also contemplated that the

methods described may be used to dramatically increase expression of *B. thuringiensis*  $\delta$ -endotoxins including and related to Cry2A, thus increasing its effectiveness against target pests and decreasing the likelihood of evolved resistance to these proteins. In one embodiment of the present invention, the Cry2Ab  $\delta$ -endotoxin is expressed. Target pests of this protein and their common hosts are shown below  
 5 in Table 1.

Table 1  
 Cry2Ab Target Pests and Common Plant Hosts of those Pests

Pests	Hosts	Reference
<i>Ostrina nubilalis</i>	<i>Zea mays</i>	Donovan
<i>Diatraea grandiosella</i>	<i>Gossypium hirsutum</i>	U. S. Patent 5,338,544
<i>Helicoverpa zea</i>	<i>Glycine max</i>	
<i>Heliothis virescens</i>		
<i>Pectinophora gossypiella</i>		
<i>Anticarsia gemmatilis</i>		
<i>Pseudoplusia includens</i>		
<i>Epinotia aporema</i>		

The method of expressing a Cry2A *B. thuringiensis*  $\delta$ -endotoxin in a plant disclosed herein  
 10 includes the steps of: (1) obtaining nucleic acid segment comprising a promoter operably linked to a first polynucleotide sequence encoding a plastid transit peptide, and a second polynucleotide sequence, encoding a Cry2A *B. thuringiensis*  $\delta$ -endotoxin lacking Dipteran activity, to yield a fusion protein comprised of an amino-terminal plastid transit peptide and a Cry2A *B. thuringiensis*  $\delta$ -endotoxin lacking Dipteran activity; and (2) transforming the plant with the DNA construct of step 1 so that the plant  
 15 expresses the protein fusion. In a preferred embodiment, the nucleic acid segment employed in step (1) of this method is structured so that the 5' end of the second polynucleotide sequence is operably linked in the same translational reading frame to the 3' end of the first polynucleotide sequence.

The plant or plant cell transformed by the method disclosed herein may be either a monocotyledonous plant or a dicotyledonous plant. Where the plant is a monocotyledonous plant, it may  
 20 be any one of a variety of species. Preferred monocotyledonous species encompassed by the present invention may include maize, rice, wheat, barley, oats, rye, millet, sorghum, sugarcane, asparagus, turfgrass, or any of a number of other grains or cereal plants. In preferred embodiments, the monocot is a maize plant.

The present invention also contemplates a process by which a variety of dicotyledonous plants or  
 25 plant cells are transformed. Such dicotyledonous plants may include plants such as cotton, soybean, tomato, potato, citrus, tobacco, sugar beet, alfalfa, fava bean, pea, bean, apple, cherry, pear, strawberry,

raspberry, or any other legume, tuber, or fruit plant. In preferred embodiments, the dicot is a soybean plant, a tobacco plant or cell, or a cotton plant or cell.

### 2.6.2 Method of expressing a Cry2Ab $\delta$ -endotoxin in a Progeny Plant

As noted with regard to other embodiments disclosed in the present invention, many of the plants intended to be transformed according to the disclosed invention are commercial crop plants. The commercial form of these plants may be the original plants, or their offspring which have inherited desired transgenes. Accordingly, the inventors further contemplate that the method disclosed herein includes a method of producing a transgenic progeny plant or progeny plant cell. The method of producing such progeny includes: The method of expressing a Cry2A *B. thuringiensis*  $\delta$ -endotoxin in a plant disclosed herein includes the steps of: (1) obtaining nucleic acid segment comprising a promoter operably linked to a first polynucleotide sequence encoding a plastid transit peptide, and a second polynucleotide sequence, encoding a Cry2A *B. thuringiensis*  $\delta$ -endotoxin lacking Dipteran activity, to yield a fusion protein comprised of an amino-terminal plastid transit peptide and a Cry2A *B. thuringiensis*  $\delta$ -endotoxin lacking Dipteran activity; (2) obtaining a second plant; and (3) crossing the first and second plants to obtain a crossed transgenic progeny plant or plant cell which has inherited the nucleic acid segments from the first plant. The present invention specifically encompasses the progeny, progeny plant or seed from any of the monocotyledonous or dicotyledonous plants, including those noted in sections 2.5 and 2.6.1 above.

### 2.6.3 Method of Co-Expressing Cry2Ab and other Cry *B. thuringiensis* $\delta$ -endotoxins in a Plant and a Progeny Plant

In another preferred embodiment, the method of expressing the Dipteran-inactive Cry2A *B. thuringiensis*  $\delta$ -endotoxin disclosed herein includes co-expression of the disclosed DNA construct in any of its various embodiments, along with a Cry1 *B. thuringiensis*  $\delta$ -endotoxin. The method of expressing these Cry *B. thuringiensis*  $\delta$ -endotoxins together is expected to achieve increased insecticidal properties in the transformed plant through increased expression and decreased development of insect resistance - all of which are desired results not present in existing technologies. This co-expression may be in the original transformant, or in any number of generations of progeny of the original transformant which have inherited the genes to co-express the proteins encoded for by any of the DNA constructs disclosed herein.

### 3.0 Brief Description of the Drawings

The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

FIG. 1. Schematic illustration of elements of monocot plant *cry2Ab* expression vectors pMON30464, pMON30463, and pMON26800.

FIG. 2. Schematic illustration of elements of dicot *cry2Ab* expression vectors pMON33830, pMON33827, pMON33828, and pMON33829.

FIG. 3. Schematic illustration of elements of dicot *cry2Aa* expression vectors pMON33803, pMON33812, pMON33811, and pMON33806.

FIG. 4. Plasmid designated pMON30464.

FIG. 5. Plasmid designated pMON33827.

FIG. 6. Plasmid designated pMON33828.

FIG. 7. Plasmid designated pMON33829.

#### 4.0 DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

The following detailed description of the invention is provided to aid those skilled in the art in practicing the present invention. Even so, the following detailed description should not be construed to unduly limit the present invention as modifications and variations in the embodiments discussed herein may be made by those of ordinary skill in the art without departing from the spirit or scope of the present inventive discovery.

#### 4.1 IDENTIFICATION OF SEQUENCES

SEQ ID NO:1. Nucleic acid sequence of a *cry2Ab* gene.

SEQ ID NO:2. Amino acid sequence of a *Cry2Ab B. thuringiensis*  $\delta$ -endotoxin.

SEQ ID NO:3. Nucleic acid sequence of a zmSSU plastid transit peptide.

SEQ ID NO:4. Amino acid sequence of a zmSSU plastid transit peptide.

SEQ ID NO:5. Nucleic acid sequence of a plastid transit peptide 1 (PTP1).

SEQ ID NO:6. Amino acid sequence of a PTP1.

SEQ ID NO:7. Nucleic acid sequence of a plastid transit peptide 1 $\Delta$  (PTP1 $\Delta$ ).

SEQ ID NO:8. Amino acid sequence of a PTP1 $\Delta$ .

SEQ ID NO:9. Nucleic acid sequence of a plastid transit peptide 2 (PTP2).

SEQ ID NO:10. Amino acid sequence of a PTP2.

SEQ ID NO:11. Nucleic acid sequence of a *cry2Aa* gene.

SEQ ID NO:12. Amino sequence of a *Cry2Aa* polypeptide.

SEQ ID NO:13. pMON33827.

SEQ ID NO:14. pMON33828.

SEQ ID NO:15. pMON33829.

SEQ ID NO:16. pMON30464.

SEQ ID NO: 17. *Bacillus thuringiensis cry2Ab* gene sequence, UWGCG accession number M23724 (Widner and Whiteley).



SEQ ID NO:18.

*Bacillus thuringiensis* cry2Ab amino acid sequence  
translated from SEQ ID NO:17.

#### 4.2 DEFINITIONS

The following words and phrases herein have the meanings as set forth below.

5 **Biological functional equivalents.** As used herein such equivalents with respect to the insecticidal proteins of the present invention are peptides, polypeptides and proteins that contain a sequence or moiety exhibiting sequence similarity to the novel peptides of the present invention, such as Cry2Ab, and which exhibit the same or similar functional properties as that of the polypeptides disclosed herein, including insecticidal activity. Biological equivalents also include peptides, polypeptides and  
10 proteins that react with, *i.e.* specifically bind to antibodies raised against Cry2Ab and that exhibit the same or similar insecticidal activity, including both monoclonal and polyclonal antibodies.

**Chloroplast or plastid localized,** as used herein, refers to a biological molecule, either polynucleotide or polypeptide, which is positioned within the chloroplast or plastid such that the molecule is isolated from the cellular cytoplasmic milieu, and functions within the chloroplast or plastid cytoplasm  
15 to provide the effects claimed in the instant invention. Localization of a biological molecule to the chloroplast or plastid can occur, with reference to polynucleotides, by artificial mechanical means such as electroporation, mechanical microinjection, or by polynucleotide coated microprojectile bombardment, or with reference to polypeptides, by secretory or import means wherein a natural, synthetic, or heterologous plastid or chloroplast targeting peptide sequence is used which functions to target, insert, assist, or  
20 localize a linked polypeptide into a chloroplast or plastid.

**Combating or Controlling Insect Damage** in an agricultural context refers to reduction in damage to a crop caused by infection by an insect pest. More generally, this phrase refers to reduction in the adverse effects caused by the presence of an undesired insect in any particular location.

**Event** refers to a transgenic plant derived from the insertion of foreign DNA into one or more  
25 unique sites in the nuclear genomic DNA.

**Expression:** The combination of intracellular processes, including transcription, translation, and other intracellular protein and RNA processing and stabilization functions, undergone by a coding DNA molecule such as a structural gene to produce a polypeptide.

**Insecticidal polypeptide** refers to a polypeptide having insecticidal properties, *e.g.*, a polypeptide  
30 which inhibits the growth, development, viability or fecundity of target insect pests.

**Operably Linked:** Nucleic acid coding segments connected in frame so that the properties of one influence the expression of the other.

**Plant-Expressible Coding Regions:** Coding regions which are expressible *in planta* because they contain typical plant regulatory elements to facilitate the expression of the gene of interest.

35 **Plastid Transit Peptide:** Any amino acid sequence useful in targeting or localizing a linked amino acid, such as a protein fusion, to a subcellular compartment or organelle such as a plastid.

**Progeny:** "Progeny" includes any offspring or descendant of the transgenic plant, or any subsequent plant which has the transformant in its lineage. Progeny is not limited to one generation, but rather encompasses the transformant's descendants so long as they contain or express the transgene. Seeds containing transgenic embryos as well as seeds from the transgenic plants and their offspring or descendants are also important parts of the invention.

**Promoter:** A recognition site on a DNA sequence or group of DNA sequences that provide an expression control element for a structural gene and to which RNA polymerase specifically binds and initiates RNA synthesis (transcription) of that gene.

**R<sub>0</sub>** is the primary regenerant plant derived from transformation of plant tissue or cells in culture. Subsequent progeny or generations derived from the R<sub>0</sub> are referred to as R<sub>1</sub> (first generation), R<sub>2</sub> (second generation), *etc.*

**Regeneration:** The process of growing a plant from a plant cell (*e.g.*, plant protoplast or explant).

**Stably maintained within a plant plastid or chloroplast** refers to the introduction by electroporation, transformation, transduction, or micelle or liposome-like fusion of a polynucleotide or nucleic acid into a chloroplast or plastid in such a way that the nucleic acid remains within the recipient chloroplast or plastid and within all subsequent progeny of the recipient chloroplast or plastid either by incorporation by recombination into the chloroplast or plastid genome, or as an autonomously replicating covalently closed circular replicon residing within the chloroplast or plastid by means of growth of any plant, plant cell, or plant tissue containing such transformed chloroplast or plastid and while in the presence of a chemical or compound which requires one or more genes present on and expressed from the replicon in order to ensure the survival of the transformed plastid or chloroplast and their progeny plastids or chloroplasts within the plant, plant cell, or plant tissue.

**Structural Coding Sequence** refers to a DNA sequence that encodes a peptide, polypeptide, or protein that is made by a cell following transcription of the structural coding sequence to messenger RNA (mRNA), followed by translation of the mRNA to the desired peptide, polypeptide, or protein product.

**Structural gene:** A gene that is expressed to produce a polypeptide.

**Substantial homology:** As this term is used herein, it refers to nucleic acid or polypeptide sequences which are about 86% homologous, to about 90% homologous, to about 95% homologous, to about 99% homologous. More specifically, the inventors envision substantial homologues to be about 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, and 99 percent homologous to the referent nucleic acid sequence of polypeptide.

**Substantial temporal or spatial regulation** refers to the expression of a gene within a plant or plant tissue from a plant operable promoter. With reference to temporal regulation, a promoter may be regulated for expression only during specific times during plant cell or tissue or even whole plant growth and development. A promoter which is actively expressing one or more genes only during seed

germination would be one example of temporal regulation. Other examples could include promoters which are actively expressing one or more genes only during times when the plant, plant cell or plant tissue is exposed to certain light intensities or during total darkness. Substantial temporal regulation refers to a promoter which is actively expressed at a certain time but which may or may not be completely suppressed at other times, such that expression may still be detected by monitoring for the presence of some indicator such as an enzyme produced from a coding sequence linked to such promoter, or as measured by the increase or decrease in some gene product such as an mRNA produced at various times throughout plant growth, differentiation, and development and/or in response to various environmental stimuli. Substantial spatial regulation refers to the expression of a gene linked to a promoter from which expression proceeds only during growth and development of certain cells or tissues within a plant. For example, a tapetal promoter would only be expected to be expressed during flower growth and development. Similarly, a root specific or root enhanced promoter would only be expected to be expressed from within root cells or root tissues. Substantially spatially regulated also refers to the level of expression from a particular tissue specific promoter in that particular tissue and as related to levels of expression from that or a similar promoter in other tissues, wherein expression may also be detected in tissues other than the particular tissue in which the promoter expression is preferred, but at significantly lower expression levels as measured by the production of an enzyme produced from a coding sequence linked to the promoter or by the appearance of some detectable gene product. Promoters can also be both substantially temporally and substantially spatially regulated together and simultaneously in a coordinately regulated manner.

**Synthetic gene:** Synthetic genes encoding the *B. thuringiensis*  $\delta$ -endotoxins of the present invention are those prepared in a manner involving any sort of genetic isolation or manipulation. This includes isolation of the gene from its naturally occurring state, manipulation of the gene as by codon modification (as described herein), or site-specific mutagenesis (as described herein), truncation of the gene or any other manipulative or isolative method.

**Terminator:** The 3' end transcription termination and polyadenylation sequence.

**Transformation:** A process of introducing an exogenous DNA sequence (e.g., a vector, or a recombinant DNA molecule) into a cell or protoplast in which that exogenous DNA is incorporated into a chromosome or is capable of autonomous replication.

**Transformed cell:** A cell which has been altered by the introduction of one or more exogenous DNA molecules into that cell.

**Transgene:** A gene construct or DNA segment comprising a gene which is desired to be expressed in the recipient cell, tissue or organism. This may include an entire plasmid, or other vector, or may simply include the functional coding section, region, domain, or segment of the transferred DNA construct.

**Transgenic cell:** Any cell derived or regenerated from a transformed cell or derived from a transgenic cell. Exemplary transgenic cells include plant calli derived from a transformed plant cell and particular cells such as leaf, root, stem, *e.g.*, somatic cells, or reproductive (germ) cells obtained from a transgenic plant.

5       **Transgenic event:** A plant or progeny thereof derived from the insertion of foreign DNA into the nuclear genome of a plant cell or protoplast.

**Transgenic plant:** A plant or progeny thereof which has been genetically modified to contain and express heterologous DNA sequences as proteins. As specifically exemplified herein, a transgenic soybean plant is genetically modified to contain and express at least one heterologous DNA sequence  
10 operably linked to and under the regulatory control of transcriptional control sequences which function in plant cells or tissue or in whole plants. A transgenic plant may also be referred to as a transformed plant. A transgenic plant also refers to progeny of the initial transgenic plant where those progeny contain and are capable of expressing the heterologous coding sequence under the regulatory control of the plant-expressible transcription control sequences described herein.

15       **Vector:** A DNA molecule capable of replication in a host cell and/or to which another DNA segment can be operatively linked so as to bring about replication of the linked segment. A plasmid is an exemplary vector.

#### 4.3       SYNTHESIS AND ISOLATION OF A NUCLEIC ACID SEGMENT ENCODING A *B. THURINGIENSIS* $\delta$ -ENDOTOXIN AND PLASTID TARGETING SEQUENCES

20       The present invention discloses novel DNA constructs comprising polynucleotide sequences encoding *B. thuringiensis*  $\delta$ -endotoxins, as well as plastid targeting sequences. Methods for the construction and expression of synthetic *B. thuringiensis* genes in plants are well known by those of skill in the art and are described in detail in U. S. Patent 5,500,365. The present invention contemplates the use of Cry2A *B. thuringiensis* genes in the transformation of both monocotyledonous and dicotyledonous  
25 plants. To potentiate the expression of these genes, the present invention provides DNA constructs comprising polynucleotide segments encoding plastid targeting peptides positioned upstream of the polynucleotide sequences encoding the desired *B. thuringiensis*  $\delta$ -endotoxins. In particular, sequences encoding *B. thuringiensis*  $\delta$ -endotoxins lacking substantial Dipteran species inhibitory activity are contemplated.

#### 30       4.4       Probes and Primers

      In one aspect, nucleotide sequence information provided by the invention allows for the preparation of relatively short DNA sequences having the ability to specifically hybridize to gene sequences of the selected polynucleotides disclosed herein. In these aspects, nucleic acid probes of an appropriate length are prepared based on a consideration of selected polypeptide sequences encoding  
35 Cry2A  $\delta$ -endotoxin polypeptides, *e.g.*, a sequence such as that shown in SEQ ID NO:1. These nucleic

acid probes may also be prepared based on a consideration of selected polynucleotide sequences encoding a plastid targeting peptide, such as those shown in SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, and SEQ ID NO:9. The ability of such nucleic acid probes to specifically hybridize to a gene sequence encoding a  $\delta$ -endotoxin polypeptide or a plastid targeting peptide sequence lends to them particular utility in a variety of embodiments. Most importantly, the probes may be used in a variety of assays for detecting the presence of complementary sequences in a given sample.

In certain embodiments, it is advantageous to use oligonucleotide primers. The sequence of such primers is designed using a polynucleotide of the present invention for use in detecting, amplifying or mutating a defined segment of a crystal protein gene from *B. thuringiensis* using PCR<sup>TM</sup> technology. The process may also be used to detect, amplify or mutate a defined segment of the polynucleotide encoding a plastid targeting peptide. Segments of genes related to the polynucleotides encoding the  $\delta$ -endotoxin polypeptides and plastid targeting peptides of the present invention may also be amplified by PCR<sup>TM</sup> using such primers.

To provide certain of the advantages in accordance with the present invention, a preferred nucleic acid sequence employed for hybridization studies or assays includes sequences that are complementary to at least a 14 to 30 or so long nucleotide stretch of a polynucleotide sequence encoding a crystal protein, such as that shown in SEQ ID NO:1, or sequences that are complementary to at least a 14 to 30 or so long nucleotide stretch of a sequence encoding a plastid targeting peptide, such as those shown in SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, and SEQ ID NO:9.

A size of at least 14 nucleotides in length helps to ensure that the fragment will be of sufficient length to form a duplex molecule that is both stable and selective. Molecules having complementary sequences over segments greater than 14 bases in length are generally preferred. In order to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained, one will generally prefer to design nucleic acid molecules having gene-complementary sequences of 14 to 20 nucleotides, or even longer where desired. Such fragments may be readily prepared by, for example, directly synthesizing the fragment by chemical means, by application of nucleic acid reproduction technology, such as the PCR<sup>TM</sup> technology of U. S. Patents 4,683,195, and 4,683,202 (each specifically incorporated herein by reference), or by excising selected DNA fragments from recombinant plasmids containing appropriate inserts and suitable restriction sites.

#### 4.5 Expression Vectors

The present invention also contemplates an expression vector comprising a polynucleotide of the present invention. Thus, in one embodiment an expression vector is an isolated and purified DNA molecule comprising a promoter operatively linked to a coding region that encodes a polypeptide of the present invention, which coding region is operatively linked to a transcription-terminating region, whereby the promoter drives the transcription of the coding region. The coding region may include a

segment encoding a *B. thuringiensis*  $\delta$ -endotoxin and a segment encoding a plastid target peptide. The DNA molecule comprising the expression vector may also contain a functional intron

As used herein, the terms "operatively linked" or "operably linked" mean that a promoter is connected to a coding region in such a way that the transcription of that coding region is controlled and regulated by that promoter. Means for operatively linking a promoter to a coding region to regulate both upstream and downstream are well known in the art.

Preferred plant transformation vectors include those derived from a Ti plasmid of *Agrobacterium tumefaciens*, as well as those disclosed, e.g., by Herrera-Estrella (1983), Bevan (1983), Klee (1985) and Eur. Pat Appl. No. EP 0120516 (each specifically incorporated herein by reference).

Promoters that function in bacteria are well known in the art. Exemplary and preferred promoters for the *B. thuringiensis* crystal proteins include the *sigA*, *sigE*, and *sigK* gene promoters. Alternatively, native, mutagenized, heterologous, or recombinant crystal protein-encoding gene promoters themselves can be used.

Where an expression vector of the present invention is to be used to transform a plant, a promoter is selected that has the ability to drive expression in that particular species of plant. Promoters that function in different plant species are also well known in the art. Promoters useful in expressing the polypeptide in plants are those which are inducible, viral, synthetic, or constitutive as described (Odell *et al.*, 1985), and/or temporally regulated, spatially regulated, and spatio-temporally regulated. Preferred promoters include the enhanced CaMV35S promoters, and the FMV35S promoter.

#### 4.5.1 Vectors with Plastid Targeting Peptide-Encoding Segments

In accordance with the present invention, expression vectors designed to specifically potentiate the expression of the polypeptide in the transformed plant may include certain regions encoding plastid targeting peptides (PTP). These regions allow for the cellular processes involved in transcription, translation and expression of the encoded protein to be fully exploited when associated with certain *B. thuringiensis*  $\delta$ -endotoxins. Such plastid targeting peptides function in a variety of ways, such as for example, by transferring the expressed protein to the cell structure in which it most effectively operates, or by transferring the expressed protein to areas of the cell in which cellular processes necessary for expression are concentrated.

The use of PTPs may also increase the frequency of recovery of morphologically normal plants, and the frequency at which transgenic plants may be recovered. Given that commercially viable expression of both Cry1A and Cry3A-type *B. thuringiensis*  $\delta$ -endotoxins have been achieved by expression of forms of the proteins that remain localized in the cytosol (*i.e.* non-targeted forms), expression of non-targeted forms of both Cry2Aa and Cry2Ab were also initially attempted in transgenic cotton, tobacco, and corn.

In corn, non-targeted Cry2Ab expression transformation vectors yield relatively few transgenic events (*i.e.* independent insertion events into the corn genome) with Cry2Ab expression levels sufficient

for commercially acceptable insect control. Moreover, many of the corn transformants expressing non-targeted Cry2Ab exhibited obvious growth defects such as severe reduction in stature (stunting) or severe yellowing of the leaves (chlorosis) that rendered the plants commercially unacceptable. Expression levels of non-targeted Cry2Ab in corn were no higher than approximately 15 ppm, a level minimally required for Cry2Ab-mediated control of European corn borer (ECB).

Although studies involving expression of plastid targeted Cry1A-type *B. thuringiensis*  $\delta$ -endotoxins in transgenic plants have been described (Wong *et al.*, 1992), targeting of the non-homologous Cry2A or Cry2A proteins has not previously been described. One report of plastid targeted Cry1Ac expression indicated that such targeting results in little or no increase in Cry1Ac expression (U.S. Patent No. 5,500,365). Another report indicated that an increase in expression of a plastid targeted form of Cry1Ac required the inclusion of a new 5' untranslated leader sequence (Wong *et al.*, 1992) and that the effect of the leader and targeting sequences on expression was highly dependent on the coding sequence of the structural gene. Wong *et al.* concluded that inclusion of both the leader sequence and plastid transit peptide increased Cry1Ac expression 18-fold, but the same sequences increased  $\beta$ -glucuronidase expression only 6-fold. Finally, none of the previous reports predicted that plastid targeting would result in increased recovery of morphologically normal *B. thuringiensis* expressing plants.

The present invention discloses that transgenic corn plants expressing Dipteran inactive Cry2A  $\delta$ -endotoxins, such as Cry2Ab, at levels up to 10-fold higher than required for ECB control were recovered at significantly higher frequencies when a plastid targeted form of the Cry2A was used. In the case of Cry2Ab, elevated expression is critical in obtaining transgenic corn with ECB control since the  $LC_{50}$  of Cry2Ab against ECB is significantly higher than the  $LC_{50}$  ECB of the Cry1Ab *B. thuringiensis* currently used to control ECB in transgenic corn (U. S. Patent 5,338,544, 1994; MacIntosh *et al.*, 1990; Armstrong *et al.*, 1995).

Increased expression is also especially valuable in that it provides additional protection against development of resistance via a high dose strategy (McGaughey and Whalon, 1993; Roush, 1994). High level expression is even further desirable as it provides sustained insect protection in instances where insecticidal gene expression decreases due to environmental conditions. Additionally and unexpectedly, corn plants transformed with plastid targeted Cry2Ab expression vectors exhibited normal growth and development.

A significant distinction between targeted and non-targeted (cytosolic) expression of Cry2Ab was the dramatic increase in levels of Cry2Ab protein in plants transformed with the plastid targeted Cry2Ab expression vector relative to plants transformed with the cytosolic Cry2Ab vector. This result was very unexpected. Also, in contrast to the teachings of previous work, the invention disclosed herein reveals

that enhanced recovery of phenotypically normal transgenic plants can be achieved using the disclosed methods of plastid targeted expression.

An example of a plastid targeting peptide (PTP) is a chloroplast targeting peptide. Chloroplast targeting peptides have been found particularly useful in the glyphosate resistant selectable marker system. In this system, plants transformed to express a protein conferring glyphosate resistance are transformed with a PTP that targets the peptide to the cell's chloroplasts. Glyphosate inhibits the shikimic acid pathway which leads to the biosynthesis of aromatic compounds including amino acids and vitamins. Specifically, glyphosate inhibits the conversion of phosphoenolpyruvic acid and 3-phosphoshikimic acid to 5-enolpyruvyl-3-phosphoshikimic acid by inhibiting the enzyme 5-enolpyruvyl-3-phosphoshikimic acid synthase (EPSP synthase or EPSPS). Supplemental EPSPS, conferred via insertion of a transgene encoding this enzyme, allows the cell to resist the effects of the glyphosate. Thus, as the herbicide glyphosate functions to kill the cell by interrupting aromatic amino acid biosynthesis, particularly in the cell's chloroplast, the PTP allows increased resistance to the herbicide by concentrating what glyphosate resistance enzyme the cell expresses in the chloroplast, *i.e.* in the target organelle of the cell. Exemplary herbicide resistance enzymes include EPSPS as noted above, glyphosate oxido-reductase (GOX) and the *aroA* gene (see U.S. Patent No. 4,535,060, specifically incorporated herein by reference in its entirety).

PTPs can target proteins to chloroplasts and other plastids. For example, the target organelle may be the amyloplast. Preferred PTPs of the present invention include those targeting both chloroplasts as well as other plastids. Specific examples of preferred PTPs include the maize RUBISCO SSU protein PTP, and functionally related peptides such as PTP1, PTPA, and PTP2. These PTPs are exemplified by the polypeptides shown in SEQ ID NO:4, SEQ ID NO:6, SEQ ID NO:8, and SEQ ID NO:10. Polynucleotide sequences encoding for these polypeptides are shown in SEQ ID NO:3, SEQ ID NO:5, SEQ ID NO:7, and SEQ ID NO:9.

Recombinant plants, cells, seeds, and other plant tissues could also be produced in which only the mitochondrial or chloroplast DNA has been altered to incorporate the molecules envisioned in this application. Promoters which function in chloroplasts have been known in the art (Hanley-Bowden et al., Trends in Biochemical Sciences 12:67-70, 1987). Methods and compositions for obtaining cells containing chloroplasts into which heterologous DNA has been inserted has been described by Daniell et al., U.S. Pat. No. 5,693,507 (1997). McBride et al. (WO 95/24492) disclose localization and expression of genes encoding Cry1A  $\delta$ -endotoxin protein in tobacco plant chloroplast genomes. As disclosed herein, localization of Cry2Aa to the chloroplast or plastid results in decreased levels of expression as measured by accumulation of Cry2Aa  $\delta$ -endotoxin, which is in contrast to the improved expression of chloroplast or plastid localized Cry2Ab  $\delta$ -endotoxin.



#### 4.5.2 Use of Promoters in Expression Vectors

The expression of a gene which exists in double-stranded DNA form involves transcription of messenger RNA (mRNA) from the coding strand of the DNA by an RNA polymerase enzyme, and the subsequent processing of the mRNA primary transcript inside the nucleus. Transcription of DNA into mRNA is regulated by a region of DNA usually referred to as the "promoter". The promoter region contains a sequence of bases that signals RNA polymerase to associate with the DNA and to initiate the transcription of mRNA using one of the DNA strands as a template to make a corresponding strand of RNA. The particular promoter selected should be capable of causing sufficient expression of the enzyme coding sequence to result in the production of an effective insecticidal amount of the *B. thuringiensis* protein.

The 3' non-translated region of the chimeric plant genes of the present invention also contains a polyadenylation signal which functions in plants to cause the addition of adenylate nucleotides to the 3' end of the RNA. Examples of preferred 3' regions are (1) the 3' transcribed, non-translated regions containing the polyadenylation signal of *Agrobacterium* tumor-inducing (Ti) plasmid genes, such as the nopaline synthase (NOS) gene and (2) the 3' ends of plant genes such as the pea ssRUBISCO E9 gene (Fischhoff *et al.*, 1987).

A promoter is selected for its ability to direct the transformed plant cell's or transgenic plant's transcriptional activity to the coding region, to ensure sufficient expression of the enzyme coding sequence to result in the production of insecticidal amounts of the *B. thuringiensis* protein. Structural genes can be driven by a variety of promoters in plant tissues. Promoters can be near-constitutive (*i.e.* they drive transcription of the transgene in all tissue), such as the CaMV35S promoter, or tissue-specific or developmentally specific promoters affecting dicots or monocots. Where the promoter is a near-constitutive promoter such as CaMV35S or FMV35S, increases in polypeptide expression are found in a variety of transformed plant tissues and most plant organs (*e.g.*, callus, leaf, seed and root). Enhanced or duplicate versions of the CaMV35S and FMV35S promoters are particularly useful in the practice of this invention (Kay *et al.*, 1987; Rogers, U. S. Patent 5,378,619).

Those skilled in the art will recognize that there are a number of promoters which are active in plant cells, and have been described in the literature. Such promoters may be obtained from plants or plant viruses and include, but are not limited to, the nopaline synthase (NOS) and octopine synthase (OCS) promoters (which are carried on tumor-inducing plasmids of *A. tumefaciens*), the cauliflower mosaic virus (CaMV) 19S and 35S promoters, the light-inducible promoter from the small subunit of ribulose 1,5-bisphosphate carboxylase (ssRUBISCO, a very abundant plant polypeptide), the rice *Act1* promoter and the Figwort Mosaic Virus (FMV) 35S promoter. All of these promoters have been used to create various types of DNA constructs which have been expressed in plants (see *e.g.*, McElroy *et al.*, 1990, U. S. Patent 5,463,175).

In addition, it may also be preferred to bring about expression of the *B. thuringiensis*  $\delta$ -endotoxin in specific tissues of the plant by using plant integrating vectors containing a tissue-specific promoter. Specific target tissues may include the leaf, stem, root, tuber, seed, fruit, *etc.*, and the promoter chosen should have the desired tissue and developmental specificity. Therefore, promoter function should be optimized by selecting a promoter with the desired tissue expression capabilities and approximate promoter strength and selecting a transformant which produces the desired insecticidal activity in the target tissues. This selection approach from the pool of transformants is routinely employed in expression of heterologous structural genes in plants since there is variation between transformants containing the same heterologous gene due to the site of gene insertion within the plant genome (commonly referred to as "position effect"). In addition to promoters which are known to cause transcription (constitutive or tissue-specific) of DNA in plant cells, other promoters may be identified for use in the current invention by screening a plant cDNA library for genes which are selectively or preferably expressed in the target tissues and then determine the promoter regions.

An exemplary tissue-specific promoter is the lectin promoter, which is specific for seed tissue. The lectin protein in soybean seeds is encoded by a single gene (*Le1*) that is only expressed during seed maturation and accounts for about 2 to about 5% of total seed mRNA. The lectin gene and seed-specific promoter have been fully characterized and used to direct seed specific expression in transgenic tobacco plants (Vodkin *et al.*, 1983; Lindstrom *et al.*, 1990). An expression vector containing a coding region that encodes a polypeptide of interest can be engineered to be under control of the lectin promoter and that vector may be introduced into plants using, for example, a protoplast transformation method (Dhir *et al.*, 1991). The expression of the polypeptide would then be directed specifically to the seeds of the transgenic plant.

A transgenic plant of the present invention produced from a plant cell transformed with a tissue specific promoter can be crossed with a second transgenic plant developed from a plant cell transformed with a different tissue specific promoter to produce a hybrid transgenic plant that shows the effects of transformation in more than one specific tissue.

Other exemplary tissue-specific promoters are corn sucrose synthetase 1 (Yang *et al.*, 1990), corn alcohol dehydrogenase 1 (Vogel *et al.*, 1989), corn light harvesting complex (Simpson, 1986), corn heat shock protein (Odell *et al.*, 1985), *pca* small subunit RuBP carboxylase (Poulsen *et al.*, 1986; Cashmore *et al.*, 1983), Ti plasmid mannopine synthase (McBride and Summerfelt, 1989), Ti plasmid nopaline synthase (Langridge *et al.*, 1989), petunia chalcone isomerase (Van Tunen *et al.*, 1988), bean glycine rich protein 1 (Keller *et al.*, 1989), CaMV 35S transcript (Odell *et al.*, 1985) and Potato patatin (Wenzler *et al.*, 1989). Preferred promoters are the cauliflower mosaic virus (CaMV 35S) promoter and the S-E9 small subunit RuBP carboxylase promoter.

The promoters used in the DNA constructs of the present invention may be modified, if desired, to affect their control characteristics. For example, the CaMV35S promoter may be ligated to the portion

of the ssRUBISCO gene that represses the expression of ssRUBISCO in the absence of light, to create a promoter which is active in leaves but not in roots. The resulting chimeric promoter may be used as described herein. For purposes of this description, the phrase "CaMV35S" promoter thus includes variations of CaMV35S promoter, *e.g.*, promoters derived by means of ligation with operator regions, random or controlled mutagenesis, *etc.* Furthermore, the promoters may be altered to contain multiple "enhancer sequences" to assist in elevating gene expression. Examples of such enhancer sequences have been reported by Kay *et al.* (1987). Chloroplast or plastid specific promoters are known in the art (Daniell *et al.*, US Pat. No. 5,693,507; herein incorporated by reference), for example promoters obtainable from chloroplast genes, such as the *psbA* gene from spinach or pea, the *rbcL* and *atpB* promoter region from maize, and rRNA promoters. Any chloroplast or plastid operable promoter is within the scope of the present invention.

The RNA produced by a DNA construct of the present invention also contains a 5' non-translated leader sequence. This sequence can be derived from the promoter selected to express the gene, and can be specifically modified so as to increase translation of the mRNA. The 5' non-translated regions can also be obtained from viral RNAs, from suitable eukaryotic genes, or from a synthetic gene sequence. The present invention is not limited to constructs wherein the non-translated region is derived from the 5' non-translated sequence that accompanies the promoter sequence. As shown below, a plant gene leader sequence which is useful in the present invention is the petunia heat shock protein 70 (hsp70) leader (Winter *et al.*, 1988).

An exemplary embodiment of the invention involves the plastid targeting or plastid localization of the *B. thuringiensis* amino acid sequence. Plastid targeting sequences have been isolated from numerous nuclear encoded plant genes and have been shown to direct importation of cytoplasmically synthesized proteins into plastids (reviewed in Keegstra and Olsen, 1989). A variety of plastid targeting sequences, well known in the art, including but not limited to ADPGPP, EPSP synthase, or ssRUBISCO, may be utilized in practicing this invention. In alternative embodiments preferred, plastidic targeting sequences (peptide and nucleic acid) for monocotyledonous crops may consist of a genomic coding fragment containing an intron sequence as well as a duplicated proteolytic cleavage site in the encoded plastidic targeting sequences.

The most preferred nucleic acid sequence, referred to herein as zmSSU PTP (SEQ ID NO:3), consists of a genomic coding fragment containing an intron sequence as well as a duplicated proteolytic cleavage site in the encoded plastidic targeting sequences, was derived from plastid targeting sequence zmS1 (Russell *et al.*, 1993). Direct translational fusions of zmSSU PTP peptide sequence (SEQ ID NO:4) to the amino terminus of the sequences are useful in obtaining elevated levels of the polypeptide in transgenic maize. In-frame fusions of the zmSSU PTP nucleic acid sequence (SEQ ID NO:3) to the

*cry2Ab* gene (SEQ ID NO:1) can be effected by ligation of the *NcoI* site at the 3' (C-terminal encoding) end of the *zmSSU* PTP sequence with the 5' *NcoI* site (N-terminal encoding) of the *cry2Ab* sequence.

The preferred sequence for dicotyledonous crops referred to herein as PTP2 (SEQ ID NO:9), consists of a genomic coding fragment containing the chloroplast targeting peptide sequence from the EPSP synthase gene of *Arabidopsis thaliana* in which the transit peptide cleavage site of the pea ssRUBISCO PTP replaces the native EPSP synthase PTP cleavage site (Klee *et al.*, 1987).

As noted above, the 3' non-translated region of the chimeric plant genes of the present invention contains a polyadenylation signal which functions in plants to cause the addition of adenylate nucleotides to the 3' end of the RNA. Examples of preferred 3' regions are (1) the 3' transcribed, non-translated regions containing the polyadenylate signal of *Agrobacterium* tumor-inducing (Ti) plasmid genes, such as the nopaline synthase (NOS) gene and (2) plant genes such as the pea ssRUBISCO E9 gene (Fischhoff *et al.*, 1987).

#### 4.5.3 Use of Introns in Expression Vectors

For optimized expression in monocotyledonous plants, an intron may also be included in the DNA expression construct. Such an intron is typically placed near the 5'-end of the mRNA in untranslated sequence. This intron could be obtained from, but not limited to, a set of introns consisting of the maize Heat Shock Protein (HSP) 70 intron (U. S. Patent 5,424,412; 1995), the rice *Act1* intron (McElroy *et al.*, 1990), the *Adh* intron 1 (Callis *et al.*, 1987), or the sucrose synthase intron (Vasil *et al.*, 1989). As shown herein, the maize HSP70 intron is useful in the present invention.

#### 4.5.4 USE OF TERMINATORS IN EXPRESSION VECTORS

RNA polymerase transcribes a nuclear genome coding DNA sequence through a site where polyadenylation occurs. Typically, DNA sequences located a few hundred base pairs downstream of the polyadenylation site serve to terminate transcription. Those DNA sequences are referred to herein as transcription-termination regions. Those regions are required for efficient polyadenylation of transcribed messenger RNA (mRNA). For coding sequences introduced into a chloroplast or plastid, or into a chloroplast or plastid genome, mRNA transcription termination is similar to methods well known in the bacterial gene expression art. For example, either in a polycistronic or a monocistronic sequence, transcription can be terminated by stem and loop structures or structures similar to rho dependent sequences.

Constructs will typically include the gene of interest along with a 3' end DNA sequence that acts as a signal to terminate transcription and, in constructs intended for nuclear genome expression, allow for the polyadenylation of the resultant mRNA. The most preferred 3' elements are contemplated to be those from the nopaline synthase gene of *A. tumefaciens* (nos 3'end) (Bevan *et al.*, 1983), the terminator for the T7 transcript from the octopine synthase gene of *A. tumefaciens*, and the 3' end of the protease inhibitor i

or ii genes from potato or tomato. Regulatory elements such as TMV  $\Omega$  element (Gallie, *et al.*, 1989), may further be included where desired.

#### 4.5.5 Other Expression-Enhancing Elements

Another type of element which can regulate gene expression is the DNA sequence between the transcription initiation site and the start of the coding sequence, termed the untranslated leader sequence. The leader sequence can influence gene expression. Compilations of leader sequences have been made to predict optimum or sub-optimum sequences and generate "consensus" and preferred leader sequences (Joshi, 1987). Preferred leader sequences are contemplated to include those which comprise sequences predicted to direct optimum expression of the linked structural gene, *i.e.* to include a preferred consensus leader sequence which may increase or maintain mRNA stability and prevent inappropriate initiation of translation. The choice of such sequences will be known to those of skill in the art in light of the present disclosure. Sequences that are derived from genes that are highly expressed in plants, and in maize in particular, will be most preferred. One particularly useful leader may be the petunia HSP70 leader.

Transcription enhancers or duplications of enhancers could be used to increase expression. These enhancers often are found 5' to the start of transcription in a promoter that functions in eukaryotic cells, but can often be inserted in the forward or reverse orientation 5' or 3' to the coding sequence. Examples of enhancers include elements from the CaMV 35S promoter, octopine synthase genes (Ellis *et al.*, 1987), the rice actin gene, and promoter from non-plant eukaryotes (*e.g.*, yeast; Ma *et al.*, 1988).

#### 4.5.6 Multigene Vector Constructs and IRES

In certain embodiments of the invention, the use of internal ribosome binding sites (IRES) elements are used to create multigene, or polycistronic, messages. IRES elements are able to bypass the ribosome scanning model of 5' methylated Cap dependent translation and begin translation at internal sites (Pelletier and Sonenberg, 1988). IRES elements from two members of the picornavirus family (polio and encephalomyocarditis) have been described (Pelletier and Sonenberg, 1988), as well an IRES from a mammalian message (Macejak and Sarnow, 1991). IRES elements can be linked to heterologous open reading frames. Multiple open reading frames can be transcribed together, each separated by an IRES, creating polycistronic messages. By virtue of the IRES element, each open reading frame is accessible to ribosomes for efficient translation. Multiple genes can be efficiently expressed using a single promoter/enhancer to transcribe a single message.

Any heterologous open reading frame can be linked to IRES elements. This includes genes for secreted proteins, multi-subunit proteins, encoded by independent genes, intracellular or membrane-bound proteins and selectable markers. In this way, expression of several proteins can be simultaneously engineered into a cell with a single construct and a single selectable marker.

Constructs intended for expression from within a chloroplast or plastid utilizing chloroplast or plastid specific transcriptional and translational machinery can contain either mono- or polycistronic sequences.

#### 4.5.7 Construction of the Expression Vector

5 The choice of which expression vector and ultimately to which promoter a polypeptide coding region is operatively linked depends directly on the functional properties desired, e.g., the location and timing of protein expression, and the host cell to be transformed. These are well known limitations inherent in the art of constructing recombinant DNA molecules. However, a vector useful in practicing the present invention is capable of directing the expression of the polypeptide coding region to which it is  
10 operatively linked.

Typical vectors useful for expression of genes in higher plants are well known in the art and include vectors derived from the tumor-inducing (Ti) plasmid of *A. tumefaciens* described (Rogers *et al.*, 1987). However, several other plant integrating vector systems are known to function in plants including pCaMVN transfer control vector described (Fromm *et al.*, 1986). pCaMVN (available from  
15 Pharmacia, Piscataway, NJ) includes the CaMV35S promoter.

In preferred embodiments, the vector used to express the polypeptide includes a selection marker that is effective in a plant cell, preferably a drug resistance selection marker. One preferred drug resistance marker is the gene whose expression results in kanamycin resistance; i.e. the chimeric gene containing the nopaline synthase promoter, Tn5 neomycin phosphotransferase II (*npII*) and nopaline  
20 synthase 3' non-translated region described (Rogers *et al.*, 1988).

Means for preparing expression vectors are well known in the art. Expression (transformation) vectors used to transform plants and methods of making those vectors are described in U. S. Patents 4,971,908, 4,940,835, 4,769,061 and 4,757,011 (each of which is specifically incorporated herein by reference). Those vectors can be modified to include a coding sequence in accordance with the present  
25 invention.

A variety of methods have been developed to operatively link DNA to vectors *via* complementary cohesive termini or blunt ends. For instance, complementary homopolymer tracts can be added to the DNA segment to be inserted and to the vector DNA. The vector and DNA segment are then joined by hydrogen bonding between the complementary homopolymeric tails to form recombinant DNA  
30 molecules.

A coding region that encodes a polypeptide having the ability to confer insecticidal activity to a cell is preferably a polynucleotide encoding a *B. thuringiensis*  $\delta$ -endotoxin or a functional equivalent of such a polynucleotide. In accordance with such embodiments, a coding region comprising the DNA sequence of SEQ ID NO:1 is also preferred.

35 Specific *B. thuringiensis*  $\delta$ -endotoxin polypeptide-encoding genes that have been shown to successfully transform plants in conjunction with plastid targeting peptide-encoding genes, to express the

*B. thuringiensis*  $\delta$ -endotoxins at high levels are those genes comprised within the plasmid vectors. Preferred plasmids containing plastid targeting sequences include pMON30464, pMON33827, pMON33828, pMON33829. These plasmids are encoded for by the sequences shown in SEQ ID NO:16, SEQ ID NO:13, SEQ ID NO:14, SEQ ID NO:15. More preferably, plants may be successfully transformed with any vector containing expression cassettes comprising the nucleotide sequences of nucleotide 1781 to 5869 of SEQ ID NO:16, nucleotide 17 to 3182 of SEQ ID NO:13, nucleotide 17 to 3092 of SEQ ID NO:14 or nucleotide 17 to 3155 of SEQ ID NO:15.

The work described herein has identified methods of potentiating *in planta* expression of *B. thuringiensis*  $\delta$ -endotoxins, which confer resistance to insect pathogens when incorporated into the nuclear, plastid, or chloroplast genome of susceptible plants. U. S. Patent 5,500,365 (specifically incorporated herein by reference) describes a method for synthesizing plant genes to optimize the expression level of the protein for which the synthesized gene encodes. This method relates to the modification of the structural gene sequences of the exogenous transgene, to make them more "plant-like" and therefore more likely to be translated and expressed by the plant. A similar method for enhanced expression of transgenes, preferably in monocotyledonous plants, is disclosed in U. S. Patent 5,689,052 (specifically incorporated herein by reference). Agronomic, horticultural, ornamental, and other economically or commercially useful plants can be made in accordance with the methods described herein, to express *B. thuringiensis*  $\delta$ -endotoxins at levels high enough to confer resistance to insect pathogens.

Such plants may co-express the *B. thuringiensis*  $\delta$ -endotoxin polypeptide along with other antifungal, antibacterial, or antiviral pathogenesis-related peptides, polypeptides, or proteins; insecticidal proteins; proteins conferring herbicide resistance; and proteins involved in improving the quality of plant products or agronomic performance of plants. Simultaneous co-expression of multiple proteins in plants is advantageous in that it exploits more than one mode of action to control plant pathogenic damage. This can minimize the possibility of developing resistant pathogen strains, broaden the scope of resistance, and potentially result in a synergistic insecticidal effect, thereby enhancing plants ability to resist insect infestation (WO 92/17591).

Specifically contemplated for use in accordance with the present invention are vectors which include the *ocs* enhancer element. This element was first identified as a 16 bp palindromic enhancer from the octopine synthase (*ocs*) gene of *Agrobacterium* (Ellis *et al.*, 1987), and is present in at least 10 other promoters (Bouchez *et al.*, 1989). It is proposed that the use of an enhancer element, such as the *ocs* element and particularly multiple copies of the element, may be used to increase the level of transcription from adjacent promoters when applied in the context of monocot transformation.

It is contemplated that introduction of large DNA sequences comprising more than one gene may be desirable. Introduction of such sequences may be facilitated by use of bacterial or yeast artificial

chromosomes (BACs or YACs, respectively), or even plant artificial chromosomes. For example, the use of BACs for *Agrobacterium*-mediated transformation was disclosed by Hamilton *et al.* (1996).

Ultimately, the most desirable DNA segments for introduction into a monocot genome may be homologous genes or gene families which encode a desired trait (for example, increased yield), and which are introduced under the control of novel promoters or enhancers, *etc.*, or perhaps even homologous or tissue specific (e.g., root-collar/sheath-, whorl-, stalk-, earshank-, kernel- or leaf-specific) promoters or control elements. Indeed, it is envisioned that a particular use of the present invention may be the production of transformants comprising a transgene which is targeted in a tissue-specific manner. For example, insect resistant genes may be expressed specifically in the whorl and collar/sheath tissues which are targets for the first and second broods, respectively, of ECB. Likewise, genes encoding proteins with particular activity against rootworm may be targeted directly to root tissues.

Vectors for use in tissue-specific targeting of gene expression in transgenic plants typically will include tissue-specific promoters and also may include other tissue-specific control elements such as enhancer sequences. Promoters which direct specific or enhanced expression in certain plant tissues will be known to those of skill in the art in light of the present disclosure.

It also is contemplated that tissue specific expression may be functionally accomplished by introducing a constitutively expressed gene (all tissues) in combination with an antisense gene that is expressed only in those tissues where the gene product is not desired. For example, a gene coding for the crystal toxin protein from *B. thuringiensis* may be introduced such that it is expressed in all tissues using the 35S promoter from Cauliflower Mosaic Virus. Alternatively, a rice actin promoter or a histone promoter from a dicot or monocot species also could be used for constitutive expression of a gene. Furthermore, it is contemplated that promoters combining elements from more than one promoter may be useful. For example, U. S. Patent 5,491,288 discloses combining a Cauliflower Mosaic Virus promoter with a histone promoter. Therefore, expression of an antisense transcript of the Bt gene in a maize kernel, using for example a zein promoter, would prevent accumulation of the  $\delta$ -endotoxin in seed. Hence the protein encoded by the introduced gene would be present in all tissues except the kernel. It is specifically contemplated by the inventors that a similar strategy could be used with the instant invention to direct expression of a screenable or selectable marker in seed tissue.

Alternatively, one may wish to obtain novel tissue-specific promoter sequences for use in accordance with the present invention. To achieve this, one may first isolate cDNA clones from the tissue concerned and identify those clones which are expressed specifically in that tissue, for example, using Northern blotting. Ideally, one would like to identify a gene that is not present in a high copy number, but which gene product is relatively abundant in specific tissues. The promoter and control elements of corresponding genomic clones may this be localized using the techniques of molecular biology known to those of skill in the art.



It is contemplated that expression of some genes in transgenic plants will be desired only under specified conditions. For example, it is proposed that expression of certain genes that confer resistance to environmentally stress factors such as drought will be desired only under actual stress conditions. It further is contemplated that expression of such genes throughout a plants development may have detrimental effects. It is known that a large number of genes exist that respond to the environment. For example, expression of some genes such as *rbcS*, encoding the small subunit of ribulose biphosphate carboxylase, is regulated by light as mediated through phytochrome. Other genes are induced by secondary stimuli. For example, synthesis of abscisic acid (ABA) is induced by certain environmental factors, including but not limited to water stress. A number of genes have been shown to be induced by ABA (Skriver and Mundy, 1990). It also is expected that expression of genes conferring resistance to insect predation would be desired only under conditions of actual insect infestation. Therefore, for some desired traits, inducible expression of genes in transgenic plants will be desired.

It is proposed that, in some embodiments of the present invention, expression of a gene in a transgenic plant will be desired only in a certain time period during the development of the plant. Developmental timing frequently is correlated with tissue specific gene expression. For example expression of zein storage proteins is initiated in the endosperm about 15 days after pollination.

It also is contemplated that it may be useful to target DNA itself with a cell. For example, it may be useful to target introduced DNA to the nucleus as this may increase the frequency of transformation. Within the nucleus itself it would be useful to target a gene in order to achieve site specific integration. For example, it would be useful to have a gene introduced through transformation replace an existing gene in the cell.

#### 4.6 Identification and Isolation of Insecticidal *B. thuringiensis* $\delta$ -Endotoxins and Genes

It is contemplated that the method described in this invention could be used to obtain substantially improved expression of a number of novel *B. thuringiensis* endotoxins isolated as described below. Identification of new *Bacillus thuringiensis* strains encoding crystalline endotoxins with insecticidal activity has been described previously (Donovan *et al.*, 1992). Isolation of the *B. thuringiensis* endotoxin, followed by amino terminal amino acid sequencing, back-translation of the amino acid sequence to design an oligonucleotide probe or use of a related *B. thuringiensis* gene as a probe, followed by cloning of the gene encoding the endotoxin by hybridization are familiar to those skilled in the art and have been described, (see *e.g.*, Donovan *et al.*, 1992); U. S. Patent 5,264,364, each specifically incorporated herein by reference.

Improved expression of Dipteran-inactive Cry2A *B. thuringiensis*  $\delta$ -endotoxins in transgenic plants can be achieved *via* the methods described in this invention. One protein for which improved expression is obtained is Cry2Ab.

Previous work indicated that certain Cry2A  $\delta$ -endotoxins were capable of wider host range specificity than other closely related Cry2A  $\delta$ -endotoxins wherein not only Lepidopteran species, but

Dipteran species also were particularly susceptible to very low toxin doses. In contrast, the closely related Cry2A endotoxins not displaying substantial Dipteran inhibitory activity were thus shown to be more narrow in their host range specificity (Widner et al., 1989, J. Bacteriol. 171:965-974; Widner et al. (a), 1990, J. Bacteriol. 172:2826-2832). These works indicated that Cry2Ab as used herein does not  
5 totally lack Dipteran inhibitory activity, but is simply much less potent than other closely related Cry2A *B. thuringiensis*  $\delta$ -endotoxins. Those works indicated that Cry2Ab in particular was much less effective than Cry2Aa, and hence lacked Dipteran activity when tested against *Aedes aegypti*. There is no one single acceptable means for distinguishing between closely related  $\delta$ -endotoxins, however, as indicated herein, selection of an appropriate Cry2A could be accomplished by using one or a combination of  
10 several methods including but not limited to comparisons in overall amino acid sequence homology, narrowly focused similarity comparisons between Cry2A's in the region specified by amino acid sequence 307-382, or based on IC50 data. Widner et al. demonstrated 50-100 times more Cry2Ab than Cry2Aa was required to obtain a similar IC50 effect on a Dipteran species. Thus, the range of susceptibility of a Dipteran species toward a Cry2A protein could be used as one means of measuring and  
15 distinguishing target insect susceptibility differences between different classes of Cry2A proteins. For example, an IC50 PPM value of about 3-fold greater than that exhibited by Cry2Aa against *Aedes aegypti* could be utilized as a feature for excluding certain Cry2A proteins as lacking substantial Dipteran species inhibitory activity. However, utilizing an approach based on IC50 inhibitory activity ranges should be used with caution, as these values are very dependent upon a number of highly variable conditions  
20 including but not limited to the methods and materials used for assaying the proteins and the physical conditioning of the insects assayed. An alternative means for distinguishing Cry2A  $\delta$ -endotoxins lacking substantial Dipteran species inhibitory activity from  $\delta$ -endotoxins which are not within the scope of the present invention could encompass excluding Cry2A proteins which are greater than about 87% similar in amino acid sequence to Cry2Aa, or more preferentially excluding Cry2A proteins which are greater than  
25 about 90% similar in amino acid sequence to Cry2Aa. In particular, the region of Cry2Aa corresponding to amino acid residues from about 307 to about 382 are believed to be critical for the Dipteran inhibitory activity of the protein, and when substituted for the complementary region of dissimilarity in Cry2Ab, confers Dipteran inhibitory activity to Cry2Ab protein. Thus, an additional means for distinguishing Cry2A  $\delta$ -endotoxins which are within the scope of the present invention could encompass a similarity  
30 comparison of this region of the protein, taking into consideration the level of homology to be avoided when comparing any particular Cry2A  $\delta$ -endotoxins to this region in Cry2Aa. The variable amino acids within this 76 amino acid sequence domain, Cry2A  $\delta$ -endotoxins which are intended to be within the scope of the present invention would preferably be those which are more than from about 80 to about 99 percent similar to Cry2Aa within this sequence, or more preferably those which are more than from about  
35 60 to about 79 percent similar to Cry2Aa within this sequence, or those which are more than from about

40 to about 59 percent similar to Cry2Aa within this sequence, or even more preferably those which are more than from about 24 to about 39 percent similar to Cry2Aa within this sequence, or most preferably those Cry2A  $\delta$ -endotoxins which are more than from about 0 to about 23 percent similar to Cry2Aa within this sequence.

#### 5 4.7 Transformed Plant Cells and Transgenic Plants

A plant transformed with an expression vector of the present invention is also contemplated. A transgenic plant derived from such a transformed or transgenic cell is also contemplated. Those skilled in the art will recognize that a chimeric plant gene containing a structural coding sequence of the present invention can be inserted into the genome of a plant by methods well known in the art. Such methods for DNA transformation of plant cells include *Agrobacterium*-mediated plant transformation, the use of liposomes, transformation using viruses or pollen, electroporation, protoplast transformation, gene transfer into pollen, injection into reproductive organs, injection into immature embryos and particle bombardment. Each of these methods has distinct advantages and disadvantages. Thus, one particular method of introducing genes into a particular plant strain may not necessarily be the most effective for another plant strain, but it is well known which methods are useful for a particular plant strain.

There are many methods for introducing transforming DNA segments into cells, but not all are suitable for delivering DNA to plant cells. Suitable methods are believed to include virtually any method by which DNA can be introduced into a cell, such as infection by *A. tumefaciens* and related *Agrobacterium* strains, direct delivery of DNA such as, for example, by PEG-mediated transformation of protoplasts (Omirulleh *et al.*, 1993), by desiccation/inhibition-mediated DNA uptake, by electroporation, by agitation with silicon carbide fibers, by acceleration of DNA coated particles, *etc.* In certain embodiments, acceleration methods are preferred and include, for example, microprojectile bombardment and the like.

Technology for introduction of DNA into cells is well-known to those of skill in the art. Four general methods for delivering a gene into cells have been described: (1) chemical methods (Graham and van der Eb, 1973); (2) physical methods such as microinjection (Capecchi, 1980), electroporation (Wong and Neumann, 1982; Fromm *et al.*, 1985) and the gene gun (Johnston and Tang, 1994; Fynan *et al.*, 1993); (3) viral vectors (Clapp, 1993; Lu *et al.*, 1993; Eglitis and Anderson, 1988a; 1988b); and (4) receptor-mediated mechanisms (Curiel *et al.*, 1991; 1992; Wagner *et al.*, 1992).

##### 30 4.7.1 Electroporation

The application of brief, high-voltage electric pulses to a variety of animal and plant cells leads to the formation of nanometer-sized pores in the plasma membrane. DNA is taken directly into the cell cytoplasm either through these pores or as a consequence of the redistribution of membrane components that accompanies closure of the pores. Electroporation can be extremely efficient and can be used both for transient expression of cloned genes and for establishment of cell lines that carry integrated copies of the gene of interest. Electroporation, in contrast to calcium phosphate-mediated transfection and

protoplast fusion, frequently gives rise to cell lines that carry one, or at most a few, integrated copies of the foreign DNA.

The introduction of DNA by means of electroporation is well-known to those of skill in the art. To effect transformation by electroporation, one may employ either friable tissues such as a suspension culture of cells, or embryogenic callus, or alternatively, one may transform immature embryos or other organized tissues directly. One would partially degrade the cell walls of the chosen cells by exposing them to pectin-degrading enzymes (pectolyases) or mechanically wounding in a controlled manner, rendering the cells more susceptible to transformation. Such cells would then be recipient to DNA transfer by electroporation, which may be carried out at this stage, and transformed cells then identified by a suitable selection or screening protocol dependent on the nature of the newly incorporated DNA.

#### 4.7.2 Microprojectile Bombardment

A further advantageous method for delivering transforming DNA segments to plant cells is microprojectile bombardment. In this method, particles may be coated with nucleic acids and delivered into cells by a propelling force. Exemplary particles include those comprised of tungsten, gold, platinum, and the like. Using these particles, DNA is carried through the cell wall and into the cytoplasm on the surface of small metal particles as described (Klein *et al.*, 1987; Klein *et al.*, 1988; Kawata *et al.*, 1988). The metal particles penetrate through several layers of cells and thus allow the transformation of cells within tissue explants. The microprojectile bombardment method is preferred for the identification of chloroplast or plastid directed transformation events.

An advantage of microprojectile bombardment, in addition to it being an effective means of reproducibly stably transforming plant cells, is that neither the isolation of protoplasts (Cristou *et al.*, 1988) nor the susceptibility to *Agrobacterium* infection is required. An illustrative embodiment of a method for delivering DNA into plant cells by acceleration is a Biolistics Particle Delivery System, which can be used to propel particles coated with DNA or cells through a screen, such as a stainless steel or Nytex screen, onto a filter surface covered with the plant cultured cells in suspension. The screen disperses the particles so that they are not delivered to the recipient cells in large aggregates. It is believed that a screen intervening between the projectile apparatus and the cells to be bombarded reduces the size of projectiles aggregate and may contribute to a higher frequency of transformation by reducing damage inflicted on the recipient cells by projectiles that are too large.

For the bombardment, cells in suspension are preferably concentrated on filters or solid culture medium. Alternatively, immature embryos or other target cells may be arranged on solid culture medium. The cells to be bombarded are positioned at an appropriate distance below the microprojectile stopping plate. If desired, one or more screens are also positioned between the acceleration device and the cells to be bombarded. Through the use of techniques set forth herein one may obtain up to 1000 or more foci of cells transiently expressing a marker gene. The number of cells in a focus which express the exogenous gene product 48 hours post-bombardment often range from 1 to 10 and average 1 to 3.

In bombardment transformation, one may optimize the prebombardment culturing conditions and the bombardment parameters to yield the maximum numbers of stable transformants. Both the physical and biological parameters for bombardment are important in this technology. Physical factors are those that involve manipulating the DNA/microprojectile precipitate or those that affect the flight and velocity of either the macro- or microprojectiles. Biological factors include all steps involved in manipulation of cells before and immediately after bombardment, the osmotic adjustment of target cells to help alleviate the trauma associated with bombardment, and also the nature of the transforming DNA, such as linearized DNA or intact supercoiled plasmids. It is believed that pre-bombardment manipulations are especially important for successful transformation of immature plant embryos.

Accordingly, it is contemplated that one may desire to adjust various of the bombardment parameters in small scale studies to fully optimize the conditions. One may particularly wish to adjust physical parameters such as gap distance, flight distance, tissue distance, and helium pressure. One may also minimize the trauma reduction factors (TRFs) by modifying conditions which influence the physiological state of the recipient cells and which may therefore influence transformation and integration efficiencies. For example, the osmotic state, tissue hydration and the subculture stage or cell cycle of the recipient cells may be adjusted for optimum transformation. The execution of other routine adjustments will be known to those of skill in the art in light of the present disclosure.

The methods of particle-mediated transformation is well-known to those of skill in the art. U. S. Patent 5,015,580 (specifically incorporated herein by reference) describes the transformation of soybeans using such a technique.

#### 4.7.3 *Agrobacterium*-Mediated Transfer

*Agrobacterium*-mediated transfer is a widely applicable system for introducing genes into plant cells because the DNA can be introduced into whole plant tissues, thereby bypassing the need for regeneration of an intact plant from a protoplast. The use of *Agrobacterium*-mediated plant integrating vectors to introduce DNA into plant cells is well known in the art. See, for example, the methods described (Fraley *et al.*, 1985; Rogers *et al.*, 1987). The genetic engineering of cotton plants using *Agrobacterium*-mediated transfer is described in U. S. Patent 5,004,863 (specifically incorporated herein by reference); like transformation of lettuce plants is described in U. S. Patent 5,349,124 (specifically incorporated herein by reference); and the *Agrobacterium*-mediated transformation of soybean is described in U. S. Patent 5,416,011 (specifically incorporated herein by reference). Further, the integration of the Ti-DNA is a relatively precise process resulting in few rearrangements. The region of DNA to be transferred is defined by the border sequences, and intervening DNA is usually inserted into the plant genome as described (Spielmann *et al.*, 1986; Jorgensen *et al.*, 1987).

Modern *Agrobacterium* transformation vectors are capable of replication in *E. coli* as well as *Agrobacterium*, allowing for convenient manipulations as described (Klee *et al.*, 1985). Moreover, recent technological advances in vectors for *Agrobacterium*-mediated gene transfer have improved the

arrangement of genes and restriction sites in the vectors to facilitate construction of vectors capable of expressing various polypeptide coding genes. The vectors described (Rogers *et al.*, 1987), have convenient multi-linker regions flanked by a promoter and a polyadenylation site for direct expression of inserted polypeptide coding genes and are suitable for present purposes. In addition, *Agrobacterium* containing both armed and disarmed Ti genes can be used for the transformations. In those plant varieties where *Agrobacterium*-mediated transformation is efficient, it is the method of choice because of the facile and defined nature of the gene transfer.

*Agrobacterium*-mediated transformation of leaf disks and other tissues such as cotyledons and hypocotyls appears to be limited to plants that *Agrobacterium* naturally infects. *Agrobacterium*-mediated transformation is most efficient in dicotyledonous plants. Few monocots appear to be natural hosts for *Agrobacterium*, although transgenic plants have been produced in asparagus using *Agrobacterium* vectors as described (Bytner *et al.*, 1987). Other monocots recently have also been transformed with *Agrobacterium*. Included in this group are corn (Ishida *et al.*) and rice (Cheng *et al.*).

A transgenic plant formed using *Agrobacterium* transformation methods typically contains a single gene on one chromosome. Such transgenic plants can be referred to as being heterozygous for the added gene. However, inasmuch as use of the word "heterozygous" usually implies the presence of a complementary gene at the same locus of the second chromosome of a pair of chromosomes, and there is no such gene in a plant containing one added gene as here, it is believed that a more accurate name for such a plant is an independent segregant, because the added, exogenous gene segregates independently during mitosis and meiosis.

An independent segregant may be preferred when the plant is commercialized as a hybrid, such as corn. In this case, an independent segregant containing the gene is crossed with another plant, to form a hybrid plant that is heterozygous for the gene of interest.

An alternate preference is for a transgenic plant that is homozygous for the added structural gene; *i.e.* a transgenic plant that contains two added genes, one gene at the same locus on each chromosome of a chromosome pair. A homozygous transgenic plant can be obtained by sexually mating (selfing) an independent segregant transgenic plant that contains a single added gene, germinating some of the seed produced and analyzing the resulting plants produced for gene of interest activity and mendelian inheritance indicating homozygosity relative to a control (native, non-transgenic) or an independent segregant transgenic plant.

Two different transgenic plants can be mated to produce offspring that contain two independently segregating added, exogenous genes. Selfing of appropriate progeny can produce plants that are homozygous for both added, exogenous genes that encode a polypeptide of interest. Back-crossing to a parental plant and out-crossing with a non-transgenic plant are also contemplated.

Transformation of plant protoplasts can be achieved using methods based on calcium phosphate precipitation, polyethylene glycol treatment, electroporation, and combinations of these treatments (see e.g., Potrykus *et al.*, 1985; Lorz *et al.*, 1985; Fromm *et al.*, 1985; Uchimiya *et al.*, 1986; Callis *et al.*, 1987; Marcotte *et al.*, 1988).

5 Application of these systems to different plant germplasm depends upon the ability to regenerate that particular plant variety from protoplasts. Illustrative methods for the regeneration of cereals from protoplasts are described (see, e.g., Fujimura *et al.*, 1985; Toriyama *et al.*, 1986; Yamada *et al.*, 1986; Abdullah *et al.*, 1986).

To transform plant germplasm that cannot be successfully regenerated from protoplasts, other  
10 ways to introduce DNA into intact cells or tissues can be utilized. For example, regeneration of cereals from immature embryos or explants can be effected as described (Vasil, 1988).

#### 4.8 Gene Expression in Plants

Unmodified bacterial genes are often poorly expressed in transgenic plant cells. Plant codon usage more closely resembles that of humans and other higher organisms than unicellular organisms, such  
15 as bacteria. Several reports have disclosed methods for improving expression of recombinant genes in plants (Murray *et al.*, 1989; Diehn *et al.*, 1996; Iannacone *et al.*, 1997; Rouwendal *et al.*, 1997; Futterer *et al.*, 1997; and Futterer and Hohn, 1996). These reports disclose various methods for engineering coding sequences to represent sequences which are more efficiently translated based on plant codon frequency tables, improvements in codon third base position bias, using recombinant sequences which avoid suspect  
20 polyadenylation or A/T rich domains or intron splicing consensus sequences. While these methods for synthetic gene construction are notable, synthetic genes of the present invention were prepared according to the method of Brown *et al.* (US Pat. No. 5,689,052; 1997), which is herein incorporated in its entirety by reference. Thus, the present invention provides a method for preparing synthetic plant genes express  
in planta a desired protein product at levels significantly higher than the wild-type genes. Briefly,  
25 according to Brown *et al.*, the frequency of rare and semi-rare monocotyledonous codons in a polynucleotide sequence encoding a desired protein are reduced and replaced with more preferred monocotyledonous codons. Enhanced accumulation of a desired polypeptide encoded by a modified polynucleotide sequence in a monocotyledonous plant is the result of increasing the frequency of preferred codons by analyzing the coding sequence in successive six nucleotide fragments and altering the  
30 sequence based on the frequency of appearance of the six-mers as to the frequency of appearance of the rarest 284, 484, and 664 six-mers in monocotyledonous plants. Furthermore, Brown *et al.* disclose the enhanced expression of a recombinant gene by applying the method for reducing the frequency of rare codons with methods for reducing the occurrence of polyadenylation signals and intron splice sites in the nucleotide sequence, removing self-complementary sequences in the nucleotide sequence and replacing  
35 such sequences with nonself-complementary nucleotides while maintaining a structural gene encoding the polypeptide, and reducing the frequency of occurrence of 5'-CG-3' dinucleotide pairs in the nucleotide

sequence. These steps are performed sequentially and have a cumulative effect resulting in a nucleotide sequence containing a preferential utilization of the more-preferred monocotyledonous codons for monocotyledonous plants for a majority of the amino acids present in the desired polypeptide.

The work described herein has identified methods of potentiating *in planta* expression of *B. thuringiensis*  $\delta$ -endotoxins, which confer resistance to insect pathogens when incorporated into the nuclear, plastid, or chloroplast genome of susceptible plants. U. S. Patent 5,500,365 (specifically incorporated herein by reference) describes a method for synthesizing plant genes to optimize the expression level of the protein for which the synthesized gene encodes. This method relates to the modification of the structural gene sequences of the exogenous transgene, to make them more "plant-like" and therefore more likely to be translated and expressed by the plant, monocot or dicot. However, the method as disclosed in U. S. Patent 5,689,052 provides for enhanced expression of transgenes, preferably in monocotyledonous plants.

#### 4.9 Production of Insect-Resistant Transgenic Plants

Thus, the amount of a gene coding for a polypeptide of interest (*i.e.* a bacterial crystal protein or  $\delta$ -endotoxin polypeptide and a plastid targeting peptide) can be increased in plants by transforming those plants using transformation methods such as those disclosed herein at Section 4.7. In particular, chloroplast or plastid transformation can result in desired coding sequences being present in up to about 10,000 copies per cell in tissues containing these subcellular organelle structures (McBride *et al.*, *Bio/Technology* 13:362-365, 1995).

DNA can also be introduced into plants by direct DNA transfer into pollen as described (Zhou *et al.*, 1983; Hess, 1987). Expression of polypeptide coding genes can be obtained by injection of the DNA into reproductive organs of a plant as described (Pena *et al.*, 1987). DNA can also be injected directly into the cells of immature embryos and the rehydration of desiccated embryos as described (Neuhaus *et al.*, 1987; Benbrook *et al.*, 1986).

##### 4.9.1 Selection of Transformed Cells

After effecting delivery of exogenous DNA to recipient cells, the next step to obtain a transgenic plant generally concern identifying the transformed cells for further culturing and plant regeneration. As mentioned herein, in order to improve the ability to identify transformants, one may desire to employ a selectable or screenable marker gene as, or in addition to, the expressible gene of interest. In this case, one would then generally assay the potentially transformed cell population by exposing the cells to a selective agent or agents, or one would screen the cells for the desired marker gene trait.

An exemplary embodiment of methods for identifying transformed cells involves exposing the transformed cultures to a selective agent, such as a metabolic inhibitor, an antibiotic, herbicide or the like. Cells which have been transformed and have stably integrated a marker gene conferring resistance to the selective agent used, will grow and divide in culture. Sensitive cells will not be amenable to further culturing. One example of a preferred marker gene confers resistance to glyphosate. When this gene is



used as a selectable marker, the putatively transformed cell culture is treated with glyphosate. Upon treatment, transgenic cells will be available for further culturing while sensitive, or non-transformed cells, will not. This method is described in detail in U. S. Patent 5,569,834, which is specifically incorporated herein by reference. Another example of a preferred selectable marker system is the neomycin phosphotransferase (*npII*) resistance system by which resistance to the antibiotic kanamycin is conferred, as described in U. S. Patent 5,569,834 (specifically incorporated herein by reference). Again, after transformation with this system, transformed cells will be available for further culturing upon treatment with kanamycin, while non-transformed cells will not. Yet another preferred selectable marker system involves the use of a gene construct conferring resistance to paromomycin. Use of this type of a selectable marker system is described in U. S. Patent 5,424,412 (specifically incorporated herein by reference).

All contemplated assays are nondestructive and transformed cells may be cultured further following identification. Another screenable marker which may be used is the gene coding for green fluorescent protein.

Transplastonomic selection (selection of plastid or chloroplast transformation events) is simplified by taking advantage of the sensitivity of chloroplasts or plastids to spectinomycin, an inhibitor of plastid or chloroplast protein synthesis, but not of protein synthesis by the nuclear genome encoded cytoplasmic ribosomes. Spectinomycin prevents the accumulation of chloroplast proteins required for photosynthesis and so spectinomycin resistant transformed plant cells may be distinguished on the basis of their difference in color: the resistant, transformed cells are green, whereas the sensitive cells are white, due to inhibition of plastid-protein synthesis. Transformation of chloroplasts or plastids with a suitable bacterial *aad* gene, or with a gene encoding a spectinomycin resistant plastid or chloroplast functional ribosomal RNA provides a means for selection and maintenance of transplastonomic events (Maliga, Trends in Biotechnology 11:101-106, 1993).

It is further contemplated that combinations of screenable and selectable markers will be useful for identification of transformed cells. In some cell or tissue types a selection agent, such as glyphosate or kanamycin, may either not provide enough killing activity to clearly recognize transformed cells or may cause substantial nonselective inhibition of transformants and nontransformants alike, thus causing the selection technique to not be effective. It is proposed that selection with a growth inhibiting compound, such as glyphosate at concentrations below those that cause 100% inhibition followed by screening of growing tissue for expression of a screenable marker gene such as kanamycin would allow one to recover transformants from cell or tissue types that are not amenable to selection alone. It is proposed that combinations of selection and screening may enable one to identify transformants in a wider variety of cell and tissue types.

#### 4.9.2 Regeneration of Transformants

The development or regeneration of plants from either single plant protoplasts or various explants is well known in the art (Weissbach and Weissbach, 1988). This regeneration and growth process typically includes the steps of selection of transformed cells, culturing those individualized cells through the usual stages of embryonic development through the rooted plantlet stage. Transgenic embryos and seeds are similarly regenerated. The resulting transgenic rooted shoots are thereafter planted in an appropriate plant growth medium such as soil.

The development or regeneration of plants containing the foreign, exogenous gene that encodes a polypeptide of interest introduced by *Agrobacterium* from leaf explants can be achieved by methods well known in the art such as described (Horsch *et al.*, 1985). In this procedure, transformants are cultured in the presence of a selection agent and in a medium that induces the regeneration of shoots in the plant strain being transformed as described (Fraley *et al.*, 1983). In particular, U. S. Patent 5,349,124 (specification incorporated herein by reference) details the creation of genetically transformed lettuce cells and plants resulting therefrom which express hybrid crystal proteins conferring insecticidal activity against *Lepidopteran* larvae to such plants.

This procedure typically produces shoots within two to four months and those shoots are then transferred to an appropriate root-inducing medium containing the selective agent and an antibiotic to prevent bacterial growth. Shoots that rooted in the presence of the selective agent to form plantlets are then transplanted to soil or other media to allow the production of roots. These procedures vary depending upon the particular plant strain employed, such variations being well known in the art.

Preferably, the regenerated plants are self-pollinated to provide homozygous transgenic plants, or pollen obtained from the regenerated plants is crossed to seed-grown plants of agronomically important, preferably inbred lines. Conversely, pollen from plants of those important lines is used to pollinate regenerated plants. A transgenic plant of the present invention containing a desired polypeptide is cultivated using methods well known to one skilled in the art.

A transgenic plant of this invention thus has an increased amount of a coding region encoding a *B. thuringiensis*  $\delta$ -endotoxin polypeptide and a plastid targeting peptide. A preferred transgenic plant is an independent segregant and can transmit that gene and its activity to its progeny. A more preferred transgenic plant is homozygous for that gene, and transmits that gene to all of its offspring on sexual mating. Seed from a transgenic plant may be grown in the field or greenhouse, and resulting sexually mature transgenic plants are self-pollinated to generate true breeding plants. The progeny from these plants become true breeding lines that are evaluated for increased expression of the *B. thuringiensis* transgene.

#### 4.10 Identification of Transgenic Plant Events with Insect Tolerance

To identify a transgenic plant expressing high levels of the  $\delta$ -endotoxin of interest, it is necessary to screen the herbicide or antibiotic resistant transgenic, regenerated plants ( $R_0$  generation) for insecticidal

activity and/or expression of the gene of interest. This can be accomplished by various methods well known to those skilled in the art, including but not limited to: 1) obtaining small tissue samples from the transgenic  $R_0$  plant and directly assaying the tissue for activity against susceptible insects in parallel with tissue derived from a non-expressing, negative control plant. For example,  $R_0$  transgenic corn plants expressing *B. thuringiensis* endotoxins such as Cry2Ab can be identified by assaying leaf tissue derived from such plants for activity against ECB; 2) analysis of protein extracts by enzyme linked immunoassays (ELISAs) specific for the gene of interest (Cry2Ab); or 3) reverse transcriptase PCR<sup>TM</sup> (RT PCR<sup>TM</sup>) to identify events expressing the gene of interest.

#### 4.11 Isolating Homologous Gene and Gene Fragments

The genes and  $\delta$ -endotoxins according to the subject invention include not only the full length sequences disclosed herein but also fragments of these sequences, or fusion proteins, which retain the characteristic insecticidal activity of the sequences specifically exemplified herein.

It should be apparent to a person of skill in this art that insecticidal  $\delta$ -endotoxins can be identified and obtained through several means. The specific genes, or portions thereof, may be obtained from a culture depository, or constructed synthetically, for example, by use of a gene machine. Variations of these genes may be readily constructed using standard techniques for making point mutations. Also, fragments of these genes can be made using commercially available exonucleases or endonucleases according to standard procedures. For example, enzymes such as *Bal31* or site-directed mutagenesis can be used to systematically cut off nucleotides from the ends of these genes. Also, genes which code for active fragments may be obtained using a variety of other restriction enzymes. Proteases may be used to directly obtain active fragments of these  $\delta$ -endotoxins.

Equivalent  $\delta$ -endotoxins and/or genes encoding these  $\delta$ -endotoxins can also be isolated from *Bacillus* strains and/or DNA libraries using the teachings provided herein. For example, antibodies to the  $\delta$ -endotoxins disclosed and claimed herein can be used to identify and isolate other  $\delta$ -endotoxins from a mixture of proteins. Specifically, antibodies may be raised to the portions of the  $\delta$ -endotoxins which are most constant and most distinct from other *B. thuringiensis*  $\delta$ -endotoxins. These antibodies can then be used to specifically identify equivalent  $\delta$ -endotoxins with the characteristic insecticidal activity by immunoprecipitation, enzyme linked immunoassay (ELISA), or Western blotting.

A further method for identifying the  $\delta$ -endotoxins and genes of the subject invention is through the use of oligonucleotide probes. These probes are nucleotide sequences having a detectable label. As is well known in the art, if the probe molecule and nucleic acid sample hybridize by forming a strong bond between the two molecules, it can be reasonably assumed that the probe and sample are essentially identical. The probe's detectable label provides a means for determining in a known manner whether hybridization has occurred. Such a probe analysis provides a rapid method for identifying insecticidal  $\delta$ -endotoxin genes of the subject invention.

The nucleotide segments which are used as probes according to the invention can be synthesized by use of DNA synthesizers using standard procedures. In the use of the nucleotide segments as probes, the particular probe is labeled with any suitable label known to those skilled in the art, including radioactive and non-radioactive labels. Typical radioactive labels include  $^{32}\text{P}$ ,  $^{125}\text{I}$ ,  $^{35}\text{S}$ , or the like. A probe labeled with a radioactive isotope can be constructed from a nucleotide sequence complementary to the DNA sample by a conventional nick translation reaction, using a DNase and DNA polymerase. The probe and sample can then be combined in a hybridization buffer solution and held at an appropriate temperature until annealing occurs. Thereafter, the membrane is washed free of extraneous materials, leaving the sample and bound probe molecules typically detected and quantified by autoradiography and/or liquid scintillation counting.

Non-radioactive labels include, for example, ligands such as biotin or thyroxine, as well as enzymes such as hydrolyses or peroxidases, or the various chemiluminescers such as luciferin, or fluorescent compounds like fluorescein and its derivatives. The probe may also be labeled at both ends with different types of labels for ease of separation, as, for example, by using an isotopic label at the end mentioned above and a biotin label at the other end.

Duplex formation and stability depend on substantial complementarity between the two strands of a hybrid, and, as noted above, a certain degree of mismatch can be tolerated. Therefore, the probes of the subject invention include mutations (both single and multiple), deletions, insertions of the described sequences, and combinations thereof, wherein said mutations, insertions and deletions permit formation of stable hybrids with the target polynucleotide of interest. Mutations, insertions, and deletions can be produced in a given polynucleotide sequence in many ways, by methods currently known to an ordinarily skilled artisan, and perhaps by other methods which may become known in the future.

The potential variations in the probes listed is due, in part, to the redundancy of the genetic code. Because of the redundancy of the genetic code, more than one coding nucleotide triplet (codon) can be used for most of the amino acids used to make proteins. Therefore different nucleotide sequences can code for a particular amino acid. Thus, the amino acid sequences of the *B. thuringiensis*  $\delta$ -endotoxins and peptides, and the plastid targeting peptides and the polynucleotides which code for them, can be prepared by equivalent nucleotide sequences encoding the same amino acid sequence of the protein or peptide. Accordingly, the subject invention includes such equivalent nucleotide sequences. Also, inverse or complement sequences are an aspect of the subject invention and can be readily used by a person skilled in this art. In addition it has been shown that proteins of identified structure and function may be constructed by changing the amino acid sequence if such changes do not alter the protein secondary structure (Kaiser and Kezdy, 1984). Thus, the subject invention includes mutants of the amino acid sequence depicted herein which do not alter the protein secondary structure, or if the structure is altered, the biological activity is substantially retained. Further, the invention also includes mutants of organisms hosting all or part of a gene encoding a  $\delta$ -endotoxin and gene encoding a plastid targeting peptide, as

discussed in the present invention. Such mutants can be made by techniques well known to persons skilled in the art. For example, UV irradiation can be used to prepare mutants of host organisms. Likewise, such mutants may include asporogenous host cells which also can be prepared by procedures well known in the art.

#### 5 4.12 Site-Specific Mutagenesis

Site-specific mutagenesis is a technique useful in the preparation of individual peptides, or biologically functional equivalent proteins or peptides, through specific mutagenesis of the underlying DNA. The technique further provides a ready ability to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide  
10 sequence changes into the DNA. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 17 to 25 nucleotides in length is preferred, with about 5 to 10 residues on  
15 both sides of the junction of the sequence being altered.

In general, the technique of site-specific mutagenesis is well known in the art, as exemplified by various publications. As will be appreciated, the technique typically employs a phage vector which exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage. These phage are readily commercially available and their use is  
20 generally well known to those skilled in the art. Double stranded plasmids are also routinely employed in site directed mutagenesis which eliminates the step of transferring the gene of interest from a plasmid to a phage.

In general, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector or melting apart of two strands of a double stranded vector which includes within  
25 its sequence a DNA sequence which encodes the desired peptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically. This primer is then annealed with the single-stranded vector, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the  
30 desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement.

The preparation of sequence variants of the selected peptide-encoding DNA segments using site-directed mutagenesis is provided as a means of producing potentially useful species and is not meant to be  
35 limiting as there are other ways in which sequence variants of peptides and the DNA sequences encoding them may be obtained. For example, recombinant vectors encoding the desired peptide sequence may be

treated with mutagenic agents, such as hydroxylamine, to obtain sequence variants. Such procedures may favorably change the protein's biochemical and biophysical characteristics or its mode of action. These include, but are not limited to: 1) improved  $\delta$ -endotoxin formation, 2) improved protein stability or reduced protease degradation, 3) improved insect membrane receptor recognition and binding, 4) improved oligomerization or channel formation in the insect midgut endothelium, and 5) improved insecticidal activity or insecticidal specificity due to any or all of the reasons stated above.

#### 4.13 Biological Functional Equivalents

Modification and changes may be made in the structure of the peptides of the present invention and DNA segments which encode them and still obtain a functional molecule that encodes a protein or peptide with desirable characteristics. The biologically functional equivalent peptides, polypeptides, and proteins contemplated herein should possess about 80% or greater sequence similarity, preferably about 85% or greater sequence similarity, and most preferably about 90% or greater sequence similarity, to the sequence of, or corresponding moiety within, the fundamental cry2Ab amino acid sequence.

The following is a discussion based upon changing the amino acids of a protein to create an equivalent, or even an improved, second-generation molecule. In particular embodiments of the invention, mutated crystal proteins are contemplated to be useful for increasing the insecticidal activity of the protein, and consequently increasing the insecticidal activity and/or expression of the recombinant transgene in a plant cell. The amino acid changes may be achieved by changing the codons of the DNA sequence, according to the codons given in Table 3.

Table 3

Amino Acid			Codons						
Alanine	Ala	A	GCA	GCC	GCG	GCU			
Cysteine	Cys	C	UGC	UGU					
Aspartic acid	Asp	D	GAC	GAU					
Glutamic acid	Glu	E	GAA	GAG					
Phenylalanine	Phe	F	UUC	UUU					
Glycine	Gly	G	GGA	GGC	GGG	GGU			
Histidine	His	H	CAC	CAU					
Isoleucine	Ile	I	AUA	AUC	AUU				
Lysine	Lys	K	AAA	AAG					
Leucine	Leu	L	UUA	UUG	CUA	CUC	CUG	CUU	
Methionine	Met	M	AUG						
Asparagine	Asn	N	AAC	AAU					

Amino Acid			Codons					
Proline	Pro	P	CCA	CCC	CCG	CCU		
Glutamine	Gln	Q	CAA	CAG				
Arginine	Arg	R	AGA	AGG	CGA	CGC	CGG	CGU
Serine	Ser	S	AGC	AGU	UCA	UCC	UCG	UCU
Threonine	Thr	T	ACA	ACC	ACG	ACU		
Valine	Val	V	GUA	GUC	GUG	GUU		
Tryptophan	Trp	W	UGG					
Tyrosine	Tyr	Y	UAC	UAU				

For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of interactive binding capacity with structures such as, for example, antigen-binding regions of antibodies or binding sites on substrate molecules. Since it is the interactive capacity and nature of a protein that defines that protein's biological functional activity, certain amino acid sequence substitutions can be made in a protein sequence, and, of course, its underlying DNA coding sequence, and nevertheless obtain a protein with like properties. It is thus contemplated by the inventors that various changes may be made in the peptide sequences of the disclosed compositions, or corresponding DNA sequences which encode said peptides without appreciable loss of their biological utility or activity.

In making such changes, the hydropathic index of amino acids may be considered. The importance of the hydropathic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte and Doolittle, 1982, incorporate herein by reference). It is accepted that the relative hydropathic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like.

Each amino acid has been assigned a hydropathic index on the basis of their hydrophobicity and charge characteristics (Kyte and Doolittle, 1982), these are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydropathic index or score and still result in a protein with similar biological activity, *i.e.* still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino

acids whose hydrophobic indices are within  $\pm 2$  is preferred, those which are within  $\pm 1$  are particularly preferred, and those within  $\pm 0.5$  are even more particularly preferred.

It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U. S. Patent 4,554,101, incorporated herein by reference, states that the  
5 greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein.

As detailed in U. S. Patent 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0  $\pm$  1); glutamate (+3.0  $\pm$  1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5  $\pm$  1); alanine (-  
10 0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4).

It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent, and in particular, an immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within  $\pm 2$  is preferred,  
15 those which are within  $\pm 1$  are particularly preferred, and those within  $\pm 0.5$  are even more particularly preferred.

As outlined above, amino acid substitutions are generally therefore based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions which take various of the foregoing characteristics  
20 into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

Polynucleotides encoding  $\delta$ -endotoxins derived from *B. thuringiensis* are known by those skilled in the art, to be poorly expressed when incorporated into the nuclear DNA of transgenic plants (reviewed by Diehn *et al.*, 1996). Preferably, a nucleotide sequence encoding the  $\delta$ -endotoxin of interest is designed  
25 essentially as described in U. S. Patent 5,500,365 and 5,689,052 (each specifically incorporated herein by reference). Examples of nucleotide sequences useful for expression include but are not limited to, *cry2Ab* (SEQ ID NO:1).

Peptides, polypeptides, and proteins biologically functionally equivalent to *Cry2Ab* include amino acid sequences containing conservative amino acid changes in the fundamental sequence shown in  
30 SEQ ID NO:2. In such amino acid sequences, one or more amino acids in the fundamental sequence is (are) substituted with another amino acid(s), the charge and polarity of which is similar to that of the native amino acid, *i.e.* a conservative amino acid substitution, resulting in a silent change.

Substitutes for an amino acid within the fundamental polypeptide sequence can be selected from other members of the class to which the naturally occurring amino acid belongs. Amino acids can be  
35 divided into the following four groups: (1) acidic amino acids; (2) basic amino acids; (3) neutral polar



amino acids; and (4) neutral non-polar amino acids. Representative amino acids within these various groups include, but are not limited to: (1) acidic (negatively charged) amino acids such as aspartic acid and glutamic acid; (2) basic (positively charged) amino acids such as arginine, histidine, and lysine; (3) neutral polar amino acids such as glycine, serine, threonine, cysteine, cystine, tyrosine, asparagine, and glutamine; (4) neutral nonpolar (hydrophobic) amino acids such as alanine, leucine, isoleucine, valine, proline, phenylalanine, tryptophan, and methionine.

Conservative amino acid changes within the fundamental polypeptide sequence can be made by substituting one amino acid within one of these groups with another amino acid within the same group. Biologically functional equivalents of *cry2Ab* can have 10 or fewer conservative amino acid changes, more preferably seven or fewer conservative amino acid changes, and most preferably five or fewer conservative amino acid changes. The encoding nucleotide sequence (gene, plasmid DNA, cDNA, or synthetic DNA) will thus have corresponding base substitutions, permitting it to encode biologically functional equivalent forms of *cry2Ab*.

## 5.0 Examples

The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

### 5.1 Example 1 - Increased Expression of Cry2Ab by Targeted Vectors

Expression of the Cry2Ab protein in corn plants transformed with targeted and non-targeted Cry2Ab expression vectors was compared and was significantly higher in plants with the targeted vector. Untargeted Cry2Ab plant expression vectors pMON26800 and pMON30463 contain an expression cassette composed of an enhanced CaMV35S promoter, a maize hsp70 intron, a synthetic *cry2Ab* gene with translational initiation and termination codons (SEQ ID NO:1), and a nopaline synthase polyadenylation site.

The targeted plant expression vector pMON30464 (SEQ ID NO:16) contains an expression cassette including enhanced CaMV35S promoter, a maize hsp70 intron, a maize ssRUBISCO chloroplast transit peptide (SEQ ID NO:3) fused in frame to a synthetic *cry2Ab* gene, and a nopaline synthase polyadenylation site.

All vectors (pMON26800, pMON30463, and pMON30464) also contain a cassette conferring paromomycin resistance to transformed plant tissue. In the case of pMON26800, this cassette consists of an enhanced CaMV35S promoter, a maize hsp70 intron, a neomycin phosphotransferase gene with a translational initiation and termination codons, and a nopaline synthase polyadenylation site. In the case

of pMON30463 and pMON30464, this cassette consists of a CaMV35S promoter, a neomycin phosphotransferase gene with a translational initiation and termination codons, and a nopaline synthase polyadenylation site. Transgenic corn plants resistant to paromomycin were derived essentially as described in U. S. Patent 5,424,412 (specifically incorporated herein by reference).

5 Leaf tissue from independently transformed transgenic events in the R<sub>0</sub> stage was subjected to quantitative analysis of Cry2Ab protein levels by a quantitative ELISA assay. This ELISA used a direct sandwich technique that used a monoclonal capture antibody raised against Cry2Aa, a different Cry2Aa monoclonal antibody conjugated to alkaline phosphatase as the secondary antibody, and purified Cry2Aa protein as a standard.

10 Comparison of Cry2Ab expression levels in pMON30463 (non-targeted) and pMON30464 (targeted) corn plants show that non-targeted Cry2Ab expression does not exceed 15 ppm while targeted expression is frequently higher than 100 ppm (Table 4). Protein blot analyses confirm that the increased level of cross reactive material produced by pMON30464 (targeted) were due to increased accumulation of an approximately Mr 71,000 protein that co-migrates with Cry2Ab produced by pMON30463 (non-  
15 targeted) and Cry2Aa standard from *B. thuringiensis*. This data indicates that the targeting peptide fused to the N-terminus of Cry2Ab protein was efficiently processed or removed.

Increased expression of Cry2Ab in pMON30464 (targeted) vectors relative to pMON26800 (non-targeted) vectors was also observed in R<sub>1</sub> progeny plants derived from the original R<sub>0</sub> transgenic events, indicating that high expression is heritable (Table 5).

20 Table 4

Expression of Cry2Ab in R<sub>0</sub> Corn Transformed with Targeted (pMON30464) and Untargeted (pMON30463) Expression Vectors: Distribution of Expression Levels in Different Events

Vector	Total Events	Total ECB +	0 ppm	0-5 ppm	5-15 ppm	15-50 ppm	50- 100 ppm	100- 200 ppm	>200 ppm
non- targeted (30463)	16	3 (19%)	0	0	3	0	0	0	0
targeted (30464)	40	14 (35%)	0	2	2	0	0	4	5

Table 5

Expression of Cry2Ab In R<sub>1</sub> Corn Transformed with Targeted (pMON30464) and Untargeted (pMON26800) Expression Vectors: Distribution of Expression Levels in Different Events

Vector	Total # events assayed	0 ppm	0-5 ppm	5-15 ppm	15-50 ppm	50-100 ppm	100-200 ppm	>200 ppm
non-targeted (26800)	28	0	18	10	0	0	0	0
targeted (30464)	33	5	3	2	0	2	4	17

To effectively control insects that feed on a variety of corn tissues, it is critical that the insecticidal protein be expressed at high levels throughout all potential feeding sites. To determine if the increases in targeted expression of Cry2Ab occur in other tissues, independent targeted and non-targeted transgenic events representing the high expressing lines obtained with the respective vector types were assayed for Cry2Ab expression levels in parallel. Expression of Cry2Ab is increased in virtually all of the corn tissues attacked by pests such as *Ostrina nubilalis* and *Helicoverpa zea* by targeted expression (Table 6). Uniform high level expression of this type is especially valuable in that it is less likely to permit evolved resistance of target pests *via* behavioral (feeding) adaptation.

Table 6

Targeted and Untargeted Cry2Ab Expression in Transgenic Maize

Vector	Event	N	Root	Leaf	sheath	stalk	shank	husk	silk	cob	kernel
N30pMON 30464	#1	1	13.1	117.6	140.8	514.9	397.5	121.8	130.5	165.2	106.9
	#2	4	11.3 ± 4.5	105 ± 12	121 ± 25	96 ± 18	134 ± 38	52 ± 9.1	101 ± 11	113 ± 45	170 ± 36
N30pMON 26800	#1	2	1.2 ± 0.4	10 ± 5.3	20 ± 12	28 ± 5.6	29 ± 7.5	7.6 ± 7.6	46 ± 9.9	9.6 ± 9.6	10.9 ± 4.6

Expression in µg Cry2Ab / gm fresh weight (root and leaf) or dry weight tissue (sheath, stalk, shank, husk, silk, cob, kernel) shown ± standard deviation (L30464 #2) or range (L26800#1).

Further analyses indicate that the increased levels of Cry2Ab protein produced by pMON30464 result in a commensurate increase in the level of bioactivity as measured directly in feeding assays. To assess the level of insecticidal activity produced, corn leaf tissue from control (non-transgenic), targeted

(pmon30464), and non-targeted (pMON30464) plants was assayed for activity against *Heliothis virescens* in tissue diet overlay studies (Table 7). Two concentrations of tissue (0.0016 and 0.0031 %) were bioassayed and the same sample of tissue used in the diet overlay was also subjected to quantitative ELISA determinations of Cry2Ab levels. The 7.5-fold increase in Cry2Ab levels in targeted (pMON30464) samples relative to the non-targeted (pMON30463) samples clearly correlates with the corresponding 6-fold difference in mean larval weight observed at both concentration rates. These data thus indicate that the increased levels of Cry2Ab produced by pMON30464 result in commensurate increases in the level of bioactivity.

Table 7

Correlation of Increased Cry2Ab Expression Levels with Increased Bioactivity in *Heliothis virescens* Tissue Diet Overlay Bioassay

Tissue Sample	Cry2Ab Conc. (ppm)	Tissue Conc. 1 (0.0031% Tissue)	Tissue Conc. 2 (0.0016% Tissue)
		Mean Larval Wt. (mg)	Mean Larval Wt. (mg)
Control	0.0	22.00	24.6
Targeted	444	1.2	2.1
Untargeted	60	7.3	12.7

## 5.2 Example 2 - Plastid Targeting of Cry2Ab Increases Frequency of Agronomically-Normal Plants Recovered from Transformation

To obtain a commercially viable transgene-based insect control trait, it is crucial that an event with normal plant growth characteristics be obtained. In most instances a fairly large number of independent transgenic events are advanced into field tests to insure that an event that meets all of the key criteria (effective insect control, normal Mendelian behavior of the transgene, and normal growth characteristics or agronomics) will be identified. Methods that increase the frequency with which normal events are obtained are clearly valuable as they increase the odds of identifying an event that can be commercialized. It is also useful to enlarge the pool size of prospective events for screening by increasing the percentage of  $R_0$  events (primary regenerated plants) with fertility. As plant transformation is labor intensive, any method that decreases the number of  $R_0$  events that must be produced in order to obtain a transgenic event with appropriate performance and growth characteristics is also valuable.

Large populations of independent transgene  $R_0$  insertion events of the non-targeted pMON26800 and pMON30463 vectors, and the targeted pMON30464 vector, were generated and scored for fertility. It was observed that a higher percentage of the  $R_0$  events generated with the targeted vector were fertile (Table 8). Progeny of fertile  $R_0$  events were subsequently introduced into field tests where they were scored for European corn borer resistance (ECB1) and normal segregation.

Methods for determination of ECB1 ratings and segregation values were essentially as described (Armstrong *et al.*, 1995). Events that passed the ECB1 and segregation criteria were subsequently scored for stunting or height reductions. While 60% of the non-targeted events displayed height reductions, only 3% of the targeted events were stunted (Table 8). Improved fertility and reduced stunting resulted in significantly improved (37% vs. 8%) recovery of unstunted ECB1 positive events with the targeted Cry2Ab vector. In summary, 4-fold more non-targeted R<sub>0</sub> events must be produced and screened to obtain the same number of normal, ECB+ R<sub>0</sub> events obtained with the targeted Cry2Ab vector in a transformation study.

Table 8

Comparisons of Percentage of Fertile, Stunted, and Normal Maize Plants Obtained with Untargeted and Targeted Cry2Ab Expression Vectors

Vector	# ECB LD + R <sub>0</sub> Events <sup>a</sup>	% Fertile Events <sup>b</sup>	% Stunted <sup>c</sup>	% Normal, ECB1 + <sup>d</sup>
Untargeted	192	66	63	7
Targeted	78	85	4	31

<sup>a</sup>#ECB LD + R<sub>0</sub> events are the # of R<sub>0</sub> events that were positive by an ECB leaf disk feeding assay.

<sup>b</sup>% of the ECB LD+ R<sub>0</sub> events yielding viable R1 progeny (seed).

<sup>c</sup>% Stunted is the % of the ECB1 positive and properly segregating events with reduced stature. (Total ECB1 positive and properly segregating for non-targeted was 38; for targeted was 25).

<sup>d</sup>4)% normal, ECB1 + is the % normal, ECB+ events obtained relative to the total number of ECB LD+ R<sub>0</sub> events screened.

### 5.3 Example 3 - Plastid Targeting of Cry2Ab Increases Frequency of High Level European Corn Borer Control in Transgenic Corn

The previously described populations of independently transformed events derived from both targeted (pMON30464) and non-targeted (pMON30463 or pMON26800) Cry2Ab expression vectors were also screened for resistance to second generation European corn borer infestations (ECB2). To facilitate these studies, the commercially efficacious transgenic corn event MON810 (Yieldgard™) transformed with the Cry1Ab gene was included as a positive control. Efficacy against ECB2 was tested in field tests essentially as described (Armstrong *et al.*, 1995). In the 1996 field test, 18 independent non-targeted pMON26800 events were compared to MON810 (Cry1Ab). Of these 18 events, only one delivered ECB2 protection that was both statistically indistinguishable from MON810 and significantly less than the non-transgenic negative control (event UT1 in Table 9). In the 1997 field test, 18 independent targeted events (pMON30464) were tested in parallel with 3 non-targeted events (1 pMON30463 event and the two pMON26800 events derived from the 1996 tests) and MON810 (Table 10). Nine of the eighteen targeted pMON30464 events delivered ECB2 protection that was statistically indistinguishable from ECB2 protection conferred by the commercially efficacious Cry1Ab-expressing

MON810 (Yieldgard™) event and all had significantly less ECB2 damage than the non-transgenic negative control (Table 10).

These data sets indicate that the absolute number and frequency of commercially efficacious Cry2Ab lines obtained from the targeted pMON30464 vector is much greater than that obtained from the non-targeted pMON26800 vector. While 9 of 18 targeted Cry2Ab events (50%) delivered ECB2 control that was both statistically indistinguishable from the MON810 Cry1Ab commercial standard and significantly less than the non-transgenic negative control, only 1 of 18 non-targeted Cry2Ab events (6%) displayed ECB2 control that was both statistically indistinguishable from the MON810 *cry1Ab* commercial standard and significantly less than the non-transgenic negative control. The superiority of the targeted Cry2Ab expression vector is especially evident if one considers that 9 commercially efficacious Cry2Ab events were obtained from a total of 78 ECB leaf disk positive R<sub>0</sub> plants for an 11.5% frequency of recovery while only 3 commercially efficacious Cry2Ab events were obtained from a total of 192 ECB leaf disk feeding positive R<sub>0</sub>s for a 1.6% recovery frequency (R<sub>0</sub> ECB data from Table 6).

**Table 9**  
Comparison of ECB2 Protection in Untargeted (UT) Cry2Ab Transgenic Corn Relative to MON810 Cry1Ab Yieldgard™ Transgenic Corn in Field Tests

Event	Sample Size	Stalk Tunneling (inches)
MON810 (+ ctrl.)	20	0.3 <sup>a</sup>
UT1	10	0.7 <sup>a</sup>
UT2	10	1.9 <sup>a</sup>
UT3	10	2.0 <sup>a</sup>
UT4	10	2.5 <sup>b</sup>
UT5	8	2.6 <sup>b</sup>
UT6	10	2.9 <sup>b</sup>
UT7	10	3.1 <sup>b</sup>
UT8	10	3.4 <sup>b</sup>
UT9	10	3.4 <sup>b</sup>
UT10	10	3.5 <sup>b</sup>
UT11	4	3.6 <sup>b</sup>
Wild type	10	3.7 <sup>b</sup>
UT12	10	3.8 <sup>b</sup>
UT13	10	4.6 <sup>b</sup>
UT14	10	5.8 <sup>b</sup>
UT15	10	6.8 <sup>c</sup>
UT16	10	7.6 <sup>c</sup>
UT17	10	9.3 <sup>c</sup>
UT18	10	10.1 <sup>c</sup>

<sup>a, b</sup> Values marked with the same superscript (a) are statistically indistinguishable from MON810 in planned comparisons at  $P = 0.05$ . Values with superscripts (b) are statistically distinct. Events with stalk tunneling values significantly greater than the Cry1Ab commercial standard MON810 are shown in boldface. Genetic background of all events is identical (B73 × H99).

5 <sup>c, \*</sup> Values marked with an asterisk are significantly lower than the wild-type non-transgenic negative control in planned comparisons with the negative control ( $P = 0.05$ ). Values marked with superscript (c) are significantly greater than the wild-type non-transgenic negative control in planned comparisons with the negative control ( $P = 0.05$ ). UT1-UT18: Untargeted pMON26800 events #1-18.

Table 10

Comparison of ECB2 Protection in Targeted (T) and Untargeted (UT) Cry2Ab Transgenic Corn Relative to MON810 Cry1Ab Yieldgard<sup>TM</sup> Transgenic Corn in Field Tests

Event	Sample Size	Stalk Tunneling (inches)
T1	9	0.6 <sup>a</sup>
T2	10	0.6 <sup>a</sup>
MON810 (+ ctrl.)	30	0.9 <sup>a</sup>
T3	14	1 <sup>a</sup>
T4	12	1.3 <sup>a</sup>
T5	7	1.4 <sup>a</sup>
<u>UT1</u>	<u>10</u>	<u>1.6<sup>a</sup></u>
T6	13	1.6 <sup>a</sup>
T7	11	1.6 <sup>a</sup>
T8	10	1.7 <sup>a</sup>
<u>UT2</u>	<u>10</u>	<u>1.8<sup>a</sup></u>
T9	10	2.4 <sup>a</sup>
T10	12	2.5 <sup>b</sup>
T11	7	2.6 <sup>b</sup>
T12	9	2.9 <sup>b</sup>
T13	10	3.2 <sup>b</sup>
T14	11	3.3 <sup>b</sup>
T15	10	3.5 <sup>b</sup>
T16	10	4.0 <sup>b</sup>
T17	10	4.3 <sup>b</sup>
<u>UT3</u>	<u>8</u>	<u>4.8<sup>b</sup></u>
T18	8	5.4 <sup>b</sup>
wild type (- ctrl.)	20	13.7 <sup>c</sup>

<sup>a, b, c</sup>Values marked with the superscript (a) are statistically indistinguishable from MON810 in planned comparisons at  $P = 0.05$ . Values with superscripts are statistically distinct. Events with stalk tunneling values significantly greater than the Cry1Ab commercial standard MON810 are shown in boldface; all transgenic events display significantly less tunneling than the wild type non-transgenic negative control in planned comparisons to the negative control ( $P = 0.5$ ). Genetic background of all events is identical (B73  $\times$  H99).

T1-T18: Targeted pMON30464 events #1-18. UT1-UT3: Untargeted pMON30463 and pMON26800 events #1-3. UT1 in the 1997 field test is the same pMON26800 event as UT3 in the 1996 field test.

#### 5.4 Example 4 - Plastid Targeting of the Cry2Ab Protein Results in Increased Expression in Transgenic Cotton Callus Tissue

Levels of the Cry2Ab protein in cotton callus tissue transformed with plastid-targeted and non-targeted Cry2Ab expression vectors were compared. Cry2Ab levels were significantly higher in callus  
5 that had been transformed with the plastid-targeted genes (Table 11).

Plant expression vector pMON33830 contained a Cry2Ab expression cassette consisting of the following genetic elements operably linked to produce functional Cry2Ab protein in plant cells: an enhanced CaMV 35S promoter, a petunia hsp70 5' untranslated leader, a synthetic *cry2Ab* gene with a translation initiation codon (SEQ ID NO:1), and transcription termination and polyadenylation sequences  
10 from the nopaline synthase (NOS) gene of *A. tumefaciens*.

Plant expression vectors pMON33827 (SEQ ID NO:13), pMON33828 (SEQ ID NO:14) and pMON33829 (SEQ ID NO:15) contained Cry2Ab expression cassettes similar to that occurring in pMON33830 except that in each a different chloroplast targeting sequence was translationally fused to the N-terminus of the synthetic *cry2Ab* gene. pMON33827 contained the coding sequence for PTP1 (SEQ  
15 ID NO:5) which consists of an *Arabidopsis thaliana* ssRUBISCO (SSU) chloroplast targeting sequence and sequences coding for the first 24 amino acids of ssRUBISCO (SSU) protein (Wong *et al.*, 1992). SEQID NO:6 represents the PTP1 targeting peptide sequence. This peptide contains the complete native targeting sequence including the plastid targeting peptide cleavage site along with the first twenty-four amino acids of the mature RUBISCO SSU protein sequentially linked to a duplicated sequence of amino  
20 acids (SEQID NO:6 amino acids position No. 50-57) containing the RUBISCO SSU plastid targeting peptide cleavage site (SEQID NO:6 amino acids position No. 80-87). PTP1 therefor contains a duplicated plastid targeting peptide cleavage site. The polynucleotide cassette containing this PTP coding sequence is linked at its 3' end to an *NcoI* restriction site which allows for insertions of coding sequences which are translationally in-frame with the PTP coding sequence, for example, those which encode  
25 Cry2Ab, Cry2Aa, variants of these, and other useful polypeptide encoding sequences.



pMON33828 contained the coding sequence for PTP1Δ (SEQ ID NO:7), a modification of PTP1 in which the 24 amino acids of SSU between the two transit peptide cleavage sites was removed by cleavage with the restriction enzyme *SphI*, which cuts once within each copy of the transit peptide cleavage site, and re-ligation, resulting in the presence of only the transit peptide portion of PTP1 followed by a single copy of the transit peptide cleavage site and an *NcoI* site. The peptide sequence for PTP1Δ is designated SEQ ID NO:8.

pMON33829 contained the coding sequence for PTP2 (SEQ ID NO:9), the transit peptide sequence from the EPSP synthase gene of *Arabidopsis thaliana*. The peptide sequence for PTP2 is designated SEQ ID NO:10.

All of the above plant transformation expression vectors also contained a selectable marker gene cassette which confers kanamycin resistance to transformed plant cells.

Cotton callus tissue from 12 randomly chosen, independent transgenic events from transformations with each of pMON33827, pMON33828, pMON33829 and pMON33830 was subjected to quantitative analysis of Cry2Ab protein levels using a quantitative ELISA assay. This ELISA used a direct sandwich technique that used a monoclonal capture antibody raised against Cry2Aa, a different Cry2Aa monoclonal antibody conjugated to alkaline phosphatase as the secondary antibody, and purified Cry2Aa protein as a standard. Comparison of Cry2Ab expression levels in targeted and non-targeted callus tissue showed a significant increase in expression when a chloroplast targeting sequence was included (Table 11). PTP1Δ provided a significantly greater mean expression level when compared to non-targeted Cry2Ab as determined by applying a t test ( $t = 2.31$ ,  $p = 0.03$ ). PTP2 provided a significantly greater probability of obtaining callus lines expressing higher levels of Cry2Ab as determined by applying a G test ( $G^2/X^2 = 5.6$ ,  $p = 0.02$ ).

Table 11

Cry2Ab Levels in Independent Transformed Cotton Callus Lines Comparing Chloroplast-Targeted and Untargeted *cry2Ab* genes

Cotton Callus Lines	Cry2Ab ng/mL of Callus Extract
Non-transformed callus	
Line 1	0
Line 2	0
Line 3	0
Line 4	0
pMON33827, PTP1- <i>cry2Ab</i> gene	
Line 1	464
Line 2	61

Cotton Callus Lines	Cry2Ab ng/mL of Callus Extract
Line 3	0
Line 4	25
Line 5	0
Line 6	368
Line 7	74
Line 8	101
Line 9	20
Line 10	652
Line 11	0
Line 12	0
pMON33828, PTP1Δ-cry2Ab Gene	
Line 1	252
Line 2	235
Line 3	0
Line 4	416
Line 5	0
Line 6	0
Line 7	0
Line 8	101
Line 9	393
Line 10	587
Line 11	788
Line 12	277
pMON33829, PTP2-cry2Ab Gene	
Line 1	60
Line 2	0
Line 3	2220
Line 4	2036
Line 5	0
Line 6	38
Line 7	674
Line 8	2440
Line 9	15
Line 10	91

Cotton Callus Lines	Cry2Ab ng/mL of Callus Extract
Line 11	290
Line 12	71
pMON33830, <i>cry2Ab</i> Gene	
Line 1	19
Line 2	166
Line 3	47
Line 4	20
Line 5	33
Line 6	47
Line 7	781
Line 8	35
Line 9	31
Line 10	0
Line 11	0
Line 12	136

### 5.5 Example 5 - Targeting the Cry2Aa Protein to Plastids Results in Decreased Expression in Transgenic Cotton Callus Tissue

In contrast to Example 4 above, and exemplifying that the increase in expression obtained using plastid targeting sequences is specific to particular *cry* genes, the inventors discovered that the same plastid targeting sequences described above, PTP1, PTP1 $\Delta$  and PTP2, resulted in significantly lower levels of expression of the closely related *cry2Aa* gene in transgenic cotton callus (Table 12). Plant expression vector pMON33803 contained a *cry2Aa* expression cassette consisting of the following genetic elements operably linked to produce functional Cry2Aa protein in plant cells: a FMV35S promoter, a petunia heat shock HSP70 5' untranslated leader, a synthetic *cry2Aa* gene (SEQ ID NO:11) with a translation initiation codon and *Nco*I restriction enzyme site at the 5'-end, and transcription termination and polyadenylation sequences from the E9 SSU gene from pea. The peptide sequence for the Cry2Aa protein is designated SEQ ID NO:12. pMON33812, pMON33811, and pMON33806 contained *cry2Aa* expression cassettes similar to that occurring in pMON33803 except that in each case a different chloroplast targeting sequence (PTP1, PTP1 $\Delta$ , and PTP2, respectively) was transitionally fused to the N-terminus of the synthetic *cry2Aa* gene. All of these vectors also contained a selectable marker gene cassette conferring glyphosate resistance to transformed plant cells.

Cotton callus tissue from 10 randomly chosen independent transgenic events from transformations with each of pMON33803, pMON33812, pMON33811 and pMON33806 was subjected

to quantitative analysis of Cry2Aa protein levels using the quantitative Cry2 ELISA assay. Comparison of Cry2Aa expression levels in targeted and non-targeted callus tissue showed a significant decrease in expression when chloroplast targeting sequences were included (Table 12). The non-targeted *cry2Aa* gene conferred expression levels that differed significantly from those achieved using any of the three plastid targeted *cry2Aa* genes, as determined by using a Tukey-Kramer HSD test ( $\alpha = 0.05$ ).

**Table 12**  
**Cry2Aa Levels In Independent Transformed Cotton Callus Lines Comparing Chloroplast-Targeted And Untargeted Cry2Aa Genes**

Cotton Callus Lines	Cry2Aa ng/mL of Extract
<b>Non-transformed callus</b>	
Line 1	0
Line 2	0
Line 3	0
Line 4	0
<b>pMON33812, PTP1-Cry2Aa Gene</b>	
Line 1	29
Line 2	32
Line 3	22
Line 4	41
Line 5	24
Line 6	47
Line 7	43
Line 8	49
Line 9	0
Line 10	23
<b>pMON33811, PTP1Δ-Cry2Aa Gene</b>	
Line 1	0
Line 2	59
Line 3	48
Line 4	72
Line 5	29
Line 6	37
Line 7	44
Line 8	32
Line 9	20

Cotton Callus Lines	Cry2Aa ng/mL of Extract
Line 10	0
pMON33806, PTP2-Cry2Aa Gene	
Line 1	27
Line 2	0
Line 3	10
Line 4	84
Line 5	205
Line 6	0
Line 7	13
Line 8	6
Line 9	0
Line 10	8
pMON33803, Cry2Aa Gene	
Line 1	63
Line 2	2278
Line 3	181
Line 4	3131
Line 5	3752
Line 6	851
Line 7	303
Line 8	1365
Line 9	1601
Line 10	1648

#### 5.6 Example 6 - Targeting the Cry2Aa Protein to Plastids Results in Decreased Expression and Increased Phytotoxicity in Transgenic Tobacco Plants

Transformed tobacco plants were generated using pMON33803, the non-targeted *cry2Aa* plant expression vector and pMON33806, the chloroplast-targeted PTP2-*cry2Aa* plant expression vector. Leaf tissue samples of equivalent weight from 48 pMON33803 plants and 41 pMON33806 plants were extracted in equal volumes of extraction buffer and the relative levels of *cry2Aa* were determined using a qualitative ELISA (Table 13). This ELISA used a direct sandwich technique that used polyclonal capture antibody raised against Cry2Aa, the same polyclonal antibody conjugated to alkaline phosphatase as the secondary antibody, and purified Cry2Aa protein as a standard.

The proportion of the total number of plants recovered from transformation that expressed non-targeted Cry2Aa at high levels was greater than the proportion of plants recovered that expressed targeted

Cry2Aa at high levels. Conversely, the proportion of the total number of plants recovered from transformation that failed to express detectable targeted Cry2Aa was greater than the proportion of plants recovered that failed to express non-targeted Cry2Aa. All of the PTP2-Cry2Aa plants that had detectable levels of Cry2Aa expression exhibited a severely abnormal phenotype; these plants were extremely stunted, had shortened internodes, had deformed, wrinkled leaves, and were infertile. All of the PTP2-Cry2Aa plants that lacked Cry2Aa expression appeared normal. In contrast, only some of the high expressing non-targeted Cry2Aa plants displayed a stunted phenotype.

**Table 13**  
Cry2Aa Levels in Independent Transformed Tobacco Plants Comparing  
Chloroplast-Targeted and Untargeted *cry2Aa* Genes

Transgenic Plants	ELISA O.D.	Transgenic Plants	ELISA O.D.
pMON33803_ <i>cry2Aa</i> gene		pMON33806 PTP2- <i>cry2Aa</i> gene	
Plant 1	2.5	Plant 1	0
Plant 2	1.1	Plant 2	0
Plant 3	2.7	Plant 3	0
Plant 4	0.1	Plant 4	0
Plant 5	0.1	Plant 5	0
Plant 6	2.3	Plant 6	0.9
Plant 7	1.9	Plant 7	0.4
Plant 8	2.4	Plant 8	0.4
Plant 9	0	Plant 9	0
Plant 10	2.1	Plant 10	0.6
Plant 11	0.1	Plant 11	0
Plant 12	0.5	Plant 12	0.5
Plant 13	2.4	Plant 13	0.4
Plant 14	0.1	Plant 14	0.7
Plant 15	2.2	Plant 15	1.5
Plant 16	0.2	Plant 16	0.6
Plant 17	2.6	Plant 17	0
Plant 18	2.5	Plant 18	0
Plant 19	2.5	Plant 19	0
Plant 20	1.4	Plant 20	0
Plant 21	2.4	Plant 21	0
Plant 22	2.1	Plant 22	0

Transgenic Plants	ELISA O.D.	Transgenic Plants	ELISA O.D.
Plant 23	0.5	Plant 23	0.6
Plant 24	2.1	Plant 24	0
Plant 25	0.3	Plant 25	0
Plant 26	0	Plant 26	0.6
Plant 27	0.3	Plant 27	0
Plant 28	2.2	Plant 28	0.7
Plant 29	0	Plant 29	0.5
Plant 30	1.5	Plant 30	0
Plant 31	0.1	Plant 31	0
Plant 32	0.1	Plant 32	0
Plant 33	0.7	Plant 33	0
Plant 34	0	Plant 34	0
Plant 35	0	Plant 35	0
Plant 36	0	Plant 36	0
Plant 37	0.2	Plant 37	0
Plant 38	2.1	Plant 38	0
Plant 39	0	Plant 39	0
Plant 40	1.9	Plant 40	0
Plant 41	1.5	Plant 41	0
Plant 42	2.8		
Plant 43	0.6		
Plant 44	2.1		
Plant 45	0.9		
Plant 46	0		
Plant 47	0		
Plant 48	0		

### 5.7 Example 7 - Transformation of Tobacco Chloroplast with a Cry2Ab gene

Recombinant plants can be produced in which only the mitochondrial or chloroplast DNA has been altered to incorporate the molecules envisioned in this application. Promoters which function in chloroplasts have been known in the art (Hanley-Bowden et al., Trends in Biochemical Sciences **12**:67-70, 1987). Methods and compositions for obtaining cells containing chloroplasts into which heterologous DNA has been inserted have been described, for example by Daniell et al. (U.S. Pat. No. 5,693,507; 1997) and Maliga et al. (U.S. Pat. No. 5,451,513; 1995). A vector can be constructed which contains an expression cassette from which a Cry2A protein could be produced. A cassette could contain a

chloroplast operable promoter sequence driving expression of a *cry2A* crystal protein gene, constructed in much the same manner as other polynucleotides herein, using thermal amplification methodologies, restriction endonuclease digestion, and ligation etc. A chloroplast expressible gene would provide a promoter and a 5' untranslated region from a heterologous gene or chloroplast gene such as *psbA*, which would provide for transcription and translation of a DNA sequence encoding a Cry2A protein in the chloroplast; a DNA sequence encoding Cry2A protein; and a transcriptional and translational termination region such as a 3' inverted repeat region of a chloroplast gene that could stabilize an expressed *cry2A* mRNA. Expression from within the chloroplast would enhance *cry2A* gene product accumulation. A host cell containing chloroplasts or plastids can be transformed with the expression cassette and then the resulting cell containing the transformed chloroplasts can be grown to express the Cry2A protein. A cassette may also include an antibiotic, herbicide tolerance, or other selectable marker gene in addition to the *cry2A* gene. The expression cassette may be flanked by DNA sequences obtained from a chloroplast DNA which would facilitate stable integration of the expression cassette into the chloroplast genome, particularly by homologous recombination. Alternatively, the expression cassette may not integrate, but by including an origin of replication obtained from a chloroplast DNA, would be capable of providing for replication of the heterologous *cry2A* gene in the chloroplast. Plants can be generated from cells containing transformed chloroplasts and can then be grown to produce seeds, from which additional plants can be generated. Such transformation methods are advantageous over nuclear genome transformation, in particular where chloroplast transformation is effected by integration into the chloroplast genome, because chloroplast genes in general are maternally inherited. This provides environmentally "safer" transgenic plants, virtually eliminating the possibility of escapes into the environment. Furthermore, chloroplasts can be transformed multiple times to produce functional chloroplast genomes which express multiple desired recombinant proteins, whereas nuclear genomic transformation has been shown to be rather limited when multiple genes are desired. Segregational events are thus avoided using chloroplast or plastid transformation. Unlike plant nuclear genome expression, expression in chloroplasts or plastids can be initiated from only one promoter and continue through a polycistronic region to produce multiple peptides from a single mRNA.

The expression cassette would be produced in much the same way that other plant transformation vectors are constructed. Plant chloroplast operable DNA sequences can be inserted into a bacterial plasmid and linked to DNA sequences expressing desired gene products, such as Cry2A proteins, so that Cry2A protein is produced within the chloroplast, obviating the requirement for nuclear gene regulation, capping, splicing, or polyadenylation of nuclear regulated genes, or chloroplast or plastid targeting sequences. An expression cassette comprising a *cry2A* gene, which is either synthetically constructed or a native gene derived directly from a *B. thuringiensis* genome or a *B. thuringiensis* episomal element, would be inserted into a restriction site in a vector constructed for the purpose of chloroplast or plastid transformation. The cassette would be flanked upstream by a chloroplast or plastid functional promoter



and downstream by a chloroplast or plastid functional transcription and translation termination sequence. The resulting cassette could be incorporated into the chloroplast or plastid genome using well known homologous recombination methods.

Alternatively, chloroplast or plastid transformation could be obtained by using an autonomously replicating plasmid or other vector capable of propagation within the chloroplast or plastid. One means of effectuating this method would be to utilize a portion of the chloroplast or plastid genome required for chloroplast or plastid replication initiation as a means for maintaining the plasmid or vector in the transformed chloroplast or plastid. A sequence enabling stable replication of a chloroplast or plastid epigenetic element could easily be identified from random cloning of a chloroplast or plastid genome into a standard bacterial vector which also contains a chloroplast or plastid selectable marker gene, followed by transformation of chloroplasts or plastids and selection for transformed cells on an appropriate selection medium. Introduction of an expression cassette as described herein into a chloroplast or plastid replicable epigenetic element would provide an effective means for localizing a Cry2A *B. thuringiensis*  $\delta$ -endotoxin to the chloroplast or plastid.

#### 6.0 References

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

- U. S. Patent No. 4,535,060; 1985
- U. S. Patent No. 4,554,101; 1985
- U. S. Patent No. 4,683,195; 1987
- U. S. Patent No. 4,683,202; 1987
- U. S. Patent No. 4,757,011; 1988
- U. S. Patent No. 4,769,061; 1988
- U. S. Patent No. 4,940,835; 1990
- U. S. Patent No. 4,971,908; 1990
- U. S. Patent No. 5,004,863; 1991
- U. S. Patent No. 5,015,580; 1991
- U. S. Patent No. 5,023,179; 1991
- U. S. Patent No. 5,225,341; 1993
- U. S. Patent No. 5,264,364; 1993
- U. S. Patent No. 5,276,269; 1994
- U. S. Patent No. 5,322,687; 1994
- U. S. Patent No. 5,338,544; 1994
- U. S. Patent No. 5,349,124; 1994
- U. S. Patent No. 5,378,619; 1995
- U. S. Patent No. 5,384,253; 1995

- U. S. Patent No. 5,416,011; 1995  
U. S. Patent No. 5,424,412; 1995  
U. S. Patent No. 5,451,513; 1995  
U. S. Patent No. 5,463,175; 1995  
5 U. S. Patent No. 5,482,852; 1996  
U. S. Patent No. 5,491,288; 1996  
U. S. Patent No. 5,500,365; 1996  
U. S. Patent No. 5,508,468; 1996  
U. S. Patent No. 5,569,834; 1996  
10 U. S. Patent No. 5,689,052; 1997  
U. S. Patent No. 5,693,507; 1997  
EPO 0120516  
WO 92/17591  
WO 95/24492  
15 Armstrong *et al.*, *Plant Cell Rep.*, 9:335-339, 1990.  
Armstrong *et al.*, *Crop Science*, 35(2):550-557, 1995.  
Barton *et al.*, *Plant Physiol.*, 85:1103-1109, 1987.  
Benfey *et al.*, *EMBO J.*, 8:2195-2202, 1989.  
Bevan *et al.*, *Nature*, 304:184, 1983.  
20 Callis *et al.*, *Genes Dev.*, 1: 1183-1200; 1987.  
Cheng *et al.*, *Proc. Natl. Acad. Sci. USA*, 96(6):2767-2772, 1998.  
Dhir *et al.*, *Plant Cell Rep.*, 10: 106-10; 1991  
Diehn *et al.*, In: *Genetic Engineering*, Ed. J.K. Setlow, Plenum Press, New York, NY, 18:83-99, 1996.  
Donovan *et al.*, *Mol. Gen. Genet.*, 214:365-372, 1988.  
25 Donovan *et al.*, *Appl. Environ. Microbiol.*, 58:3921-3927, 1992.  
English *et al.*, *Insect. Biochem. Molec. Bio.*, 24(10):1025-1026, (1994)  
Feinberg and Vogelstein, *Anal. Biochem.*, 132:6-13, 1983.  
Fischhoff *et al.*, *Bio/Technology*, 5:807-813, 1987.  
Frischauf *et al.*, *Methods Enzymol.*, 153:103-115, 1987.  
30 Fromm *et al.*, *Nature*, 319:791-793, 1986.  
Fromm *et al.*, *Bio/Technology*, 8:833-839, 1990.  
Gould *et al.*, *J. Cell Biol.*, 105:2923-2931, 1987.  
Hanley-Bowden *et al.*, *Trends in Biochemical Sciences* 12:67-70; 1987.  
Herrera-Estrella *et al.*, *Nature*, 303:209, 1983.  
35 Hertig *et al.*, *Plant Mol. Biol.*, 16:171-174, 1991.  
Hofte and Whiteley, *Microbiol. Rev.*, 53(2):242-255, 1989.

- Horsch *et al.*, *Science*, 227:1229-1231, 1985.
- Horton *et al.*, *Gene*, 77:61-68, 1989.
- Ishida *et al.*, *Nat. Biotechnol.*, 14(6):745-750, 1996.
- Kay *et al.*, *Science*, 236:1299-1302, 1987.
- 5 Keegstra and Olsen, *Ann. Rev. Plant Physiol. Mol. Biol.*, 40:471-501, 1989.
- Klee *et al.*, *Bio/Technology*, 3:637-642, 1985.
- Klee *et al.*, *Mol. Gen. Genet.*, 210:437-442, 1987.
- Kochler and Ho, *Plant Cell*, 2:769-783, 1990.
- Koziel *et al.*, *Bio/Technology*, 11:194-200, 1993.
- 10 Lambert *et al.*, *Appl. Environ. Microbiol.*, 62:80-86, 1996.
- Lee *et al.*, *Science*, 239:1288-1291, 1988.
- Lindstrom *et al.*, *Dev. Genet.*, 11:160-7;1990.
- Macejak and Sarnow, *Nature*, 353:90-94, 1991.
- MacIntosh *et al.*, *J. Invert. Pathol.*, 56:258-266, 1990.
- 15 Maliga, *Trends in Biotechnology*, 11:101-106, 1993.
- McElroy *et al.*, *Plant Cell*, 2:163-171, 1990.
- McGaughey and Whalon, *Science*, 258:1451-1455, 1993.
- Nayak *et al.*, *Proc. Natl. Acad. Sci. USA*, 94(6):2111-2116, 1997.
- Odell *et al.*, *Nature*, 313: 810-12; 1985.
- 20 Pelletier and Sonenberg, *Nature*, 334:320-325, 1988.
- Perlak *et al.*, *Bio/technology*, 8:939-943, 1990.
- Perlak *et al.*, *Plant Molecular Biol.*, 22:313-321, 1993.
- Roush, *Biocontrol Sci. Technol.*, 4:501-516, 1994.
- Russell *et al.*, *Plant Cell Reports*, 13:24-27, 1993.
- 25 Shelton *et al.*, *J. Econ. Entomol.*, 86:697-705, 1993.
- Tang *et al.*, *Appl. Environ. Microbiol.*, 62:564-569, 1996.
- Tillmann *et al.*, *EMBO J.*, 8(9):2463-2467, 1989.
- Vaeck *et al.*, *Nature*, 328:33-37, 1987.
- Vasil *et al.*, *Plant Physiol.*, 91:1575-1579; 1989.
- 30 Vodkin *et al.*, *Cell*, 34:1023-1031; 1983.
- Widner *et al.*, *J. Bacteriol.*, 171:965-974; 1989.
- Widner *et al.* (a), *J. Bacteriol.*, 172:2826-2832; 1990.
- Winter *et al.*, *Mol. Gen. Genet.*, 221(2):315-319, 1988.
- Wong *et al.*, *Plant Molec. Biol.*, 20:81-93, 1992.
- 35 Xu *et al.*, *Plant Mol. Biol.*, 27:237-248, 1995.
- Yamamoto *et al.*, *Plant Cell*, 3:371-382, 1991.

All of the compositions and methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

## Claims:

1. A plant comprising a nucleic acid sequence comprising a plant functional promoter sequence operably linked to a polynucleotide sequence encoding a Cry2A *Bacillus thuringiensis*  $\delta$ -endotoxin protein which lacks substantial Dipteran species inhibitory activity, wherein expression of said nucleic acid sequence in said plant yields said protein localized to a subcellular organelle or compartment.
2. A plant comprising a nucleic acid sequence comprising a plant functional promoter sequence operably linked to a first polynucleotide sequence encoding a plastid transit peptide, which is linked in-frame to a second polynucleotide sequence encoding a Cry2A *Bacillus thuringiensis*  $\delta$ -endotoxin protein which lacks substantial Dipteran species inhibitory activity, wherein said second polynucleotide is operably linked to a plant functional 3' end transcription termination and polyadenylation sequence, wherein expression of said nucleic acid sequence in said plant yields a fusion protein comprised of an amino-terminal plastid transit peptide covalently linked to said  $\delta$ -endotoxin protein, and wherein said fusion protein functions to localize said  $\delta$ -endotoxin protein to a subcellular organelle or compartment.
3. The plant of claim 1, wherein said polynucleotide sequence encoding a Cry2A *Bacillus thuringiensis*  $\delta$ -endotoxin protein comprises a sequence encoding a Cry2Ab *Bacillus thuringiensis*  $\delta$ -endotoxin protein.
4. The plant of claim 2, wherein said polynucleotide sequence encoding a Cry2A *Bacillus thuringiensis*  $\delta$ -endotoxin protein comprises a sequence encoding a Cry2Ab *Bacillus thuringiensis*  $\delta$ -endotoxin protein.
5. The plant of claim 1, wherein said subcellular organelle or compartment is a plant plastid or chloroplast.
6. The plant of claim 2, wherein said subcellular organelle or compartment is a plant plastid or chloroplast.
7. The plant of claim 1, wherein said nucleic acid sequence is introduced into and stably maintained within a plant plastid or chloroplast.
8. A plant tissue derived from progeny of the plant according to claim 5, wherein said plant tissue comprises a plant, plant seed, or plant cells containing said polynucleotide sequence encoding said  $\delta$ -endotoxin protein.
9. A plant tissue derived from progeny of the plant according to claim 6, wherein said plant tissue comprises a plant, plant seed, or plant cells containing said polynucleotide sequence encoding said  $\delta$ -endotoxin protein.
10. The plant according to claim 1, wherein said nucleic acid sequence comprising a

promoter is a plant chloroplast or plastid functional promoter.

11. The plant according to claim 2, wherein said nucleic acid sequence comprising a plant functional promoter is a promoter sequence which is naturally expressed in plants.
12. The plant according to claim 1, wherein said polynucleotide sequence  
5 encoding a Cry2A *Bacillus thuringiensis*  $\delta$ -endotoxin protein is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:11, and SEQ ID NO:17.
13. The plant according to claim 2, wherein said polynucleotide sequence encoding a Cry2A *Bacillus thuringiensis*  $\delta$ -endotoxin protein is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:11, and SEQ ID NO:17.
- 10 14. The plant according to claim 1, wherein said Cry2Ab *Bacillus thuringiensis*  $\delta$ -endotoxin protein is selected from the group consisting of SEQ ID NO:2, SEQ ID NO:12, and SEQ ID NO:18.
15. The plant according to claim 2, wherein said Cry2Ab *Bacillus thuringiensis*  $\delta$ -endotoxin protein is selected from the group consisting of SEQ ID NO:2, SEQ ID NO:12, and SEQ  
15 ID NO:18.
16. The plant of claim 2, wherein said nucleic acid sequence further comprises a plant functional intron sequence.
17. The plant of claim 16, wherein said intron sequence is selected from the group consisting of Adh intron 1, sucrose synthase intron, TMV omega element, maize Heat Shock Protein  
20 70 intron, and the rice Act1 intron.
18. The plant of claim 16, wherein said intron sequence is the maize Heat Shock Protein 70 intron.
19. The plant of claim 2, wherein first polynucleotide sequence encodes a plastid transit peptide selected from the group consisting of zmSSU PTP, PTP1, PTP1 $\Delta$ , and PTP2.
- 25 20. The plant of claim 19, wherein said zmSSU PTP plastid transit peptide comprising SEQ ID NO:4 is encoded by the nucleic acid sequence comprising SEQ ID NO:3.
21. The plant of claim 19, wherein said PTP1 plastid transit peptide comprising SEQ ID NO:6 is encoded by the nucleic acid sequence comprising SEQ ID NO:5.
22. The plant of claim 19, wherein said PTP1 $\Delta$  plastid transit peptide comprising SEQ ID  
30 NO:8 is encoded by the nucleic acid sequence comprising SEQ ID NO:7.
23. The plant of claim 19, wherein said PTP2 plastid transit peptide comprising SEQ ID NO:10 is encoded by the nucleic acid sequence comprising SEQ ID NO:9.
24. The plant of claim 2, comprising nucleotides 17 to 3182 of SEQ ID NO:13.
25. The plant of claim 2, comprising nucleotides 17 to 3092 of SEQ ID NO:14.
- 35 26. The plant of claim 2, comprising nucleotides 17 to 3155 of SEQ ID NO:15.

27. The plant of claim 1, wherein the plant is a monocotyledonous plant.
28. The plant of claim 2, wherein the plant is a monocotyledonous plant.
29. The plant of claim 27, wherein the plant is a monocotyledonous plant selected from the group consisting of maize, rice, wheat, barley, oats, rye, millet, sorghum, sugarcane, and turfgrass.
30. The plant of claim 28, wherein the plant is a monocotyledonous plant selected from the group consisting of maize, rice, wheat, barley, oats, rye, millet, sorghum, sugarcane, and turfgrass.
31. The plant of claim 1, wherein said plant is a dicotyledonous plant.
32. The plant of claim 2, wherein said plant is a dicotyledonous plant.
33. The plant of claim 31, wherein the plant is a dicotyledonous plant selected from the group consisting of cotton, soybean, tomato, potato, citrus, tobacco, canola, and strawberry.
34. The plant of claim 32, wherein the plant is a dicotyledonous plant selected from the group consisting of cotton, soybean, tomato, potato, citrus, tobacco, canola, and strawberry.
35. The plant of claim 1, further comprising an  $R_0$  transgenic plant.
36. The plant of claim 2, further comprising an  $R_0$  transgenic plant.
37. A progeny plant of any generation of the plant of claim 35, wherein said plant has inherited said nucleic acid sequence from said  $R_0$  transgenic plant.
38. A progeny plant of any generation of the plant of claim 36, wherein said plant has inherited said nucleic acid sequence from said  $R_0$  transgenic plant.
39. The plant according to claim 1, wherein said plant further comprises an additional nucleic acid sequence comprising a plant operable promoter linked to a polynucleotide sequence encoding a CryI *B. thuringiensis*  $\delta$ -endotoxin protein.
40. The plant according to claim 2, wherein said plant further comprises an additional nucleic acid sequence comprising a plant operable promoter linked to a polynucleotide sequence encoding a CryI *B. thuringiensis*  $\delta$ -endotoxin protein.
41. A method of producing a transgenic progeny plant comprising:
- (a) obtaining a first plant containing a nucleic acid sequence comprising a plant functional promoter operably linked to a first polynucleotide sequence encoding a plastid transit peptide, which is linked in frame to a second polynucleotide sequence encoding a Cry2A *Bacillus thuringiensis*  $\delta$ -endotoxin protein lacking substantial Dipteran species inhibitory activity, wherein said second polynucleotide is operably linked to a plant functional 3' end transcription termination and polyadenylation sequence, wherein expression of said nucleic acid sequence in said plant yields a fusion protein comprising an amino-terminal

plastid transit peptide covalently linked to said  $\delta$ -endotoxin protein, and wherein said fusion protein functions to localize said  $\delta$ -endotoxin protein to a subcellular organelle or compartment;

(b) obtaining a second plant; and

5 (c) crossing said first and second plants to obtain a crossed transgenic progeny plant, said progeny plant having inherited said nucleic acid sequence from said first plant.

42. The method of claim 41, wherein said progeny plant is a monocotyledonous plant, said monocotyledonous plant being selected from the group consisting of maize, rice, wheat, barley, oats, rye, millet, sorghum, sugarcane, and turfgrass.

10 43. The method of claim 41, wherein said progeny plant is a dicotyledonous plant, said dicotyledonous plant being selected from the group consisting of cotton, soybean, tomato, potato, citrus, and tobacco.

44. A nucleic acid sequence comprising a promoter operably linked to a first polynucleotide sequence encoding a plastid transit peptide, which is linked in frame to a second  
15 polynucleotide sequence encoding a Cry2A *Bacillus thuringiensis*  $\delta$ -endotoxin protein lacking substantial Dipteran inhibitory activity, wherein expression of said nucleic acid sequence by a plant cell produces a fusion protein comprising an amino-terminal plastid transit peptide covalently linked to said  $\delta$ -endotoxin protein, and wherein said fusion protein functions to localize said  $\delta$ -endotoxin protein to a subcellular organelle or  
20 compartment.

45. The nucleic acid sequence of claim 44, wherein said second polynucleotide sequence encodes a Cry2Ab *Bacillus thuringiensis*  $\delta$ -endotoxin protein.

46. The nucleic acid sequence of claim 45, wherein said second polynucleotide sequence encodes a Cry2Ab *Bacillus thuringiensis*  $\delta$ -endotoxin protein selected from the group of  
25 sequences consisting of SEQ ID NO:2 and SEQ ID NO:18.

47. The nucleic acid sequence of claim 46, wherein said second polynucleotide sequence is selected from the group of sequences consisting of SEQ ID NO:1 and SEQ ID NO:17.

48. A plant cell comprising a nucleic acid sequence comprising a promoter operably linked to a polynucleotide sequence encoding a Cry2A *Bacillus thuringiensis*  $\delta$ -endotoxin  
30 protein lacking substantial Dipteran inhibitory activity, wherein expression of said nucleic acid sequence in said plant yields said protein localized to a subcellular organelle or compartment.

49. The plant cell of claim 48, wherein said polynucleotide sequence encoding a Cry2A *Bacillus thuringiensis*  $\delta$ -endotoxin protein encodes a Cry2Ab *Bacillus thuringiensis*  $\delta$ -  
35 endotoxin protein.



50. The plant cell of claim 49, wherein said subcellular organelle or compartment is a plant plastid or chloroplast.
51. The plant cell of claim 48, wherein said nucleic acid sequence is introduced into and stably maintained within a plant plastid or chloroplast.
- 5 52. The plant cell of claim 51 wherein said nucleic acid sequence is expressed within said plastid or chloroplast, said expression producing an insecticidally effective amount of said  $\delta$ -endotoxin protein localized to said plastid or chloroplast.
53. A plant tissue derived from progeny of the plant cell according to claim 48, wherein said plant tissue comprises a plant, plant seed, plant cells or progeny tissues thereof  
10 containing said polynucleotide sequence expressing said  $\delta$ -endotoxin localized to a plant plastid or chloroplast.
54. The plant cell according to claim 50, wherein said nucleic acid sequence comprising a promoter is a plant chloroplast or plastid functional promoter.
55. The plant cell according to claim 54, wherein said polynucleotide sequence encoding a  
15 Cry2A *Bacillus thuringiensis*  $\delta$ -endotoxin is selected from the group consisting of SEQ ID NO:1, SEQ ID NO:11, and SEQ ID NO:17.
56. The plant cell according to claim 55, wherein said Cry2Ab *Bacillus thuringiensis*  $\delta$ -endotoxin is selected from the group consisting of SEQ ID NO:2, SEQ ID NO:12, and  
20 SEQ ID NO:18.




FIG. 1A




FIG. 1B

2/7




FIG. 2A




FIG. 2B




FIG. 3A




FIG. 3B




FIG. 4




FIG. 5




FIG. 6




FIG. 7



## SEQUENCE LISTING

<110> David R. Corbin  
Charles P. Romano

<120> Improved Method for Transforming Plants to Express  
delta-Endotoxins

<130> 38-21(13547)

<140> Application No. 09/186,002

<141> Filed 11/04/98

<160> 18

<170> FastSEQ for Windows Version 3.0

<210> 1

<211> 1934

<212> DNA

<213> Artificial Sequence

<220>

<223> Completely Synthesized

<400> 1

ccatggacaa	ctccgtcctg	aactctggtc	gcaccaccat	ctgcgacgcc	tacaacgtcg	60
cggcgcgatga	tccattcagc	ttccagcaca	agagcctcga	cactgttcag	aaggagtggga	120
cggagtggaa	gaagaacaac	cacagcctgt	acctggaccc	catcgtcggc	acgggtggcca	180
gcttccttct	caagaaggtc	ggctctctcg	tcgggaagcg	catcctctcg	gaactccgca	240
acctgatctt	tccatctggc	tccaccaacc	tcattgcaaga	catcctcagg	gagaccgaga	300
agtttctcaa	ccagcgctc	aacactgata	cccttgctcg	cgtcaacgct	gagctgacgg	360
gtctgcaagc	aaacgtggag	gagttcaacc	gccaaagtga	caacttcctc	aaccccaacc	420
gcaatgcggt	gcctctgtcc	atcacttctt	ccgtgaacac	catgcaacaa	ctgttcctca	480
accgcttgcc	tcagttccag	atgcaaggct	accagctgct	cctgctgcca	ctctttgctc	540
aggctgccaa	cctgcacctc	tccttcattc	gtgacgtgat	cctcaacgct	gacgagtggg	600
gcatctctgc	agccacgctg	aggacctacc	gcgactacct	gaagaactac	accagggact	660
actccaacta	ttgcatcaac	acctaccagt	cggccttcaa	gggcctcaat	acgaggcttc	720
acgacatgct	ggagttcagg	acctacatgt	tcctgaacgt	gttcgagtac	gtcagcatct	780
ggtcgctctt	caagtaccag	agcctgcttg	tgtccagcgg	cgccaacctc	tacgccagcg	840
gctctggtcc	ccaacaaact	cagagcttca	ccagccagga	ctggccattc	ctgtattcgt	900
tgttccaagt	caactccaac	tacgtctcca	acggcttctc	tgggtgctcg	ctctccaaca	960
cttcccccaa	cattgttggc	ctccccggct	ccaccacaac	tcattgctctg	cttgttgcca	1020
gagtgaacta	ctccggcggc	atctcgagcg	gcgacattgg	tgcattcgccg	ttcaaccaga	1080
acttcaactg	ctccaccttc	ctgccgcgcg	tgtccacccc	gttcgtgagg	tcctggctcg	1140
acagcggtc	cgaccgcgag	ggcgtggcca	ccgtcaccaa	ctggcaaac	gagtccttcg	1200
agaccacctt	tggcctccgg	agcggcgcc	tcacggcgcg	tgggaattct	aactacttcc	1260
ccgactactt	catcaggaac	atctctgggt	ttctctctgt	cgtccgcaac	gaggacctcc	1320
gccgtccact	gcactacaac	gagatcagga	acatcgcttc	tcctgctggg	acgcccggag	1380
gtgcaagggc	gtacatgggtg	agcgtccata	acaggaagaa	caacatccac	gctgtgcatg	1440
agaacggctc	catgatccac	ctggcgccca	atgattacac	cggcttcacc	atctctccaa	1500
tccacgccac	ccaagtgaac	aaccagacac	gcaccttcac	ctccgagaag	ttcggcaacc	1560
agggcgactc	cctgagggtc	gagcagaaca	acaccaccgc	caggtacacc	ctgcgcggca	1620
acggcaacag	ctacaacctg	tacctgcgcg	tcagctccat	tggcaactcc	accatcaggg	1680
tcaccatcaa	cgggaggggtg	tacacagcca	ccaatgtgaa	cacgacgacc	aacaatgatg	1740
gcgtcaacga	caacggcgcc	cgcttcagcg	acatcaacat	tggcaacgtg	gtggccagca	1800
gcaactccga	cgtcccgcgtg	gacatcaacg	tgaccttgaa	ctctggcacc	cagttcgacc	1860
tcattgaacat	catgctgggtg	ccaactaaca	tctcgccgct	gtactgatag	gagctctgat	1920
ccccatggga	attc					1934

<210> 2  
 <211> 634  
 <212> PRT  
 <213> *Bacillus thuringiensis*

<400> 2

```

Met Asp Asn Ser Val Leu Asn Ser Gly Arg Thr Thr Ile Cys Asp Ala
 1           5           10           15
Tyr Asn Val Ala Ala His Asp Pro Phe Ser Phe Gln His Lys Ser Leu
 20           25           30
Asp Thr Val Gln Lys Glu Trp Thr Glu Trp Lys Lys Asn Asn His Ser
 35           40           45
Leu Tyr Leu Asp Pro Ile Val Gly Thr Val Ala Ser Phe Leu Leu Lys
 50           55           60
Lys Val Gly Ser Leu Val Gly Lys Arg Ile Leu Ser Glu Leu Arg Asn
 65           70           75           80
Leu Ile Phe Pro Ser Gly Ser Thr Asn Leu Met Gln Asp Ile Leu Arg
 85           90           95
Glu Thr Glu Lys Phe Leu Asn Gln Arg Leu Asn Thr Asp Thr Leu Ala
100           105           110
Arg Val Asn Ala Glu Leu Thr Gly Leu Gln Ala Asn Val Glu Glu Phe
115           120           125
Asn Arg Gln Val Asp Asn Phe Leu Asn Pro Asn Arg Asn Ala Val Pro
130           135           140
Leu Ser Ile Thr Ser Ser Val Asn Thr Met Gln Gln Leu Phe Leu Asn
145           150           155           160
Arg Leu Pro Gln Phe Gln Met Gln Gly Tyr Gln Leu Leu Leu Leu Pro
165           170           175
Leu Phe Ala Gln Ala Ala Asn Leu His Leu Ser Phe Ile Arg Asp Val
180           185           190
Ile Leu Asn Ala Asp Glu Trp Gly Ile Ser Ala Ala Thr Leu Arg Thr
195           200           205
Tyr Arg Asp Tyr Leu Lys Asn Tyr Thr Arg Asp Tyr Ser Asn Tyr Cys
210           215           220
Ile Asn Thr Tyr Gln Ser Ala Phe Lys Gly Leu Asn Thr Arg Leu His
225           230           235           240
Asp Met Leu Glu Phe Arg Thr Tyr Met Phe Leu Asn Val Phe Glu Tyr
245           250           255
Val Ser Ile Trp Ser Leu Phe Lys Tyr Gln Ser Leu Leu Val Ser Ser
260           265           270
Gly Ala Asn Leu Tyr Ala Ser Gly Ser Gly Pro Gln Gln Thr Gln Ser
275           280           285
Phe Thr Ser Gln Asp Trp Pro Phe Leu Tyr Ser Leu Phe Gln Val Asn
290           295           300
Ser Asn Tyr Val Leu Asn Gly Phe Ser Gly Ala Arg Leu Ser Asn Thr
305           310           315           320
Phe Pro Asn Ile Val Gly Leu Pro Gly Ser Thr Thr Thr His Ala Leu
325           330           335
Leu Ala Ala Arg Val Asn Tyr Ser Gly Gly Ile Ser Ser Gly Asp Ile
340           345           350
Gly Ala Ser Pro Phe Asn Gln Asn Phe Asn Cys Ser Thr Phe Leu Pro
355           360           365
Pro Leu Leu Thr Pro Phe Val Arg Ser Trp Leu Asp Ser Gly Ser Asp
370           375           380
Arg Glu Gly Val Ala Thr Val Thr Asn Trp Gln Thr Glu Ser Phe Glu
385           390           395           400
Thr Thr Leu Gly Leu Arg Ser Gly Ala Phe Thr Ala Arg Gly Asn Ser
405           410           415
Asn Tyr Phe Pro Asp Tyr Phe Ile Arg Asn Ile Ser Gly Val Pro Leu
420           425           430

```

Val Val Arg Asn Glu Asp Leu Arg Arg Pro Leu His Tyr Asn Glu Ile  
 435 440 445  
 Arg Asn Ile Ala Ser Pro Ser Gly Thr Pro Gly Gly Ala Arg Ala Tyr  
 450 455 460  
 Met Val Ser Val His Asn Arg Lys Asn Asn Ile His Ala Val His Glu  
 465 470 475 480  
 Asn Gly Ser Met Ile His Leu Ala Pro Asn Asp Tyr Thr Gly Phe Thr  
 485 490 495  
 Ile Ser Pro Ile His Ala Thr Gln Val Asn Asn Gln Thr Arg Thr Phe  
 500 505 510  
 Ile Ser Glu Lys Phe Gly Asn Gln Gly Asp Ser Leu Arg Phe Glu Gln  
 515 520 525  
 Asn Asn Thr Thr Ala Arg Tyr Thr Leu Arg Gly Asn Gly Asn Ser Tyr  
 530 535 540  
 Asn Leu Tyr Leu Arg Val Ser Ser Ile Gly Asn Ser Thr Ile Arg Val  
 545 550 555 560  
 Thr Ile Asn Gly Arg Val Tyr Thr Ala Thr Asn Val Asn Thr Thr Thr  
 565 570 575  
 Asn Asn Asp Gly Val Asn Asp Asn Gly Ala Arg Phe Ser Asp Ile Asn  
 580 585 590  
 Ile Gly Asn Val Val Ala Ser Ser Asn Ser Asp Val Pro Leu Asp Ile  
 595 600 605  
 Asn Val Thr Leu Asn Ser Gly Thr Gln Phe Asp Leu Met Asn Ile Met  
 610 615 620  
 Leu Val Pro Thr Asn Ile Ser Pro Leu Tyr  
 625 630

<210> 3  
 <211> 415  
 <212> DNA  
 <213> Zea mays

<400> 3  
 tctagaggat cagcatggcg cccaccgtga tgatggcctc gtcggccacc gccgctgctc 60  
 cgttcctggg gctcaagtcc accgccagcc tccccgtcgc ccgccgtccc tccagaagcc 120  
 tcggcaacgt cagcaacggc ggaaggatcc ggtgcatgca ggtaacaaat gcatacctagc 180  
 tagtagttct ttgcattgca gcagctgcag ctacgcagtt agtaatatga agggaaactga 240  
 tgatccatgc atggactgat gtgtgttgcc catcccatcc catcccatcc cccaaacgaa 300  
 ccgaaaaacac cgtactacgt gcaggtgtgg ccctacggca acaagaagtt cgagacgctg 360  
 tcgtactctgc cgccgctgct gaccggcggg cgcacccgct gcatagcaggc catgg 415

<210> 4  
 <211> 79  
 <212> PRT  
 <213> Zea mays

<400> 4  
 Met Ala Pro Thr Val Met Met Ala Ser Ser Ala Thr Ala Val Ala Pro  
 1 5 10 15  
 Phe Leu Gly Leu Lys Ser Thr Ala Ser Leu Pro Val Ala Arg Arg Ser  
 20 25 30  
 Ser Arg Ser Leu Gly Asn Val Ser Asn Gly Gly Arg Ile Arg Cys Met  
 35 40 45  
 Gln Val Trp Pro Tyr Gly Asn Lys Lys Phe Glu Thr Leu Ser Tyr Leu  
 50 55 60  
 Pro Pro Leu Ser Thr Gly Gly Arg Ile Arg Cys Met Gln Ala Met  
 65 70 75

<210> 5  
 <211> 268  
 <212> DNA

<213> Artificial Sequence

<220>

<221> transit\_peptide

<222> 1-267

<223> coding sequence for PTP1 comprising an Arabidopsis thaliana ssRUBISCO (SSU) chloroplast targeting sequence and sequences coding for the first 24 amino acids of ssRUBISCO (SSU) protein (Wong et al., 1992)

<400> 5

```

atggcttcct ctatgctctc ttccgctact atggttgctc ctccggctca ggccactatg      60
gtcgtctcctt tcaacggact taagtctctc gctgccttcc cagccaccgc caaggctaac      120
aacgacatta cttccatcac aagcaacggc ggaagagtta actgcatgca ggtgtggcct      180
ccgattggaa agaagaagtt tgagactctc tcttaccttc ctgaccttac cgattccggt      240
ggtcgctgca actgcatgca ggccatgg      268
  
```

<210> 6

<211> 89

<212> PRT

<213> Arabidopsis thaliana

<400> 6

```

Met Ala Ser Ser Met Leu Ser Ser Ala Thr Met Val Ala Ser Pro Ala
 1           5           10           15
Gln Ala Thr Met Val Ala Pro Phe Asn Gly Leu Lys Ser Ser Ala Ala
      20           25           30
Phe Pro Ala Thr Arg Lys Ala Asn Asn Asp Ile Thr Ser Ile Thr Ser
      35           40           45
Asn Gly Gly Arg Val Asn Cys Met Gln Val Trp Pro Pro Ile Gly Lys
      50           55           60
Lys Lys Phe Glu Thr Leu Ser Tyr Leu Pro Asp Leu Thr Asp Ser Gly
      65           70           75           80
Gly Arg Val Asn Cys Met Gln Ala Met
              85
  
```

<210> 7

<211> 178

<212> DNA

<213> Arabidopsis thaliana

<400> 7

```

atggcttcct ctatgctctc ttccgctact atggttgctc ctccggctca ggccactatg      60
gtcgtctcctt tcaacggact taagtctctc gctgccttcc cagccaccgc caaggctaac      120
aacgacatta cttccatcac aagcaacggc ggaagagtta actgcatgca ggccatgg      178
  
```

<210> 8

<211> 59

<212> PRT

<213> Arabidopsis thaliana

<400> 8

```

Met Ala Ser Ser Met Leu Ser Ser Ala Thr Met Val Ala Ser Pro Ala
 1           5           10           15
Gln Ala Thr Met Val Ala Pro Phe Asn Gly Leu Lys Ser Ser Ala Ala
      20           25           30
Phe Pro Ala Thr Arg Lys Ala Asn Asn Asp Ile Thr Ser Ile Thr Ser
      35           40           45
Asn Gly Gly Arg Val Asn Cys Met Gln Ala Met
      50           55
  
```

<210> 9

<211> 240  
<212> DNA  
<213> Arabidopsis thaliana

<400> 9  
atggcgcaag ttagcagaat ctgcaatggt gtgcagaacc catctcttat ctccaatctc 60  
tcgaaatcca gtcaacgcaa atctccctta tcggtttctc tgaagacgca gcagcatcca 120  
cgagcttata cgatttcgtc gtcgtgggga ttgaagaaga gtgggatgac gtttaattggc 180  
tctgagcttc gtctctttaa ggtcatgtct tctgtttcca cggcgtgcat gcttgccatg 240

<210> 10  
<211> 80  
<212> PRT  
<213> Arabidopsis thaliana

<400> 10  
Met Ala Gln Val Ser Arg Ile Cys Asn Gly Val Gln Asn Pro Ser Leu  
1 5 10 15  
Ile Ser Asn Leu Ser Lys Ser Ser Gln Arg Lys Ser Pro Leu Ser Val  
20 25 30  
Ser Leu Lys Thr Gln Gln His Pro Arg Ala Tyr Pro Ile Ser Ser Ser  
35 40 45  
Trp Gly Leu Lys Lys Ser Gly Met Thr Leu Ile Gly Ser Glu Leu Arg  
50 55 60  
Pro Leu Lys Val Met Ser Ser Val Ser Thr Ala Cys Met Leu Ala Met  
65 70 75 80

<210> 11  
<211> 1907  
<212> DNA  
<213> Artificial Sequence

<220>  
<223> completely synthesized

<400> 11  
ccatggacaa caacgtcttg aactctggta gaacaacat ctgcgacgca tacaacgtcg 60  
tggtctacga tccattcagc ttggaacaca agagcctcga cactattcag aaggagtggg 120  
tggaatggaa acgtactgac cactctctct acgtcgcacc tgtggttgga acagtgtcca 180  
gcttccttct caagaaggtc ggctctctca tcggaaaaacg tatcttgtcc gaactctggg 240  
gtatcatctt tccatctggg tccactaatc tcatgcaaga catcttgagg gagaccgaac 300  
agtttctcaa ccagcgtctc aacactgata ccttggttag agtcaacgtc gaggttgatcg 360  
gtctccaagc aaacattcgt gagttcaacc agcaagtggg caacttcttg aatccaactc 420  
agaatcctgt gcctctttcc atcacttctt ccgtgaacac tatgcagcaa ctcttctca 480  
acagattgcc tcagtttcag attcaaggct accagttgct ccttcttcca ctctttgctc 540  
aggctgcaa catgcacttg tccctcatac gtgacgtgat cctcaacgct gacgaatggg 600  
gaatctctgc agccactctt aggacataca gagactactt gaggaactac actcgtgatt 660  
actccaacta ttgcatcaac acttatcaga ctgcctttcg tggactcaat actaggcttc 720  
acgacatgct tgagttcagg acctacatgt tccctaacgt gtttgagtac gtcagcattt 780  
ggagtctctt caagtaccag agcttgatgg tgcctcttgg agccaatctc tacgctctg 840  
gcagtggacc acagcaaact cagagcttca cagctcagaa ctggccattc ttgtatagct 900  
tggtccaagt caactccaac tacattctca gtggtatctc tgggaccaga ctctccataa 960  
cctttcccaa cattggtgga cttccaggct ccactacaac ccatagcctt aactctgcca 1020  
gagtgaacta cagtggagggt gtcagctctg gattgattgg tgcaactaac ttgaaccaca 1080  
acttcaattg ctccaccgtc ttgccacctc tgagcacacc gtttgtaggg tctcggttg 1140  
acagcgttac tgatcgcgaa ggagttgcta cctctacaaa ctggcaaac gagtccttc 1200  
aaaccactct tagccttcgg tgtggagctt tctctgcacg tgggaattca aactactttc 1260  
cagactactt cattaggaac atctctgggtg ttcctctcgt catcaggaat gaagacctca 1320  
cccgtccact tcattacaac cagattagga acatcgagtc tccatccggt actccaggag 1380  
gtgcaagagc ttacctcgtg tctgtccata acaggaagaa caacatctac gctgccaacg 1440

agaatggcac	catgattcac	cttgaccacag	aagattacac	tggtaccac	atctctccaa	1500
tccatgctac	ccaagtgaac	aatcagacac	gcaccttcac	ctccgaaaag	ttcggaaatc	1560
aaggtgactc	cttgagggtc	gagcaatcca	acactaccgc	taggtacact	ttgagaggca	1620
atggaaacag	ctacaacctt	tacttgagag	ttagctccat	tggttaactcc	accatccgtg	1680
ttaccatcaa	cggacgtgtt	tacacagtct	ctaattgtgaa	cactacaacg	aacaatgatg	1740
gcgttaacga	caacggagcc	agattcagcg	acatcaacat	tggtcaacatc	gtggcctctg	1800
acaacactaa	cgttactttg	gacatcaatg	tgacctctaa	ttctggaact	ccatttgatc	1860
tcatgaacat	catgtttgtg	ccaactaacc	tcctccatt	gtactaa		1907

&lt;210&gt; 12

&lt;211&gt; 634

&lt;212&gt; PRT

<213> *Bacillus thuringiensis*

&lt;400&gt; 12

Met	Asp	Asn	Asn	Val	Leu	Asn	Ser	Gly	Arg	Thr	Thr	Ile	Cys	Asp	Ala	1	5	10	15
Tyr	Asn	Val	Val	Ala	His	Asp	Pro	Phe	Ser	Phe	Glu	His	Lys	Ser	Leu	20	25	30	
Asp	Thr	Ile	Gln	Lys	Glu	Trp	Met	Glu	Trp	Lys	Arg	Thr	Asp	His	Ser	35	40	45	
Leu	Tyr	Val	Ala	Pro	Val	Val	Gly	Thr	Val	Ser	Ser	Phe	Leu	Leu	Lys	50	55	60	
Lys	Val	Gly	Ser	Leu	Ile	Gly	Lys	Arg	Ile	Leu	Ser	Glu	Leu	Trp	Gly	65	70	75	80
Ile	Ile	Phe	Pro	Ser	Gly	Ser	Thr	Asn	Leu	Met	Gln	Asp	Ile	Leu	Arg	85	90	95	
Glu	Thr	Glu	Gln	Phe	Leu	Asn	Gln	Arg	Leu	Asn	Thr	Asp	Thr	Leu	Ala	100	105	110	
Arg	Val	Asn	Ala	Glu	Leu	Ile	Gly	Leu	Gln	Ala	Asn	Ile	Arg	Glu	Phe	115	120	125	
Asn	Gln	Gln	Val	Asp	Asn	Phe	Leu	Asn	Pro	Thr	Gln	Asn	Pro	Val	Pro	130	135	140	
Leu	Ser	Ile	Thr	Ser	Ser	Val	Asn	Thr	Met	Gln	Gln	Leu	Phe	Leu	Asn	145	150	155	160
Arg	Leu	Pro	Gln	Phe	Gln	Ile	Gln	Gly	Tyr	Gln	Leu	Leu	Leu	Leu	Pro	165	170	175	
Leu	Phe	Ala	Gln	Ala	Ala	Asn	Met	His	Leu	Ser	Phe	Ile	Arg	Asp	Val	180	185	190	
Ile	Leu	Asn	Ala	Asp	Glu	Trp	Gly	Ile	Ser	Ala	Ala	Thr	Leu	Arg	Thr	195	200	205	
Tyr	Arg	Asp	Tyr	Leu	Arg	Asn	Tyr	Thr	Arg	Asp	Tyr	Ser	Asn	Tyr	Cys	210	215	220	
Ile	Asn	Thr	Tyr	Gln	Thr	Ala	Phe	Arg	Gly	Leu	Asn	Thr	Arg	Leu	His	225	230	235	240
Asp	Met	Leu	Glu	Phe	Arg	Thr	Tyr	Met	Phe	Leu	Asn	Val	Phe	Glu	Tyr	245	250	255	
Val	Ser	Ile	Trp	Ser	Leu	Phe	Lys	Tyr	Gln	Ser	Leu	Met	Val	Ser	Ser	260	265	270	
Gly	Ala	Asn	Leu	Tyr	Ala	Ser	Gly	Ser	Gly	Pro	Gln	Gln	Thr	Gln	Ser	275	280	285	
Phe	Thr	Ala	Gln	Asn	Trp	Pro	Phe	Leu	Tyr	Ser	Leu	Phe	Gln	Val	Asn	290	295	300	
Ser	Asn	Tyr	Ile	Leu	Ser	Gly	Ile	Ser	Gly	Thr	Arg	Leu	Ser	Ile	Thr	305	310	315	320
Phe	Pro	Asn	Ile	Gly	Gly	Leu	Pro	Gly	Ser	Thr	Thr	Thr	His	Ser	Leu	325	330	335	
Asn	Ser	Ala	Arg	Val	Asn	Tyr	Ser	Gly	Gly	Val	Ser	Ser	Gly	Leu	Ile	340	345	350	
Gly	Ala	Thr	Asn	Leu	Asn	His	Asn	Phe	Asn	Cys	Ser	Thr	Val	Leu	Pro	355	360	365	

Pro Leu Ser Thr Pro Phe Val Arg Ser Trp Leu Asp Ser Gly Thr Asp  
 370 375 380  
 Arg Glu Gly Val Ala Thr Ser Thr Asn Trp Gln Thr Glu Ser Phe Gln  
 385 390 395 400  
 Thr Thr Leu Ser Leu Arg Cys Gly Ala Phe Ser Ala Arg Gly Asn Ser  
 405 410 415  
 Asn Tyr Phe Pro Asp Tyr Phe Ile Arg Asn Ile Ser Gly Val Pro Leu  
 420 425 430  
 Val Ile Arg Asn Glu Asp Leu Thr Arg Pro Leu His Tyr Asn Gln Ile  
 435 440 445  
 Arg Asn Ile Glu Ser Pro Ser Gly Thr Pro Gly Gly Ala Arg Ala Tyr  
 450 455 460  
 Leu Val Ser Val His Asn Arg Lys Asn Asn Ile Tyr Ala Ala Asn Glu  
 465 470 475 480  
 Asn Gly Thr Met Ile His Leu Ala Pro Glu Asp Tyr Thr Gly Phe Thr  
 485 490 495  
 Ile Ser Pro Ile His Ala Thr Gln Val Asn Asn Gln Thr Arg Thr Phe  
 500 505 510  
 Ile Ser Glu Lys Phe Gly Asn Gln Gly Asp Ser Leu Arg Phe Glu Gln  
 515 520 525  
 Ser Asn Thr Thr Ala Arg Tyr Thr Leu Arg Gly Asn Gly Asn Ser Tyr  
 530 535 540  
 Asn Leu Tyr Leu Arg Val Ser Ser Ile Gly Asn Ser Thr Ile Arg Val  
 545 550 555 560  
 Thr Ile Asn Gly Arg Val Tyr Thr Val Ser Asn Val Asn Thr Thr Thr  
 565 570 575  
 Asn Asn Asp Gly Val Asn Asp Asn Gly Ala Arg Phe Ser Asp Ile Asn  
 580 585 590  
 Ile Gly Asn Ile Val Ala Ser Asp Asn Thr Asn Val Thr Leu Asp Ile  
 595 600 605  
 Asn Val Thr Leu Asn Ser Gly Thr Pro Phe Asp Leu Met Asn Ile Met  
 610 615 620  
 Phe Val Pro Thr Asn Leu Pro Pro Leu Tyr  
 625 630

<210> 13  
 <211> 10339  
 <212> DNA  
 <213> Artificial Sequence

<220>  
 <221> unsure  
 <222> 3687-3760; 4382-4434;  
 <223> "n" = g, a, c, or t

<400> 13  
 ggccgcgtta actgcaggtc cgatgtgaga cttttcaaca aagggttaata tccggaaacc 60  
 tctcggatt ccattgccca gctatctgtc actttattgt gaagatagtg gaaaaggaag 120  
 gtggctccta caaatgccat cattgcgata aaggaaaggc catcgttgaa gatgcctctg 180  
 ccgacagtgg tcccaaagat ggaccccccac ccacgaggag catcgtggaa aaagaagacg 240  
 ttccaaccac gtcttcaaag caagtggatt gatgtgatgg tccgatgtga gacttttcaa 300  
 caaagggtta tatccggaaa cctcctcgga ttccattgcc cagctatctg tcactttatt 360  
 gtgaagatag tggaaaagga aggtggctcc tacaaatgcc atcattgcga taaaggaaaag 420  
 gccatcgttg aagatgcctc tgccgacagt ggtcccaaaag atggaccccc acccacgagg 480  
 agcatcgttg aaaaagaaga cgttccaacc acgtcttcaa agcaagtgga ttgatgtgat 540  
 atctccactg acgtaaggga tgacgcacaa tcccactate cttcgcaaga cecttctct 600  
 atataaggaa gttcatttca tttggagagg acacagaaaa atttgctaca ttgtttcaca 660  
 aacttcaaatt attattcatt tttttgtcag ctttcaaact ctttgtttct tgtttgttga 720  
 ttgagaatac aatggcttcc tctatgctct cttccgctac tatggttgcc tctccggctc 780  
 aggccactat ggtcgtcctt ttcaacggac ttaagtcttc cgctgccttc ccagccaccc 840  
 gcaaggctaa caacgacatt acttccatca caagcaacgg cggaagagtt aactgcatgc 900

aggtgtggcc	tccgattgga	aagaagaagt	ttgagactct	ctettacctt	cetgacctta	960
ccgattccgg	tggtcgctgc	aactgcatgc	aggccatgga	caactccgtc	ctgaactctg	1020
gtcgcaccac	catctgcgac	gcctacaacg	tgcggcgca	tgatccattc	agcttccagc	1080
acaagagcct	cgacactgtt	cagaaggagt	ggacggagtg	gaagaagaac	aaccacagcc	1140
tgtacctgga	ccccatcgtc	ggcacgggtg	ccagcttcc	tctcaagaag	gtcggctctc	1200
tgcgcgggaa	gcgcatcctc	tccgaactcc	gcaacctgat	ctttccatct	ggctccacca	1260
acctcatgca	agacatcctc	agggagaccg	agaagtttct	caaccagcgc	ctcaacactg	1320
atacccttgc	tgcgctcaac	gctgagctga	cgggtctgca	agcaaacgtg	gaggagtcca	1380
accgccaagt	ggacaacttc	ctcaacccca	accgcaatgc	gggtgcctctg	tccatcactt	1440
cttccgtgaa	caccatgcaa	caactgttcc	tcaaccgctt	gcctcagttc	cagatgcaag	1500
gctaccagct	gctcctgctg	ccactctttg	ctcaggctgc	caacctgcac	ctctccttca	1560
ttcgtgacgt	gatcctcaac	gctgacgagt	ggggcatctc	tgcagccacg	ctgaggacct	1620
accgcgacta	cctgaagaac	tacaccaggg	actactcaa	ctattgcac	aacacctacc	1680
agtggcctt	caagggcctc	aatacagagg	ttcacgacat	gctggagtcc	aggacctaca	1740
tgttccgtga	cggtttcgag	tacgtcagca	tctggctcgt	cttcaagta	cagagcctgc	1800
tggtgtccag	cgggcgcaac	ctctacgcca	gcggctctgg	tccccacaa	actcagagct	1860
tcaccagcca	ggactggcca	ttcctgtatt	cggtgttcca	agtcaactcc	aactacgtcc	1920
tcaacggctt	ctctgggtgct	cgcctctcca	acaccttccc	caacattggt	ggcctccccg	1980
gctccaccac	aactcatgct	ctgcttgctg	ccagagtga	ctactccggc	ggcatctoga	2040
gcggcgacat	tggtgcacat	ccgttcaacc	agaacttcaa	ctgctccacc	ttcctgcccgc	2100
cgctgctcac	ccggttcgtg	aggtcctggc	tcgacagcgg	ctccgaccgc	gagggcggtg	2160
ccaccgtcac	caactggcaa	accgagtcc	tcgagaccac	ccttggtccc	cgagcggtcg	2220
ccttcacggc	gcgtgggaat	tctaactact	tccccgacta	cttcatcagg	aacatctctg	2280
gtgttctct	cgctgctccg	aacgaggacc	tccgctgccc	actgcactac	aacgagatca	2340
ggaacatcgc	ctctccgtcc	gggacgcccc	gaggtgcaag	ggcgtagcatg	gtgagcgctcc	2400
ataacaggaa	gaacaacatc	cacgctgtgc	atgagaaagg	ctccatgatc	cacctggcgc	2460
ccaatgatta	caccggcttc	accatctctc	caatccacgc	cacccaagtg	aacaaccaga	2520
cacgcacctt	catctccgag	aagttcggca	accagggcga	ctccctgagg	ttcgagcaga	2580
acaacaccac	cgccaggtag	accctgcggc	gcaacggcaa	cagctacaac	ctgtacctgc	2640
ggtcagctc	cattggcaac	tccaccatca	gggtcaccat	caaaggagg	gtgtacacag	2700
ccaccaatgt	gaacacgacg	accaacaatg	atggcgctcaa	cgacaacggc	gcccgttcca	2760
gcgacatcaa	cattggcaac	gtgggtggcca	gcagcaactc	cgacgtcccc	ctggacatca	2820
acgtgacctt	gaactctggc	accagttctg	acctcatgaa	catcatgctg	gtgccaaacta	2880
acatctcgcc	gctgtactga	taggagctct	gatccccatg	ggaattcccc	atcggttcaaa	2940
catttggcaa	taaagtctct	taagattgaa	tctgttggcc	ggctcttgca	tgattatcat	3000
ataatttctg	ttgaattacg	ttaagcatgt	aataactaac	atgtaatgca	tgacgttatt	3060
tatgagatgg	gtttttatga	ttagagtccc	gcaattatac	atttaatacg	cgatagaaaa	3120
caaaatatag	cgcgcaaaact	aggataaatt	atcgcgcgcg	gtgtcatcta	tgttactaga	3180
tccgggatata	ccccggggcg	gccgctcgag	tggtggccgc	atcgatcgtg	aagtttctca	3240
tctaagcccc	catttggacg	tgaatgtaga	cacgtcgaaa	taaagatttc	cgaattagaa	3300
taattttgtt	attgctttcg	cctataaata	cgacggatcg	taatttgcg	ttttatcaaa	3360
atgtactttc	attttataat	aacgctgcgg	acatctacat	ttttgaattg	aaaaaaaatt	3420
ggttaattact	ctttcttttt	ctccatattg	accatcatac	tcattgctga	tccatgtaga	3480
tttcccgga	atgaagccat	ttacaattga	atatatcctg	ccgcccgtgc	cgctttgcac	3540
ccggtggagc	ttgcatgttg	gtttctacgc	agaactgagc	cgggttaggca	gataatttcc	3600
attgagaact	gagccatgtg	caccttcccc	ccaacacggg	gagcgacggg	gcaacggagt	3660
gateccatg	ggacttttcc	tagctttnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	3720
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	3780
ggggcaccac	ccttcggcgt	gcgcgggtcac	gcgccagggc	gcagccctgg	ttaaaaacaa	3840
ggtttataaa	tattggttta	aaagcagggt	aaaagacagg	ttagcggtgg	ccgaaaaacg	3900
ggcggaaacc	cttgcaaatg	ctggattttc	tgctgtgga	cagccctca	aatgtcaata	3960
ggtgcgcccc	tcactctgtc	tcactctgcc	cctcaagtgt	caaggatcgc	gcccctcatc	4020
tgctcagtagt	cgcgccccctc	aagtgtcaat	accgcagggc	acttatcccc	aggctgttcc	4080
acatcatctg	tgggaaaactc	gcgtaaaaatc	aggcggtttc	gccgatttgc	gaggctggcc	4140
agctccacgt	cgcgcggcca	aatcgagcct	gcccctcatc	tgtcaacgcc	gcgcggggtg	4200
agtcggcccc	tcaagtgtca	acgtccgccc	ctcatctgtc	agtgaggggc	aagttttccg	4260
cgtggatatcc	acaacgcggg	cggccggccg	cggtgtctcg	cacacggctt	cgacggcggt	4320
tctggcgcg	ttgcaggggc	atagacggcc	gccagccag	cggcgagggc	aaccagcccg	4380
gnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4440
cgaccgatgc	ccttgagagc	cttcaaccca	gtcagctcct	tccggtgggc	gcggggcatg	4500
actatcgtcg	ccgcacttat	gactgtcttc	tttatcatgc	aactcgtagg	acaggtgccg	4560



gcagcgctct	gggtcatttt	cggcgaggac	cgctttcgct	ggagcgcgac	gatgatcggc	4620
ctgtcgcttg	cggtattcgg	aatcttgcac	gccctcgctc	aagccttcgt	cactgggtccc	4680
gccaccaaac	gtttcggcga	gaagcaggcc	attatcgccg	gcattggcggc	cgacgcgctg	4740
ggctacgtct	tgctggcggt	cgcgacgcga	ggctggatgg	ccttcccat	tatgattctt	4800
ctcgcttcgg	gcggcatcgg	gatgcccggg	ttgcaggcca	tgctgtccag	gcaggtagat	4860
gacgaccatc	agggacagct	tcaaggatcg	ctcgcggtc	ttaccagcct	aacttcgatc	4920
actggaccgc	tgatcgtcac	ggcgatttat	gccgcctcgg	cgagcacatg	gaacgggttg	4980
gcattggattg	taggcgcgcg	cctatacctt	gtctgcctcc	ccgcgttgcg	tcggcggtgca	5040
tgagagccggg	ccacctcgac	ctgaatggaa	gccggcgcca	cctcgctaac	ggattcacca	5100
ctccaagaat	tgagccaat	caattcttgc	ggagaactgt	gaatgcgcaa	accaaccctt	5160
ggcagaacat	atccatcgcg	tccgccatct	ccagcagccg	cacgcggcgc	atctcgggca	5220
gcgttgggtc	ctggccacgg	gtgcgcatga	tcgtgtcct	gtcgttgagg	acccggctag	5280
gctggcgggg	ttgccttact	ggttagcaga	atgaatcacc	gatacgcgag	cgaacgtgaa	5340
gcgactgctg	ctgcaaaaacg	tctgcgacct	gagcaacaac	atgaatggtc	ttcggtttcc	5400
gtgtttcgta	aagtctggaa	acgcggaagt	cagcgccctg	caccattatg	ttcgggattc	5460
ggatcgccagg	atgctgctgg	ctaccctgtg	gaacacctac	atctgtatta	acgaagcgct	5520
ggcattgacc	ctgagtgttt	tttctctggt	cccgccgcat	ccataccgcc	agttgtttac	5580
cctcacaacg	ttccagtaac	cgggcatggt	catcatcagt	aaccctgata	gtgagcatcc	5640
tctctcggtt	catcggtatc	attaccccca	tgaacagaaa	ttccccctta	cacggaggca	5700
tcaagtgacc	aaacaggaaa	aaaccgcctt	taacatggcc	cgctttatca	gaagccagac	5760
attaacgctt	ctggagaaac	tcaacgagct	ggagcgggat	gaacaggcag	acatctgtga	5820
atcgcttcac	gaccacgctg	atgagcttta	ccgcagctgc	ctcgcgcggt	tcgggtgatga	5880
cggtgaaaac	ctctgacaca	tgacgctccc	ggagacgggt	acagcttgct	tgtaagcgga	5940
tgccggggagc	agacaagccc	gtcagggcgc	gtcagcggtg	gttggcggtg	gtcggggcgc	6000
agccatgacc	cagtcacgta	gcgatagcgg	agtgtatact	ggcttaacta	tgccggcatca	6060
gagcagattg	tactgagagt	gcaccatatg	cgggtgtgaa	taccgcacag	atgctgaagg	6120
agaaaataacc	gcatacggcg	ctcttcggct	tcctcgctca	ctgactcgct	gcgctcggtc	6180
gttcggctgc	ggcgagcggt	atcagctcac	tcaaaaggcg	taatacgggt	atccacagaa	6240
tcagggggata	acgcaggaaa	gaacatgtga	gcaaaaggcc	agcaaaaggc	caggaaccgt	6300
aaaaaggccg	cggtgctggc	gtttttccat	aggtcccgcc	cccctgacga	gcatacaaaa	6360
aatcgacgct	caagtcagag	gtggcgaaac	ccgacaggac	tataaagata	ccaggcggtt	6420
ccccctggaa	gctccctcgt	gcgctctcct	gttcggaccc	tgccgcttac	cggtatacctg	6480
tcgcgctttc	tccttcgggg	aagcgtggcg	ctttctcata	gtcacgctg	taggtatctc	6540
agttcggtgt	aggtcggttcg	ctccaagctg	ggctgtgtgc	acgaaccccc	cggtcagccc	6600
gaccgctgcg	ccttatccgg	taactatcgt	cttgagctca	acccggtgag	acacgactta	6660
tcgccactgg	cagcagccac	tggttaacagg	attagcagag	cgaggtatgt	agccggtgct	6720
acagagtctt	tgaagtgggt	gcctaactac	ggctacacta	gaaggacagt	atttggtatc	6780
tgcgctctgc	tgaagccagt	taccttcgga	aaaagagtgt	gtagctcttg	atccggcaaa	6840
caaacaccgc	ctggttagcgg	tggttttttt	gtttgcaagc	agcagattac	gcgcagaaaa	6900
aaaggatctc	aagaagatcc	tttgatcttt	tctacggggt	ctgacgctca	gtggaacgaa	6960
aactcacggt	aagggaatttt	ggctcatgaga	ttatcaaaaa	ggatcttcac	ctagatcctt	7020
ttaaattaaa	aatgaagttt	taaatcaatc	taaagtatat	atgagtaaac	ttggtctgac	7080
agttaccaat	gcttaatcag	tgaggcacct	atctcagcga	tctgtctatt	tcgttcaccc	7140
atagtgcct	gactccccgt	cgtgtagata	actacgatac	gggagggctt	accatctggc	7200
cccagtgctg	caatgatacc	gcgagaccga	cgctcaccgg	ctccagattt	atcagcaata	7260
aaccagccag	ccggaagggc	cgagcgcaga	agtgggtcctg	caactttatc	cgctccatc	7320
cagctctatta	attgttgccg	ggaagctaga	gtaagtagtt	cgccagttaa	tagtttgccg	7380
aacgttggtg	ccattgctgc	aggtcgggag	cacaggatga	cgcttaacaa	ttcattcaag	7440
ccgacaccgc	ttcgcgccgc	ggcttaattc	aggagttaaa	catcatgagg	gaagcggtga	7500
tcgcccgaagt	atcgactcaa	ctatcagagg	tagttggcgt	catcgagcgc	catctcgaa	7560
cgacgttgct	ggccgtacat	ttgtacggct	ccgcagtgga	tgccggcctg	aagccacaca	7620
gtgatattga	tttgcgtggt	acgggtgaccg	taaggcttga	tgaaacaacg	cgccgagcct	7680
tgatcaacga	ccttttgga	acttcggctt	cccctggaga	gagcgagatt	ctccgcgctg	7740
tagaagtca	cattgttggt	cacgacgaca	tcattccgtg	gcgttatcca	gctaagcgcg	7800
aactgcaatt	tgagaaatgg	cagcgcaatg	acattcttgc	aggtatcttc	gagccagcca	7860
cgatcgacat	tgatctggct	atcttgctga	caaaagcaag	agaacatagc	gttgcttggt	7920
taggtccagc	ggcgagggaa	ctctttgatc	cggttccctga	acaggatcta	tttgaggcgc	7980
taaatgaaac	cttaacgcta	tggaactcgc	cgcccgactg	ggctggcgat	gagcgaaatg	8040
tagtgcttac	gttgctccgc	atcttggtaca	gcgcagtaac	cgccaaaatc	gcgcggaagg	8100
atgtcgctga	agactgggca	atggagcgcc	tgccggccca	gtatcagccc	gtcatacttg	8160
aagctaggca	ggcttatctt	ggacaagaag	atcgcttgcc	ctcgcgcgca	gatcagttgg	8220

aagaatttgt	tcaactacgtg	aaaggcgaga	tcaccaaggt	agtcggcaaa	taatgtctaa	8280
caattcgttc	aagccgacgc	cgcttcgcgg	cgcggttaa	ctcaagcgtt	agatgctgca	8340
ggcatcgtgg	tgtaacgctc	gtcgtttgg	atggcttcat	tcagctccgg	ttcccaacga	8400
tcaaggcgag	ttacatgata	ccccatgttg	tgcaaaaaag	cggtagctc	cttcggtcct	8460
ccgatcgagg	atctttcggc	gctgcgtac	gtccgckacc	gcgttgagg	atcaagccac	8520
agcagcccac	tcgacctcta	gccgaccag	acgagccaag	ggatctttt	ggaatgctgc	8580
tccgtcgtca	ggctttccga	cgtttggtg	ggtgaacaga	agtcattatc	gtacggaatg	8640
ccaagcactc	ccgaggggaa	ccctgtggtt	ggcatgcaca	tacaaatgga	cgaacggata	8700
aaccttttca	cgccctttta	aatatccgtt	attctaataa	acgctctttt	ctcttaggtt	8760
taccgcgcaa	tatatcctgt	caaacactga	tagtttaaac	tgaaggcggg	aaacgacaat	8820
ctgatcccca	tcaagcttgg	tcagtgga	gctagcttcc	cgatcctatc	tgtaacttca	8880
tcaaaaggac	agtagaaaag	gaaggtggca	ctacaaaatgc	catcattgca	ataaaggaaa	8940
ggctatcggt	caagatgcct	ctgccgacag	tggtcccaaa	gatggacccc	caccacagag	9000
gagcatcgtg	gaaaaagaag	acgttccaac	cacgtcttca	aagcaagtgg	attgatgtga	9060
tacttccact	gacgtaagg	atgacgcaca	atcccaactat	ccttcgcaag	acccttcctc	9120
tatataagga	agttcatttc	atttgagag	gacacgctga	aatcaccagt	ctctctctac	9180
aagatcgggg	atctctagct	agacgatcgt	ttcgcatgat	tgaacaagat	ggattgcacg	9240
caggttctcc	ggccgcttgg	gtggagaggc	tattcggtta	tgactgggca	caacagacaa	9300
tcggctgctc	tgatgccgcc	gtgttcgggc	tgtaagcgca	ggggcgcccg	gttctttttg	9360
tcaagaccga	cctgtccggt	gccctgaatg	aactgcagga	cgaggcagcg	cggtatcgt	9420
ggctggccac	gacgggctgt	ccttgccgag	ctgtgctcga	cggtgtcact	gaagcgggaa	9480
gggaactggc	gctattgggc	gaagtgcgg	ggcaggatct	cctgtcatct	caccttgctc	9540
ctgccgagaa	agtatccatc	atggctgatg	caatgcggcg	gctgcatacg	cttgatccgg	9600
ctacctgccc	attcgaccac	caagcgaaac	atcgcatcga	gcgagcacgt	actcggatgg	9660
aagccggtct	tgatgatcag	gatgatctgg	acgaagagca	tcaggggctc	gcgccagccg	9720
aactgttcgc	caggctcaag	gcgcgcatgc	ccgacggcga	ggatctcgtc	gtgacccatg	9780
gcgatgcctg	cttgccgaat	atcatgggtg	aaaatggccg	cttttctgga	ttcatcgact	9840
gtggccggct	gggtgtggcg	gaccgctatc	aggacatagc	gttggtacc	cgatgatattg	9900
ctgaagagct	tggcggcgaa	tgggtcgacc	gcttctcgt	gctttacggt	atcgccgctc	9960
ccgattcgca	gcgcacgcgc	ttctatcgcc	ttcttgacga	gttcttctga	gcgggactct	10020
ggggttcgat	ccccaattcc	cgatcggtta	aacatttggc	aataaagttt	cttaagattg	10080
aatcctgttg	ccggtcttgc	gatgattatc	atataatttc	tggtgaatta	cgtaaagcat	10140
gtaataatta	acatgtaatg	catgacgtta	tttatgagat	gggtttttat	gattagagtc	10200
ccgcaattat	acatttaata	cgcgatagaa	aacaaaatat	agcgcgcaaa	ctaggataaa	10260
ttatcgcgcg	cggtgtcatc	tatgttacta	gatcggggat	cgggccactc	gaccaagcct	10320
ctgcaggtcc	tgctcgagc					10339

&lt;210&gt; 14

&lt;211&gt; 10249

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 3597-3670; 4292-4344;

&lt;223&gt; "n" = g, a, c, or t

&lt;400&gt; 14

ggccgcgtta	actgcaggtc	cgatgtgaga	cttttcaaca	aagggttaata	tccggaaacc	60
tcctcggatt	ccattgccca	gctatctgtc	actttattgt	gaagatagtg	gaaaagggaag	120
gtggctccta	caaatgccat	cattgcgata	aaggaaaggc	catcgttgaa	gatgcctctg	180
ccgacagtgg	tcccaaagat	ggacccccac	ccacgaggag	catcgtggaa	aaagaagacg	240
ttccaaccac	gtcttcaaag	caagtggatt	gatgtgatgg	tccgatgtga	gacttttcaa	300
caaagggtaa	tatccggaaa	cctcctcgga	ttccattgcc	cagctatctg	tacttttatt	360
gtgaagatag	tggaaaagga	aggtggctcc	tacaaatgcc	atcattgcga	taaaaggaaag	420
gccatcgttg	aagatgcctc	tgccgacagt	ggccccaaag	atggaccccc	acccacgagg	480
agcatcgtgg	aaaaagaaga	cgttccaacc	acgtcttcaa	agcaagtggg	ttgatgtgat	540
atctccactg	acgtaaggga	tgacgcacaa	tcccactatc	cttcgcaaga	cccttcctct	600
atataaggaa	gttcatttca	tttgagagg	acacagaaaa	atttgctaca	ttgtttcaca	660

aacttcaaat	attattcatt	tatttgcag	ctttcaaaact	ctttgtttct	tgtttgttga	720
ttgagaatac	aatggcttcc	tctatgctct	cttcggtac	tatggttgcc	tctccggctc	780
aggccactat	ggtcgctcct	ttcaacggac	tttaagtctc	cgctgccttc	ccagccaccc	840
gcaaggctaa	caacgacatt	acttccatca	caagcaacgg	cggaagagtt	aactgcatgc	900
aggccatgga	caactccgtc	ctgaactctg	gtcgaccac	catctgcgac	gcctacaacg	960
tcgcggcgca	tgatccattc	agcttccagc	acaagagcct	cgacactgtt	cagaaggagt	1020
ggacggagtg	gaagaagaac	aaccacagcc	tgtacctgga	ccccatcgtc	ggcacgggtg	1080
ccagcttctc	tctcaagaag	gtcggctctc	tcgtcgggaa	gcgcacccctc	tcggaactcc	1140
gcaacctgat	ctttccatct	ggctccacca	acctcatgca	agacatccctc	aggagagacc	1200
agaagtcttct	caaccagcgc	ctcaacactg	atacccttgc	tcgctgcaac	gctgagctga	1260
cgggtctgca	agcaaacgtg	gaggagttca	accgccaaagt	ggacaacttc	ctcaacccca	1320
accgcaatgc	ggtgcctctg	tccatcactt	cttcctgtga	caccatgcaa	caactgttcc	1380
tcaaccgctt	gcctcagttc	cagatgcaag	gctaccagct	gctcctgtg	ccactctttg	1440
ctcagggtgc	caacctgcac	ctctccttca	ttcgtgacgt	gatcctcaac	gctgacgagt	1500
ggggcactctc	tcagccacg	ctgaggacct	accgcgacta	cctgaagaac	tacaccaggg	1560
actactccaa	ctattgcac	aacacctacc	agtcggcctt	caagggcctc	aatacagggc	1620
ttcacgacat	gctggagttc	aggacctaca	tgttctcgaa	cggtgtcgag	tacgtcagca	1680
tctggctcgt	cttcaagtag	cagagcctgc	tggtgtccag	cggcgccaac	ctctacgcca	1740
cgcggtctgg	tccccaacaa	actcagagct	tcaccagcca	ggactggcca	ttcctgtatt	1800
cgttgttcca	agtcaactcc	aactacgtcc	tcaacggctt	ctctggtgct	cgctctcca	1860
acaccttccc	caacattggt	ggcctccccg	gctccaccac	aactcatgct	ctgcttctg	1920
ccagagtga	ctactccggc	ggcatctcga	gcggcgacat	tggtgcatcg	ccgttcaacc	1980
agaacttcaa	ctgctccacc	ttcctgccgc	cgctgctcac	ccggttcgtg	aggtcctggc	2040
tcgacagcgg	ctccgaccgc	gagggcggtg	ccaccgtcac	caactggcaa	accgagtcct	2100
tcgagaccac	ccttggcctc	cggagcggcg	ccttcacggc	gcgtgggaat	tctaactact	2160
tccccgacta	cttcatcagg	aacatctctg	gtgttctctc	cgctcctccg	aacgaggacc	2220
tccgcggtcc	actgcactac	aacgagatca	ggaacatcgc	ctctccgtcc	gggacgccc	2280
gaggtgcaag	ggcgtacatg	gtgagcgtcc	ataacaggaa	gaacaacatc	cacgctgtgc	2340
atgagaacgg	ctccatgac	cacctggcgc	ccaatgatta	caccggcttc	accatctctc	2400
caatccacgc	caaccaagtg	aacaaccaga	cacgcacctt	catctccgag	aagttcggca	2460
accagggcga	ctccctgagg	ttcgagcaga	acaacaccac	cgccaggtag	accctgcgcg	2520
gcaacggcaa	cagctacaac	ctgtacctgc	gcgtcagctc	cattggcaac	tccaccatca	2580
gggtcaccat	caacgggagg	gtgtacacag	ccaaccaatg	gaacacgacg	accaacaatg	2640
atggcgctcaa	cgacaacggc	gcccgttcca	gcgacatcaa	cattggcaac	gtgggtggcca	2700
gcagcaacct	cgacgtcccg	ctggacatca	acgtgacctc	gaactctggc	accagttcgc	2760
acctcatgaa	catcatgctg	gtgccaaacta	acatctcgcc	gctgtactga	taggagctct	2820
gatcccatg	ggaattccc	atcggttcaa	catttggaac	ttaaagttct	taagattgaa	2880
tcctgttgcc	ggtcttgcca	tgattatcat	ataatttctg	ttgaattacg	ttaagcatgt	2940
aataattaac	atgtaatgca	tgacgttatt	tatgagatgg	gtttttatga	ttagagtccc	3000
gcaattatac	atttaatacg	cgatagaataa	caaaatatag	cgcgcaaaact	aggataaatt	3060
atcgcgcgcg	gtgtcatcta	tgttactaga	tcggggatat	ccccggggcg	gccgctcgag	3120
tggtggccgc	atcgatcggtg	aagtttctca	tctaagcccc	catttggaacg	tgaattgtaga	3180
cacgtcgaaa	taaagatttc	cgaaattgaa	taatttgttt	attgctttcg	cctataaata	3240
cgacggatcg	taatttgtcg	ttttatcaaa	atgtactttc	attttataat	aacgctgcgg	3300
acatctacat	ttttgaattg	aaaaaaaatt	ggtaattact	ctttcttttt	ctccatattg	3360
accatcatac	tcattgctga	tccatgtaga	tttcccggac	atgaagccat	ttacaattga	3420
atatacctg	ccgcccgtgc	cgctttgcac	ccggtggagc	ttgcatgttg	gtttctacgc	3480
agaactgagc	cgttagggca	gataatttcc	attgagaact	gagccatgtg	caccttcccc	3540
ccaacacggt	gagcgacggg	gcaacggagt	gatccacatg	ggacttttcc	tagctttnnnn	3600
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	3660
nnnnnnnnnn	ccgggagggg	tcgagaaggg	ggggcaccctc	ccttcggcgt	gcgcgggtcac	3720
gcgccagggc	gcagccctgg	ttaaaaacaa	ggtttataaa	tattggttta	aaagcaggtt	3780
aaaagacagg	ttagcgggtg	ccgaaaaacg	ggcggaaccc	cttgcaaatg	ctggattttc	3840
tgccctgtgga	cagccctcca	aatgtcaata	gggtgcgccc	tcatctgtca	tcactctgcc	3900
cctcaagtgt	caaggatcgc	gcccctcatc	tgctcagtagt	cgcgcccttc	aagtgtcaat	3960
accgcagggc	acttatcccc	aggcttgtcc	acatcatctg	tgggaaactc	gcgtaaaaatc	4020
aggcgttttc	gccgattttg	gaggctggcc	agctccacgt	cgccggccga	aatcgagcct	4080
gcccctcatc	tgtcaacgcc	gcgccgggtg	agtcggcccc	tcaagtgtca	acgtccgccc	4140
ctcatctgtc	agtgaaggcc	aagttttccg	cgtgggtatcc	acaacgcggg	cggccggccg	4200
cgggtgtctg	cacacggctt	cgacggcggt	tctggcgcg	ttgcagggcc	atagacggcc	4260
gccagcccg	cggcgagggc	aaccagcccg	gnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4320

nnnnnnnnnn	nnnnnnnnnn	nnnnngtcgat	cgaccgatgc	ccttgagagc	cttcaaccca	4380
gtcagctcct	tccggtgggc	gcggggcatg	actatcgctc	ccgcacttat	gactgtcttc	4440
tttatctagc	aactcgtagg	acaggtgccc	gcagcgctct	gggtcatttt	cggcgaggac	4500
cgctttcgct	ggagcgcgac	gatgatcggc	ctgtcgcttg	cggtattcgg	aatcttgac	4560
gccctcgctc	aagccttcgt	cactgggtccc	gccaccaaac	gtttcggcga	gaagcaggcc	4620
attatcgccc	gcatggcggc	cgacgcgctg	ggctacgtct	tgctggcggt	cgcgacgcga	4680
ggctggatgg	ccttccccat	tatgattcct	ctcgcttcgg	gcggcatcgg	gatgcccgcg	4740
ttgcaggcca	tgctgtccag	gcaggtagat	gacgaccatc	agggacagct	tcaaggatcg	4800
ctcgcgctc	ttaccagcct	aacttcgatc	actggaccgc	tgatcgtcac	ggcgatttat	4860
gcgcctcgg	cgagcacatg	gaacgggttg	gcatggattg	taggcgcgcg	cctatacctt	4920
gtctgcctcc	ccgcgttgcg	tcgcgggtgca	tggagccggg	ccacctcgac	ctgaatggaa	4980
gccggcgcca	cctcgctaac	ggattcacca	ctccaagaat	tggagccaat	caattcttgc	5040
ggagaactgt	gaatgcgcaa	accaaccctt	ggcagaacat	atccatcgcg	tccgccatct	5100
ccagcagccg	cacgcggcgc	atctcgggca	gcgttgggtc	ctggccacgg	gtgcgcatga	5160
tcgtgtcctc	gtcgttgagg	acccggctag	gctggcgggg	ttgccttact	ggtagcaga	5220
atgaatcacc	gatacgcgag	cgaacgtgaa	gcgactgctg	ctgcaaaacg	tctgcgacct	5280
gagcaacaac	atgaatggtc	ttcggtttcc	gtgtttcgta	aagtctggaa	acgcggaagt	5340
cagcgccctg	caccattatg	ttccggatct	gcatcgcagg	atgctgctgg	ctacctgtg	5400
gaacacctac	atctgtatta	acgaagcgct	ggcattgacc	ctgagtgttt	tttctctggt	5460
ccgcgcgcat	ccataccgcc	agttgtttac	cctcacaaac	ttccagtaac	cgggcatggt	5520
catcatcagt	aaccgcgtatc	gtgagcatcc	tctctcggtt	catcggtatc	attaccacca	5580
tgaacagaaa	ttccccctta	cacggaggca	tcaagtgtac	aaacaggaaa	aaaccgcctt	5640
taacatggcc	cgcttttatca	gaagccagac	attaacgctt	ctggagaaac	tcaacgagct	5700
ggagcgggat	gaacaggcag	acatctgtga	atcgcttcac	gaccacgctg	atgagcttta	5760
ccgcagctgc	ctgcgcgctt	tcgggtgatga	cggtgaaaac	ctctgacaca	tgagctccc	5820
ggagacggtc	acagcttgct	tgtaaagcga	tgccgggagc	agacaagccc	gtcagggcgc	5880
gtcagcgggt	gttggcgggt	gtcggggcgc	agccatgtac	cagtacgta	gcgatagcgg	5940
agtgtatact	ggcttaacta	tgccggcatca	gagcagattg	tactgagagt	gcacatatg	6000
cggtgtgaaa	taccgcacag	atgcgttaag	agaaaatacc	gcatcaggcg	ctcttccgct	6060
tcctcgctca	ctgactcgct	gcgctcggtc	gttcggctgc	ggcgagcggg	atcagctcac	6120
tcaaaggcgg	taatacgggt	atccacagaa	tcaggggata	acgcaggaaa	gaacatgtga	6180
gcaaaaggcc	agcaaaaggc	caggaaccgt	aaaaaggccg	cggtgctggc	gtttttccat	6240
aggctccgcc	cccctgacga	gcatcacaaa	aatcgacgct	caagtcagag	gtggcgaaac	6300
ccgacaggac	tataaagata	ccaggcggtt	ccccctggaa	gctccctcgt	gcgctctcct	6360
gttccgacct	tgccgcctac	cggatacctg	tcgcgcttcc	tccttccggg	aagcgtggcg	6420
ctttctcata	gtcacgctg	taggtatctc	agttcggtgt	aggtcggtcg	ctccaagctg	6480
ggctgtgtgc	acgaaccccc	cgttcagccc	gaccgctgcg	ccttatccgg	taactatcgt	6540
cttgagtcca	acccggttaag	acacgactta	tcgccactgg	cagcagccac	tggttaacagg	6600
attagcagag	cgaggtatgt	aggcggtgct	acagagttct	tgaagtggtg	gcctaactac	6660
ggctacacta	gaaggacagt	atttggtatc	tgcgctctgc	tgaagccagt	taccttcgga	6720
aaaagagttg	gtagctcttg	atccggcaaa	caaaccaccg	ctggtagcgg	tggttttttt	6780
gtttgcaagc	agcagattac	gcgcagaaaa	aaaggatctc	aagaagatcc	tttgatcttt	6840
tctacggggt	ctgacgctca	gtggaacgaa	aactcacggt	aagggatttt	ggatcatgaga	6900
ttatcaaaaa	ggatcttcac	ctagatcctt	ttaaattaaa	aatgaagttt	taaatcaatc	6960
taaagtatat	atgagtaaac	ttggtctgac	agttaccaat	gcttaatcag	tgaggcacct	7020
atctcagcga	tctgtctatt	tcgttcaccc	atagttgcct	gactccccgt	cgtgtagata	7080
actacgatac	gggagggtct	accatctggc	ccagtgctg	caatgatacc	gcgagaccca	7140
cgctcaccgg	ctccagattt	atcagcaata	aaccagccag	ccggaagggc	cgagcgcaga	7200
agtggtcctg	caactttatc	cgctccatc	cagtctatta	attgttgccg	ggaagctaga	7260
gtaagttagt	cgccagttaa	tagtttgctc	aacgttggtg	ccattgctgc	aggtcgggag	7320
cacaggatga	cgcctaacaa	ttcattcaag	ccgacaccgc	ttcgcgcgcg	ggcttaattc	7380
aggagttaaa	catcatgagg	gaagcgggtga	tcgcccgaagt	atcgactcaa	ctatcagagg	7440
tagttggcgt	catcgagcgc	catctcgaa	cgacgttgct	ggcggtacat	ttgtacggct	7500
ccgcagtggg	tgccggcctg	aagccacaca	gtgatattga	tttgctggtt	acggtgaccg	7560
taaggcttga	tgaacaacg	cggcgagctt	tgatcaacga	ccttttgga	acttcggctt	7620
ccccggaga	gagcgagatt	ctccgcgctg	tagaagtcac	cattgttgtg	cacgacgaca	7680
tcattccgtg	gcgttatoca	gctaagcgcg	aactgcattt	tggagaatgg	cagcgcaatg	7740
acattcttgc	aggtatcttc	gagccagcca	cgatcgacat	tgatctggct	atcttgctga	7800
caaaagcaag	agaacatagc	gttgcccttg	taggtccagc	ggcgaggagg	ctctttgatc	7860
cggttcctga	acaggatcta	tttgaggcgc	taaatgaaac	cttaacgcta	tggaaactcg	7920
cgcccgactg	ggctggcgat	gagcgaaatg	tagtgcttac	gttgtcccgc	atttggtaca	7980

gcgagtaac	cggcaaaatc	gcgccgaagg	atgtcgctga	agactgggca	atggagcgcc	8040
tgcgggcccc	gtatcagccc	gtcatacttg	aagctaggca	ggcttatctt	ggacaagaag	8100
atcgcttggc	ctcgcgcgca	gatcagttgg	aagaatttgt	tcactacgtg	aaaggcgaga	8160
tcaccaagggt	agtcggcaaa	taatgtctaa	caattcgttc	aagccgacgc	cgcttcgcgg	8220
cgcggcttaa	ctcaagcggt	agatgctgca	ggcatcgctg	tgtaacgctc	gtcgtttggt	8280
atggcttcat	tcagctccgg	ttcccaacga	tcaaggcgag	ttacatgatc	ccccatgttg	8340
tgcaaaaaag	cggtttagctc	cttcgggtcct	ccgatcgagg	atttttcggc	gctgcgctac	8400
gtccgckacc	gcgttgaggg	atcaagccac	agcagcccac	tcgacctcta	gccgaccag	8460
acgagccaag	ggatcttttt	ggaatgctgc	tccgtcgctca	ggctttccga	cgtttggttg	8520
gttgaacaga	agtcattatc	gtacggaatg	ccaagcactc	ccgaggggaa	ccctgtggtt	8580
ggcatgcaca	tacaaatgga	cgaacggata	aaccttttca	cgccctttta	aatatccgtt	8640
attctaataa	acgctctttt	ctcttagggt	tacccgcca	tatatcctgt	caaacactga	8700
tagtttaaac	tgaaggcggg	aaacgacaat	ctgatcccca	tcaagcttgg	tcgagtggaa	8760
gctagcttcc	cgatcctatc	tgtaacttca	tcaaaaggac	agtagaaaag	gaagtgggca	8820
ctacaaatgc	catcattgcg	ataaaggaaa	ggctatcggt	caagatgcct	ctgccgacag	8880
tggtcccaaa	gatggacccc	cacccacgag	gagcatcggt	gaaaaagaag	acgttccaac	8940
cacgtcttca	aagcaagtgg	attgatgtga	tacttccact	gacgtaaggg	atgacgcaca	9000
atcccaactat	ccttcgcaag	acccttctct	tatataagga	agttcatttc	atttgagag	9060
gacacgctga	aatcaccagt	ctctctctac	aagatcgggg	atctctagct	agacgatcgt	9120
ttcgcatgat	tgaacaagat	ggattgcacg	caggttctcc	ggccgcttgg	gtggagaggc	9180
tattcggtta	tgactgggca	caacagacaa	tcggctgctc	tgatgcgcgc	gtgttccggc	9240
tgtaacgca	ggggcgcccg	gttctttttg	tcaagaccga	cctgtccggt	gccctgaatg	9300
aactgcagga	cgaggcagcg	cggctatcgt	ggctggccac	gacgggctt	ccttgccgag	9360
ctgtgctcga	cggttgcact	gaagcgggaa	gggactgggt	gctattgggc	gaagtgcgg	9420
ggcaggatct	cctgtcatct	caccttgctc	ctgccgagaa	agtatccatc	atggctgatg	9480
caatgcggcg	gctgcatacg	cttgatccgg	ctacctgccc	attcgaccac	caagcgaaac	9540
atcgcatcga	gcgagcacgt	actcggatgg	aagccggtct	tgtaacgag	gatgatctgg	9600
acgaagagca	tcaggggctc	gcgccagccg	aactgttcgc	caggctcaag	gcgcgcagtc	9660
ccgacggcga	ggatctcgtc	gtgacccatg	gcgatgcctg	cttgccgaat	atcatgggtg	9720
aaaatggccg	cttttctgga	ttcatcgact	gtggccgggt	gggtgtggcg	gaccgctatc	9780
aggacatagc	gttggttacc	cgtgatattg	ctgaagagct	tgccggcgaa	tgggctgacc	9840
gcttctcgt	gctttacggt	atcgccgctc	ccgattcgca	gcgcacgcgc	ttctatcgcc	9900
ttcttgacga	gttcttctga	gcgggactct	gggggttcgat	cccccaattcc	cgatcggtta	9960
aacatttggc	aataaagttt	cttaagattg	aatcctgttg	ccggtcttgc	gatgattatc	10020
atataatttc	tggtgaatta	cgtaaagcat	gtaataatta	acatgtaatg	catgacgtta	10080
tttatgagat	gggtttttat	gattagagtc	cgcgaattat	acatttaata	cgcgatagaa	10140
aacaaaatat	agcgcgcaaa	ctaggataaa	ttatcgcgcg	cggtgtcatc	tatgttacta	10200
gatcggggat	cggggccactc	gaccaagctt	ctgcaggctc	tgctcgagc		10249

&lt;210&gt; 15

&lt;211&gt; 10312

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; unsure

&lt;222&gt; 3660-3773; 4355-4407;

&lt;223&gt; "n" = g, a, c, or t

&lt;400&gt; 15

ggccgcgtta	actgcaggtc	cgatgtgaga	cttttcaaca	aagggttaata	tccggaaacc	60
tcctcggatt	ccattgcccc	gctatctgtc	actttattgt	gaagatagtg	gaaaagggaag	120
gtggctccta	caaattgccat	cattgcgata	aaggaaaggc	catcggtgaa	gatgcctctg	180
ccgacagtgg	tcccaaagat	ggacccccc	ccacgaggag	catcggtgaa	aaagaagacg	240
ttccaaccac	gtcttcaaag	caagtggatt	gatgtgatgg	tccgatgtga	gacttttcaa	300
caaagggtta	tatccggaaa	cctcctcgga	ttccattgcc	cagctatctg	tcactttatt	360
gtgaagatag	tggaaaagga	aggtggctcc	tacaaatgcc	atcattgcga	taaaggaaaag	420
gccatcggtg	aagatgcctc	tgccgcagat	gggtcccaag	atggaccccc	acccacgagg	480
agcatcggtg	aaaaagaaga	cgttccaacc	acgtcttcaa	agcaagtgga	ttgatgtgat	540

atctccactg	acgtaaggga	tgacgcacaa	tcccactatc	cttcgcaaga	cccttcctct	600
atataaggaa	gttcatttca	tttggagagg	acacagaaaa	atttgctaca	ttgtttcaca	660
aacttcaaat	attattcatt	tatttgtcag	ctttcaaaact	ctttgtttct	tgtttgttga	720
ttgagaatac	aatggcgcaa	gttagcagaa	tctgcaatgg	tgtgcagaac	ccatctctta	780
tctccaatct	ctcgaaatcc	agtcaacgca	aatctccctt	atcgggtttct	ctgaagacgc	840
agcagcatcc	acgagcttat	ccgatttcgt	cgctgtgggg	attgaagaag	agtgggatga	900
cgtaatttgg	ctctgagctt	cgctctctta	aggtcatgtc	ttctgtttcc	acggcggtgca	960
tgcttgccat	ggacaactcc	gtcttgaact	ctggctgcac	caccatctgc	gacgcctaca	1020
acgtcgcggc	gcatgatcca	ttcagcttcc	agcacaagag	cctcgacact	gttcagaagg	1080
agtggacgga	gtggaaagaag	aacaaccaca	gcctgtacct	ggaccccatc	gtcggcacgg	1140
tggccagctt	ccttctcaag	aaggctcggt	ctctcgctcg	gaagcgcatc	ctctcggaac	1200
tccgcaacct	gatctttcca	tctggctcca	ccaacctcat	gcaagacatc	ctcagggaga	1260
ccgagaagtt	tctcaaccag	cgcttcaaca	ctgataccct	tgctcgctgc	aacgctgagc	1320
tgacgggtct	gcaagcaaac	gtggaggagt	tcaaccgcca	agtggacaac	ttcttcaacc	1380
ccaaccgcaa	tgcggtgcct	ctgtccatca	cttcttccgt	gaacaccatg	caacaactgt	1440
tcctcaaccg	cttgctcag	ttccagatgc	aaggtctacca	gctgtcctg	ctgccactct	1500
ttgctcaggc	tgccaacctg	cacctctcct	tcattcgtga	cgtgacctc	aacgctgacg	1560
agtggggcat	ctctgcagcc	acgctgagga	cctaccgcga	ctacctgaag	aactacacca	1620
gggactactc	caactattgc	atcaacacct	accagtcggc	cttcaagggc	ctcaatacga	1680
ggcttcacga	catgctggag	ttcaggacct	acatgttctt	gaacgtgttc	gagtacgtca	1740
gcatctgggt	gctcttcaag	taccagagcc	tgttggtgtc	cagcgccgcc	aacctctacg	1800
ccagcggtct	tggtccccaa	caaaactcaga	gtctcaccag	ccaggactgg	ccattcctgt	1860
attcggtgtt	ccaagtcaac	tccaactacg	tcctcaacgg	cttctctggt	gctcgcctct	1920
ccaacacctt	ccccaacatt	gttggectcc	ccggctccac	cacaactcat	gctctgcttg	1980
ctgccagagt	gaactactcc	ggcggtcatct	cgagcgccga	cattgggtgca	tcgcggttca	2040
accagaactt	caactgctcc	accttctctg	cgcgctgtct	caccccgctc	gtgaggtcct	2100
ggctcgacag	cggctccgac	cgcgagggcg	tggccaccgt	caccaactgg	caaaccgagt	2160
ccttcgagac	cacccttggt	ctccggagcg	gcgccttcac	ggcgcggtgg	aattctaaact	2220
acttccccga	ctacttcatc	aggaacatct	ctggtgttcc	tctcgtcgtc	cgcaacgagg	2280
acctcgcgcg	tccactgcac	tacaacgaga	tcaggaacat	cgcctctccg	tccgggacgc	2340
ccggaggtgc	aagggcggtac	atggtgagcg	tccataacag	gaagaacaac	atccacgctg	2400
tgcatgagaa	cggctccatg	atccacctgg	cgcccaatga	ttacaccggc	ttcaccatct	2460
ctccaatcca	cgccacccaa	gtgaacaacc	agacacgcac	cttcatctcc	gagaagtctg	2520
gcaaccaggg	cgactccctg	agggttcgagc	agaacaacac	caccgccagg	tacaccctgc	2580
gcggcaacgg	caacagctac	aacctgtacc	tgcgcgtcag	ctccattggc	aactccacca	2640
tcagggtcac	catcaacggg	aggggtgaca	cagccaccaa	tgtgaacacg	acgaccaaca	2700
atgatggcgt	caacgacaac	ggcgcccgct	tcagcgacat	caacattggc	aacgtggtgg	2760
ccagcagcaa	ctccgacgtc	ccgctggaca	tcaacgtgac	cctgaactct	ggcaccaggt	2820
tcgacctcat	gaacatcatg	ctggtgccaa	ctaacatctc	gccgctgtac	tgataggagc	2880
tctgatcccc	atgggaattc	ccgatcgttc	aaacatttgg	caataaagtt	tcttaagatt	2940
gaatcctggt	gccggtcttg	cgatgattat	catataattt	ctggtgaatt	acgttaaagca	3000
tgtaataatt	aacatgtaat	gcatgacgtt	attttagaga	tgggttttta	tgattagagt	3060
cccgaatta	tacatttaat	acgcgataga	aaacaaaata	tagcgcgcaa	actaggataa	3120
attatcgctc	gcggtgtcat	ctatgttact	agatcgggga	tatccccggg	gcggccgctc	3180
gagtggtggc	cgcactgatc	gtgaagtttc	tcatctaagc	ccccatttgg	acgtgaatgt	3240
agacacgtcg	aaataaagat	ttccgaatta	gaataatttg	tttattgctt	tcgcctataa	3300
atacgacgga	tcgtaatttg	tcgtttttatc	aaaatgtact	ttcattttat	aataacgctg	3360
cggacatcta	cattttttgaa	ttgaaaaaaa	attggttaatt	actctttctt	tttctccata	3420
ttgaccatca	tactcattgc	tgatccatgt	agatttcccg	gacatgaagc	catttacaat	3480
tgaatatatc	ctgccgccgc	tgccgctttg	cacccgggtg	agcttgcatg	ttggtttcta	3540
cgcgaactg	agccgggttag	gcagataatt	tccattgaga	actgagccat	gtgcaccttc	3600
cccccaacac	ggtagcgac	ggggcaacgg	agtatccac	atgggacttt	tcttagcttn	3660
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	3720
nnnnnnnnnn	nnnccgggag	ggttcgagaa	gggggggcac	cccecttcgg	cgtgcgcggt	3780
cacgcgccag	ggcgagccc	tggttaaaaa	caaggtttat	aaatattggt	ttaaaagcag	3840
gttaaaaagac	aggttagcgg	tggccgaaaa	acggcgggaa	acccttgcaa	atgctggatt	3900
ttctgcctgt	ggacagccc	tcaaatgtca	ataggtgcgc	ccctcatctg	tcatcactct	3960
gcccctcaag	tgtcaaggat	cgcgccctc	atctgtcagt	agtcgcgcc	ctcaagtgtc	4020
aataccgcag	ggcacttate	cccaggcttg	tccacatcat	ctgtgggaaa	ctcgcgtaaa	4080
atcaggcggt	ttcgccgatt	tgcgaggtcg	gccagctcca	cgtcgccggc	cgaaatcgag	4140
cctgcccctc	atctgtcaac	gccgcgccgg	gtgagtcggc	cctcaagtg	tcaacgtccg	4200

cccccatctt	gtcagtgagg	gccaagtttt	ccgcgtggta	tccacaacgc	cggcggccgg	4260
ccgcggtgtc	tcgcacacgg	cttcgacggc	gtttctggcg	cgtttgagg	gccatagacg	4320
gccgccagcc	cagcggcgag	ggcaaccagc	ccggnnnnnn	nnnnnnnnnn	nnnnnnnnnn	4380
nnnnnnnnnn	nnnnnnnnnn	nnnnnnngtc	gatcgaccga	tgcccttgag	agccttcaac	4440
ccagtcagct	ccttcgggtg	ggcgcggggc	atgactatcg	tcgcgcgact	tatgactgtc	4500
ttctttatca	tgcaactcgt	aggacaggtg	ccggcagcgc	tctgggtcat	tttcggcgag	4560
gaccgctttc	gctggagcgc	gacgatgac	ggcctgtcgc	ttgcgggtatt	cggaatcctg	4620
cacgccctcg	ctcaagcctt	cgctactggg	ccgcgccacca	aacgtttcgg	cgagaagcag	4680
gccattatcg	ccggcatggc	ggccgacgcg	ctgggctacg	tcttgctggc	gttcgcgacg	4740
cgaggctgga	tggccttccc	cattatgatt	cttctcgcct	ccggcgggcat	cgggatgcc	4800
gcgttgagg	ccatgctgtc	caggcaggta	gatgacgacc	atcagggaca	gcttcaagg	4860
tcgctcgcg	ctcttaccag	cctaacttcg	atcactggac	cgctgatcgt	cacggcgatt	4920
tatgcccgcct	cggcgagcac	atggaaagg	ttggcatgga	ttgtaggcgc	cgccctatac	4980
cttgctgccc	tccccgcgtt	gcgtcgcggt	gcatggagcc	gggccacctc	gacctgaatg	5040
gaagccggcg	gcacctcgt	aacggattca	ccactccaag	aattggagcc	aatcaattct	5100
tgcgagagac	tgtgaatgcg	caaaccaacc	cttgccagaa	catatccatc	gcgtccgcca	5160
tctccagcag	ccgcaacggg	cgcatctcgg	gcagcgttgg	gtcctggcca	cggttgccga	5220
tgatcgtgct	cctgtcgttg	aggaccgggc	taggctggcg	gggttgccct	actggttagc	5280
agaatgaatc	accgatacgc	gagcgaacgt	gaagcgactg	ctgctgcaaa	acgtctcgca	5340
cctgagcaac	aacatgaatg	gtcttcgggt	tccgtgtttc	gtaaagtctg	gaaacgcgga	5400
agtcagcgcc	ctgcaccatt	atgttcggga	tctgcacgcg	aggatgctgc	tggtaccct	5460
gtggaacacc	tacatctgta	ttaacgaagc	gctggcattg	accctgagtg	atctttctct	5520
ggtcccgccg	catccatacc	gccagttggt	taccttcaca	acgttccagt	aaccgggcat	5580
gttcacatc	agtaaccggt	atcgtgagca	tctctctcgc	ttctcatcgt	atcattacc	5640
ccatgaacag	aaattcccc	ttacacggag	gcatacaagt	accaaaccag	aaaaaacccg	5700
ccttaacatg	gccccgttta	tcagaagcca	gacattaacg	cttctggaga	aactcaacga	5760
gctggacggc	gatgaacagg	cagacatctg	tgaatcgctt	cacgaccacg	ctgatgagct	5820
ttaccgcagc	tgccctcgcg	gtttcggtga	tgacggtgaa	aacctctgac	acatgcagct	5880
cccgagagcg	gtcacagctt	gtctgtaagc	ggatgcgggg	agcagacaag	cccgtagagg	5940
cgcgtagcgg	gggtgtggcg	gggtgcgggg	cgacgccatg	acccagtcac	gtagcgatag	6000
cggagtgtat	actggcttaa	ctatgcggca	tcagagcaga	ttgtaactgag	agtgcaccat	6060
atgcggtgtg	aaataccgca	cagatgcgta	aggagaaaaa	accgcacatc	gcgctcttcc	6120
gcttctcgcg	tcactgactc	gctgcgctcg	gtcgttcggc	tgccgagcgc	ggatcacgct	6180
cactcaaagg	cggtataacg	gttatccaca	gaatcagggg	ataacgcagg	aaagaacatg	6240
tgagcaaaa	gccagcaaaa	ggccaggaa	cgtaaaaagg	ccgcgttgct	ggcggttttc	6300
cataggctcc	gccccctga	cgagcatcac	aaaaatcgac	gctcaagtca	gaggtggcga	6360
aaaccgacag	gactataaag	ataccaggcg	tttccccctg	gaagctccct	cgtgcgctct	6420
cctgttccga	ccctgcgct	taccggatac	ctgtccgcct	ttctcccttc	gggaagcggt	6480
gcgctttctc	atagctcacg	ctgtaggat	ctcagttcgg	tgtaggtcgt	tcgctccaag	6540
ctgggctgtg	tgacgaaacc	ccccgttcag	ccgcaccgct	gcgccttatc	cggttaactat	6600
cgtcttgagt	ccaaccgggt	aagacacgac	ttatcgccac	tgccagcagc	cactggtaac	6660
aggattagca	gagcgaggta	tgtagggcgt	gctacagagt	tcttgaagt	gtggcctaac	6720
tacggctaca	ctagaaggac	agtatttggt	atctgcgctc	tgctgaagcc	agttaccttc	6780
ggaaaaagag	ttggtagctc	ttgatccggc	aaacaaacca	ccgctggtag	cggtgggttt	6840
tttgtttgca	agcagcagat	taagcgcaga	aaaaaaggat	ctcaagaaga	tcctttgatc	6900
ttttctacgg	ggtctgacgc	tcagtggaa	gaaaactcac	gttaagggat	tttgggtcat	6960
agattatcaa	aaaggatctt	cacctagatc	cttttaaat	aaaaatgaag	ttttaaatca	7020
atctaaagta	tatatgagta	aacttggtct	gacagttacc	aatgcttaat	cagtgaaggca	7080
cttatctcag	cgatctgtct	atttcgttca	tccatagttg	cctgactccc	cgtcgtgtag	7140
ataactacga	tacgggaggg	cttaccatct	ggccccagtg	ctgcaatgat	accgcgagac	7200
ccacgctcac	cggctccaga	tttatcagca	ataaaccagc	cagccggaag	ggccgagcgc	7260
agaagtggtc	ctgcaacttt	atccgcctcc	atccagtcct	ttaattgttg	ccgggaagct	7320
agagtaagta	gttcgccagt	taatatgttg	cgcaacggtg	ttgccattgc	tgccaggtcgg	7380
gagcacagga	tgacgcctaa	caattcattc	aagccgacac	cgcttcgagg	cgccgcttaa	7440
ttcaggaggt	aaacatcatg	agggaagcgg	tgatcgccga	agtatcgact	caactatcag	7500
aggtagttgg	cgtcatcgag	cgccatctcg	aaccgacgtt	gctggccgta	catttgttacg	7560
gctccgaggt	ggatggcggc	ctgaagccac	acagtgatat	tgatttgctg	gttacgggtga	7620
ccgtaaggct	tgatgaaaca	acgcgcgag	ctttgatcaa	cgaccttttg	gaaacttcgg	7680
cttccctcgg	agagagcgag	attctccgcg	ctgtagaagt	caccattggt	gtgcacgacg	7740
acatcattcc	gtggcggttat	ccagctaagc	gcgaactgca	atttgagaaa	tgccagcgca	7800
atgacattct	tgccaggtatc	ttcgagccag	ccacgatcga	cattgatctg	gctatcttgc	7860



tgacaaaagc	aagagaacat	agcgttgcc	tggtagggtcc	agcggcggag	gaactccttg	7920
atccggttcc	tgaacaggat	ctatttgagg	cgctaaatga	aaccttaacg	ctatggaact	7980
cgccgcccga	ctgggctggc	gatgagcgaa	atgtagtgct	tacgttgctc	cgcatcttggt	8040
acagcgcagt	aacccggcaaa	atcgcgcgca	aggatgtcgc	tgaagactgg	gcaatggagc	8100
gcctgccggc	ccagtatcag	cccgtcatac	ttgaagctag	gcaggcttat	cttggaacaag	8160
aagatcgctt	ggcctcgcg	gcagatcagt	tggaagaatt	tggtcactac	gtgaaaggcg	8220
agatcaccaa	ggtagtcggc	aaataatgtc	taacaattcg	ttcaagccga	cgccgcttcg	8280
ggcgcgggct	taactcaagc	gttagatgct	gcaggcatcg	tggtgtcacg	ctcgtcgttt	8340
ggtatggctt	cattcagctc	cggttcccaa	cgatcaaggc	gagttacatg	atcccccatg	8400
ttgtgcaaaa	aagcggttag	ctccttcggg	cctccgatcg	aggatttttc	ggcgctgcgc	8460
tacgtccgck	accgcgttga	gggatcaagc	cacagcagcc	cactcgacct	ctagccgacc	8520
cagacgagcc	aagggatctt	tttggaatgc	tgctccgtcg	tcaggctttc	cgacgtttgg	8580
gtgggtgaac	agaagtcatt	atcgtaacga	atgccaaagc	ctcccaggag	gaacctgtg	8640
gttggcatgc	acatacaaat	ggacgaacgg	ataaaccttt	tcacgccctt	ttaaatatcc	8700
gttattctaa	taaacgctct	tttctcttag	gtttaccgcg	caatatatcc	tgtcaaacac	8760
tgatgcttta	aactgaaggc	gggaaacgac	aatctgatcc	ccatcaagct	tggtcgagt	8820
gaagctagct	tcccgatcct	atctgtcact	tcatacaaa	gacagtagaa	aaggaaagtg	8880
gcactacaaa	tgccatcatt	gcgataaagg	aaaggctatc	gttcaagatg	cctctgccga	8940
cagtggtccc	aaagatggac	ccccacccac	gaggagcatc	gtggaaaaag	aagacgttcc	9000
aaccacgtct	tcaaagcaag	tggattgatg	tgatacttcc	actgacgtaa	gggatgacgc	9060
acaatcccac	tatccttcgc	aagacccttc	ctctatataa	ggaagttcat	ttcatcttga	9120
gaggacacgc	tgaatcacc	agtctctctc	tacaagatcg	gggatctcta	gctagacgat	9180
cgtttcgcat	gattgaacaa	gatggattgc	acgcagggtc	tcgggccgct	tggtgggaga	9240
ggctattcgg	ctatgactgg	gcacaacaga	caatcggctg	ctctgatgcc	gccgtgttcc	9300
ggctgtcagc	gcagggggcg	ccggttcttt	ttgtcaagac	cgacctgtcc	ggtgccctga	9360
atgaactgca	ggacgaggca	gcgcggctat	cgtggctggc	cacgacgggc	gttccttgcg	9420
cagctgtgct	cgacgttgct	actgaagcgg	gaagggaactg	gctgctattg	ggcgaagtgc	9480
cggggagga	tctcctgtca	tctcaccttg	ctcctgccga	gaaagtatcc	atcatggctg	9540
atgcaatgcy	gcggctgcat	acgcttgatc	cggtacactg	cccattcgac	caccaagcga	9600
aacatcgcat	cgagcgagca	cgtactcgga	tggaagccgg	tcttgtcgat	caggatgatc	9660
tggaacgaaga	gcatcagggg	ctcgcgcag	ccgaactgtt	cgccaggctc	aaggcgcgca	9720
tgcccgacgg	cgaggatctc	gtcgtgacct	atggcgatgc	ctgcttgccg	aatatcatgg	9780
tggaataatg	ccgcttttct	ggattcatcg	actgtggccg	gctgggtgtg	gcggaccgct	9840
atcaggacat	agcgttggct	acccgtgata	ttgctgaaga	gcttgccggc	gaatgggctg	9900
accgcttctc	cgtgctttac	ggtatcgccg	ctcccatttc	gcagcgcatc	gccttctatc	9960
gccttcttga	cgagttcttc	tgagcgggac	tctgggggtc	gatccccaat	tcccgatcgt	10020
tcaaacattt	ggcaataaag	tttcttaaga	ttgaatcctg	ttgccggctc	tgcatgatt	10080
atcatataat	ttctgttgaa	ttacgttaag	catgtaataa	ttaacatgta	atgcatgacg	10140
ttatztatga	gatgggtttt	tatgattaga	gtcccgaat	tatacattha	atacgcgata	10200
gaaaacaaaa	tatagcgcgc	aaactaggat	aaattatcgc	gcgcgggtgc	atctatgtta	10260
ctagatcggg	gatcggggca	ctcgaccaag	cttctgcagg	tcctgctcga	gc	10312

&lt;210&gt; 16

&lt;211&gt; 8349

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;221&gt; gene

&lt;222&gt; 3666-5573;

&lt;223&gt; completely synthesized

&lt;400&gt; 16

gcaactgttg	ggaagggcga	tcgggtcggg	cctcttctgct	attacgccag	ctggcgaaaag	60
ggggatgtgc	tgcaaggcga	ttaaagttggg	taacgccagg	gttttcccag	tcacgacgtt	120
gtaaaacgac	ggccagtga	ttgcggccac	gcgtgggtacc	aagcttcccg	atcctatctg	180
tcacttcatc	aaaaggacag	tagaaaagga	aggtggcacc	tacaaatgcc	atcattgcga	240
taaaggaaag	gctatcattc	aagatgcctc	tgccgacagt	gggtcccaaag	atggaccccc	300



accacagagg	agcatcgtgg	aaaaagaaga	cgttccaacc	acgtcttcaa	agcaagtggg	360
ttgatgtgat	acttccactg	acgtaaggga	atgacgcaca	atcccaactat	ccttcgcaag	420
acccttcctc	tatataagga	agttcatttc	atttggagag	gacacgctga	aatcaccagt	480
ctctctctac	aagatcgggg	atctctagct	agacgatcgt	ttcgcatgat	tgaacaagat	540
ggattgcacg	caggttctcc	ggcgcgttgg	gtggagaggc	tattcggtcta	tgactgggca	600
caacagacaa	tgggtgctc	tgatgccgcc	gtgttccggc	tgtagcgca	ggggcgcccg	660
gttctttttg	tcaagaccga	cctgtccggg	gccctgaatg	aactgcagga	cgaggcagcg	720
cggtatcgt	ggctggccac	gacgggcgtt	ccttgcgcag	ctgtgctcga	cgttgtcact	780
gaagcgggaa	gggactggct	gctattgggc	gaagtgcggg	ggcaggatct	cctgtcatct	840
caccttgctc	ctgccgagaa	agtatccatc	atggctgatg	caatgcggcg	gctgcatacg	900
cttgatccgg	ctacctgccc	attcgaccac	caagcgaac	atcgcatcga	gcgagcacgt	960
actcggatgg	aagccgggtc	tgtcgatcag	gatgatctgg	acgaagagca	tcaggggctc	1020
gcgcagccg	aactgttcgc	caggctcaag	gcgcgcatgc	ccgacggcga	ggatctcgtc	1080
gtgacccatg	gcgatgcctg	cttgccgaat	atcatggtgg	aaaatggccg	cttttctgga	1140
ttcatcgact	gtggccggct	gggtgtggcg	gaccgctatc	aggacatagc	gttggctacc	1200
cgtgatattg	ctgaagagct	tggcggcgaa	tgggtgacc	gcttccctcg	gctttacggt	1260
atcgccgctc	ccgattcgca	gcgcacgcc	ttctatcgcc	ttcttgacga	gttcttctga	1320
gcgggactct	gggggttcgaa	atgaccgacc	aagcgcagcc	caacctgcc	tcacgagatt	1380
tccgatccac	cgccgccttc	tatgaaaggt	tgggcttcgg	aatcgtttcc	cgggacgccc	1440
gctggatgat	cctccagcgc	ggggatctca	tgctggagtt	cttcgccccac	ccccggatcc	1500
ccatgggaat	tcccgatcgt	tcaaacattt	ggcaataaag	tttcttaaga	ttgaatcctg	1560
ttgccggctc	tgcatgatgt	atcatataat	ttctgttgaa	ttacgttaag	catgtaataa	1620
ttaacatgta	atgcatgacg	ttatttatga	gatgggtttt	tatgattaga	gtcccgcaat	1680
tatacattta	atacgcgata	gaaaaacaaa	tatagcgcgc	aaactaggat	aaattatcgc	1740
gcgcgggtgc	atctatgtta	ctagatcggg	gatatccccg	cggccgcggt	aacaagcttc	1800
tgagggtccg	atgtgagact	tttcaacaaa	gggtaatatc	cggaaacctc	ctcggattcc	1860
attgcccagc	tatctgtcac	tttattgtga	agatagtggg	aaaggaagg	ggctcctaca	1920
aatgccatca	ttgcgataaa	ggaaaggcca	tcgttgaaaga	tgccctctgc	gacagtggct	1980
ccaaagatgg	acccccaccc	acgaggagca	tcgtggaaaa	agaagacgtt	ccaaccacgt	2040
cttcaaagca	agtggattga	tgtgatggtc	cgatgtgaga	cttttcaaca	aagggttaata	2100
tccggaaacc	tccctcgatt	ccattgccca	gctatctgtc	actttattgt	gaagatagt	2160
gaaaaggaag	gtggctccta	caaatgccat	cattgcgata	aaggaaaggc	catcgttgaa	2220
gatgcctctg	ccgacagtgg	tcccaaagat	ggacccccac	ccacgaggag	catcgtggaa	2280
aaagaagacg	ttccaaaccac	gtcttcaag	caagtggatt	gatgtgatat	ctccactgac	2340
gtaagggatg	acgcacaatc	ccactatcct	tcgcaagacc	cttccctctat	ataaggaaat	2400
tcattttcatt	tgagaggagc	acgctgacaa	gctgactcta	gcagatctac	cgtcttcggt	2460
acgcgctcac	tccgcccctc	gcctttgtta	ctgccacgtt	tctctgaatg	ctctcttggt	2520
tgggtgattg	tgagagtggg	ttagctggat	ctgaattac	actctgaaat	cgtgtctctg	2580
ctgtgctgat	tacttgccgt	cctttgtagc	agcaaaatat	agggacatgg	tagtacgaaa	2640
cgaagataga	acctacacag	caatacgaga	aatgtgtaat	ttgggtgctta	gcgggtattta	2700
tttaagcaca	tggttggtgt	atagggcact	tggattcaga	agtttgctgt	taatttaggc	2760
acaggcttca	tactacatgg	gtcaatagta	tagggattca	tattataggc	gatactataa	2820
taattttgttc	gtctgcagag	cttattattt	gccaaaatta	gatattccta	ttctgttttt	2880
gtttgtgtgc	tgtaaaattg	ttaacgcctg	aaggaaataa	tataaatgac	gaaattttga	2940
tgtttatctc	tgctccttta	ttgtgacat	aagtcaagat	cagatgcact	tgtttttaaat	3000
attgttgtct	gaagaaataa	gtactgacag	tattttgatg	cattgatctg	cttggttggt	3060
gtaacaaaat	ttaaaaataa	agagtttcct	ttttgttgct	ctccttacct	cctgatggta	3120
tctagtatct	accaactgac	actatattgc	ttctctttac	atacgtatct	tgctcgatgc	3180
cttctcccta	gtgttgacca	gtgttactca	catagctctt	gctcatttca	ttgtaatgca	3240
gataccaagc	ggcctctaga	ggatcagcat	ggcgccacc	gtgatgatgg	cctcgctggc	3300
caccgcgctc	gctccgttcc	tggggctcaa	gtccaccgcc	agcctccccg	tcgcccgccg	3360
ctcctccaga	agcctcggca	acgtcagcaa	cggcggaagg	atccggtgca	tcgaggtaac	3420
aaatgcatcc	tagctagtag	ttctttgcat	tgcagcagct	gcagctagcg	agttagtaat	3480
aggaagggaa	ctgatgatcc	atgcatggac	tgatgtgtgt	tgcccatccc	atcccatccc	3540
atttcccaaa	cgaaccgaaa	acacgcgtact	acgtgcaggt	gtggccctac	ggcaacaaga	3600
agttcgagac	gctgtcgta	ctgccgcgcg	tgtagaccgg	cgggcgcac	cgctgcagtc	3660
aggccatgga	caactccgtc	ctgaactctg	gtcgaccac	catctgcgac	gcctacaacg	3720
tcgcgcgca	tgatccattc	agcttccagc	acaagagcct	cgacactgtt	cagaaggagt	3780
ggacggagtg	gaagaagaac	aaccacagcc	tgtacctgga	ccccatcgct	ggcaggggtg	3840
ccagcttcc	tctcaagaag	gtcggctctc	tcgtcgggaa	gcgcacctc	tcggaactcc	3900
gcaacctgat	ctttccatct	ggctccacca	acctcatgca	agacatcctc	agggagaccg	3960

agaagtttct	caaccagcgc	ctcaacactg	atacccttgc	tcgcgtcaac	gctgagctga	4020
cgggtctgca	agcaaacgtg	gaggagttca	accgccaaagt	ggacaacttc	ctcaacccca	4080
accgcaatgc	ggtgcctctg	tccatcactt	cttcctgtgaa	caccatgcaa	caactgttcc	4140
tcaaccgcct	gcctcagttc	cagatgcaa	gctaccagct	gctcctgctg	ccactctttg	4200
ctcagggtgc	caacctgcac	ctctccttca	ttcgtgacgt	gatcctcaac	gctgacgagt	4260
ggggcatctc	tcagccacg	ctgaggacct	accgcgacta	cctgaagaac	tacaccaggg	4320
actactccaa	ctattgcatc	aacacctacc	agtcggcctt	caagggcctc	aatacagaggc	4380
ttcacgacat	gctggagtgc	aggacctaca	tgttcctgaa	cgtgttcgag	tacgtcagca	4440
tctggctcgt	cttcaagtac	cagagcctgc	tggtgtccag	cggcgccaac	ctctacgcca	4500
gcggctctgg	tccccacaa	actcagagct	tcaccagcca	ggactggcca	ttcctgtatt	4560
cgttggtcca	agtcactcc	aactacgtcc	tcaacggctt	ctctggtgct	cgcctctcca	4620
acaccttccc	caacattggt	ggcctccccc	gtctccaccac	aactcatgct	ctgcttgctg	4680
ccagagtga	ctactccggc	ggcatctcga	gcggcgacat	tggtgcatcg	ccggtcaacc	4740
agaacttcaa	ctgctccacc	ttcctgcgcg	cgtgctcac	cccgttcgtg	aggtcctggc	4800
tcgacagcgg	ctccgaccgc	gagggcgtgg	ccaccgtcac	caactggcaa	accgagtcct	4860
tcgagaccac	ccttggcctc	cggagcggcg	ccttcacggc	gcgtgggaat	tctaactact	4920
tccccgacta	cttcatcagg	aacatctctg	gtgttctctt	cgtcgtccgc	aacgaggacc	4980
tccgcgctcc	actgcactac	aacgagatca	ggaacatcgc	ctctccgtcc	gggacgccc	5040
gaggtgcaag	ggcgatcatg	gtgagcgtcc	ataacaggaa	gaacaacatc	cacgctgtgc	5100
atgagaacgg	ctccatgatc	cacctggcgc	ccaatgatta	caccggcttc	accatctctc	5160
caatccacgc	cacccaagtg	aacaaccaga	cacgcacctt	catctccgag	aagttcggca	5220
accagggcga	ctccctgagg	ttcgagcaga	acaacaccac	cggcaggta	acctgcgcgc	5280
gcaacggcaa	cagctacaac	ctgtacctgc	gcgtcagctc	cattggcaac	tccaccatca	5340
gggtcaccat	caacgggagg	gtgtacacag	ccaccaatgt	gaacacgacg	accaacaatg	5400
atggcgctca	cgacaacggc	gcccgcctca	gcgacatcaa	cattggcaac	gtggtggcca	5460
gcagcaactc	cgacgtcccg	ctggacatca	acgtgacctt	gaactctggc	accagttcgc	5520
acctcatgaa	catcatgctg	gtgccaaacta	acatctcgcc	gctgtactga	taggagctct	5580
gatccccatg	ggaattcccg	atcggttcaa	catttggcaa	taaagtttct	taagattgaa	5640
tctgtttgcc	ggtcttgcca	tgattatcat	ataatttctg	ttgaattacg	ttaagcatgt	5700
aataattaac	atgtaatgca	tgacgttatt	tatgagatgg	gtttttatga	ttagagtcct	5760
gcaattatac	atttaatacg	cgatagaaaa	caaaatatag	cgcgcaaaact	aggataaatt	5820
atcgcgcgcg	gtgtcatcta	tgttactaga	tcggggat	ccccggggcg	gcccggggga	5880
attcggtacc	aagcttacgc	gtggccgcag	cttggcgtaa	tcattggtcat	agctgtttcc	5940
tggtgaaat	tgttatccgc	tcacaattcc	acacaacata	cgagccggaa	gcataaagtg	6000
taaagcctgg	ggtgccta	gagtgagcta	actcacatta	attgcgttgc	gctcactgcc	6060
cgttttcag	tcgggaaacc	tgctgtgcca	gctgcattaa	tgaatcgccc	aacgcgcggg	6120
gagaggcgg	ttgcgtattg	ggcgtctctc	cgttctctcg	ctcactgact	cgtctgcctc	6180
ggtcgttcgg	ctgcggcgag	cggtatcagc	tcactcaaag	gcggtataac	ggttatccac	6240
agaatcaggg	gataacgcag	gaaagaacat	gtgagcaaaa	ggccagcaaa	aggccaggaa	6300
ccgtaaaaag	gcccgcgttc	tggcgttttt	ccataggctc	cgcctccctg	acgagcatca	6360
caaaaatcga	cgctcaagtc	agaggtggcg	aaaccgcaga	ggactataaa	gataccaggc	6420
gtttccccct	ggaagctccc	tcgtgcgctc	tctgttccg	acctgcgcgc	ttaccggata	6480
cctgtccgcc	tttctccctt	cgggaagcgt	ggcgtttct	caatgctcac	gctgtaggta	6540
tctcagttcg	gtgtaggctg	ttcgtcccaa	gctgggctgt	gtgcacgaac	ccccggttca	6600
gcccgaaccg	tcgccttat	ccggtaaacta	tcgtcttgag	tccaaccccg	taagacacga	6660
cttatcgcca	ctggcagcag	ccactggtaa	caggattagc	agagcgaggt	atgtaggcgg	6720
tgctacagag	ttcttgaagt	ggtggcctaa	ctacggctac	actagaagga	cagtatttgg	6780
tatctgcgct	ctgctgaagc	cagttacctt	cggaaaaaga	gttggtagct	cttgatccgg	6840
caaacaaaacc	accgctggta	gcggtggttt	ttttgtttgc	aagcagcaga	ttacgcgcag	6900
aaaaaaagga	tctcaagaag	atcctttgat	cttttctacg	gggtctgacg	ctcagtgga	6960
cgaaaactca	cgttaaggga	ttttggtcat	gagattatca	aaaaggatct	tcacctagat	7020
ccttttgggg	tgggcgaaga	actccagcat	gagatccccg	cgtggaggga	tcattccagcc	7080
ggcgtcccgg	aaaacgattc	cgaagcccaa	cctttcatag	aaggcggcgg	tggaaatcgaa	7140
atctcgtgat	ggcaggttgg	gcgtcgcttg	gtcggctcatt	tcgaacccca	gagtcgccgt	7200
cagaagaact	cgtcaagaag	gcgatagaag	gcgatgcgct	gcgaatcggg	agcggcgata	7260
ccgtaaaagca	cgaggaagcg	gtcagcccat	tcgcgcgcaa	gctcttcagc	aatatcacgg	7320
gtagccaacg	ctatgtcctg	atagcggctc	gccacacca	gcgggccaca	gtcgatgaat	7380
ccagaaaagc	ggccattttc	caccatgata	ttcggaagc	aggcatcgcc	atgggtcacg	7440
acgagatcct	cgcgcgtcgg	catgcgcgcc	ttgagcctgg	cgaacagttc	ggctggcgcg	7500
agcccctgat	gctcttcgct	cagatcatcc	tgatcgacaa	gaccggcttc	catccgagta	7560
cgtgctcgct	cgatgcgatg	tttcgcttgg	tggtcgaatg	ggcaggtagc	cggatcaagc	7620

gtatgcagcc	gccgcattgc	atcagccatg	atggatactt	tctcggcagg	agcaagggtga	7680
gatgacagga	gacccctgcc	cggcaacttc	cccaatagca	gccagtcctt	tcccgttca	7740
gtgacaacgt	cgagcacagc	tgcgcaagga	acgcccgtcg	tggccagcca	cgatagccgc	7800
gctgcctcgt	cctgcagttc	attcagggca	cgggacaggt	cggctctgac	aaaaagaacc	7860
ggcgccccct	gcgctgacag	ccggaacacg	gcggcatcag	agcagccgat	tgtctgttgt	7920
gccagtcgat	agccgaatag	cctctccacc	caagcggccg	gagaacctgc	gtgcaatcca	7980
tcttgttcaa	tcattgcgaaa	cgatccctcat	cctgtctctt	gatcagatct	tgatccccctg	8040
cgccatcaga	tccttggcgg	caagaaagcc	atccagttta	ctttgcaggg	cttcccaacc	8100
ttaccagagg	gcgccccagc	tggcaattcc	ggttcgcttg	ctgtccataa	aaccgccag	8160
tctagctatc	gccatgtaag	cccactgcaa	gctacctgct	ttctctttgc	gcttgcgttt	8220
tcccttgtcc	agatagccca	gtagctgaca	ttcatccggg	gtcagcaccg	tttctgcgga	8280
ctggctttct	acgtgttccg	cttcccttag	cagcccttgc	gccctgagtg	cttgccggcag	8340
cgatgaagct						8349

&lt;210&gt; 17

&lt;211&gt; 1912

&lt;212&gt; DNA

<213> *Bacillus thuringiensis*

&lt;400&gt; 17

atgaatagtg	tattgaatag	cggaagaact	actatattgtg	atgcgataaa	tgtagcggct	60
catgatccat	ttagttttca	acacaaatca	ttagataccg	tacaaaagga	atggacggag	120
tggaaaaaaa	ataatcatag	tttataccta	gacccatttg	ttggaaactgt	ggctagtttt	180
ctgttaaaga	aagtggggag	tcttgtttga	aaaaggatac	taagttaggt	acggaattta	240
atatttccta	gtggtagtac	aaatctaata	caagatattt	taagagagac	agaaaaattc	300
ctgaatcaaa	gacttaatac	agacactcct	gcccgtgtaa	atgcggaatt	gacagggctg	360
caagcaaatg	tagaagagtt	taatcgacaa	gtagataaatt	ttttgaacc	taaccgaaac	420
gctgttcctt	tatcaataac	ttcttcagtt	aatacaatgc	aacaattatt	tctaaataga	480
ttaccccgat	tccagatgca	aggataccaa	ctgttattat	tacctttatt	tgcacaggca	540
gccaatttac	atctttcttt	tattagagat	gttattctaa	atgcagatga	atggggaatt	600
tcagcagcaa	cattacgtac	gtatcgagat	tacttgaaaa	attatacaag	agattactct	660
aactattgta	taaatacgta	tcaaagtgcg	tttaaagggt	taaacactcg	tttacacgat	720
atgttagaat	ttagaacata	tatgttttta	aatgtatttg	agtatgtatc	tatctggtcg	780
ttgtttaaat	atcaaagtct	tctagtatct	tccggtgcta	atttatatgc	aagtggtagt	840
ggaccacagc	agacccaatc	atttacttca	caagactggc	catttttata	ttctcttttc	900
caagttaatt	caaattatgt	gttaaattgga	tttagtggtg	ctaggctttc	taataccttc	960
cctaataatag	ttggttttacc	tggttctact	acaactcacg	cattgcttgc	tgcaagggtt	1020
aattacagtg	gaggaatttc	gtctgggtgat	ataggtgcat	ctccgtttaa	tcaaaatttt	1080
aattgtagca	catttctccc	cccattgtta	acgccatttg	ttaggagttg	gctagattca	1140
ggttcagatc	gggagggcgt	tgccaccgtt	acaaattggc	aaacagaatc	ctttgagaca	1200
acttttaggt	taaggagtgg	tgcttttaca	gctcgcggta	attcaaaacta	tttcccagat	1260
tattttattc	gtaatatattc	tggagttcct	ttagttgtta	gaaatgaaga	tttaagaaga	1320
ccgttacact	ataatgaaat	aagaaatata	gcaagtcctt	caggaacacc	tggtggagca	1380
cgagcttata	tgggtatctgt	gcataacaga	aaaaataata	tccatgctgt	tcatgaaat	1440
ggttctatga	ttcatttagc	gccaaatgac	tatacaggat	ttactatttc	gccgatacat	1500
gcaactcaag	tgaataatca	aacacgaaca	tttatttctg	aaaaatttgg	aatcaagggt	1560
gattctttta	ggtttgaaca	aaacaacacg	acagctcggt	atagccttag	aggggaatgga	1620
aatagttaca	atctttattt	aagagtttct	tcaataggaa	attccactat	tcgagttact	1680
ataaacggta	gggtatatac	tgctacaaat	gttaatacta	ctacaaataa	cgatggagtt	1740
aatgataatg	gagctcggtt	ttcagatatt	aatatcggta	atgtagtagc	aagtagtaat	1800
tctgatgtac	cattagatat	aaatgtaaca	ttaaactccg	gtactcaatt	tgatcttatg	1860
aatattatgc	ttgtaccaac	taatatattca	ccactttatt	aaggtttgag	ta	1912

&lt;210&gt; 18

&lt;211&gt; 633

&lt;212&gt; PRT

<213> *Bacillus thuringiensis*

&lt;400&gt; 18

Met Asn Ser Val Leu Asn Ser Gly Arg Thr Thr Ile Cys Asp Ala Tyr  
 1 5 10 15  
 Asn Val Ala Ala His Asp Pro Phe Ser Phe Gln His Lys Ser Leu Asp  
 20 25 30  
 Thr Val Gln Lys Glu Trp Thr Glu Trp Lys Lys Asn Asn His Ser Leu  
 35 40 45  
 Tyr Leu Asp Pro Ile Val Gly Thr Val Ala Ser Phe Leu Leu Lys Lys  
 50 55 60  
 Val Gly Ser Leu Val Gly Lys Arg Ile Leu Ser Glu Leu Arg Asn Leu  
 65 70 75 80  
 Ile Phe Pro Ser Gly Ser Thr Asn Leu Met Gln Asp Ile Leu Arg Glu  
 85 90 95  
 Thr Glu Lys Phe Leu Asn Gln Arg Leu Asn Thr Asp Thr Leu Ala Arg  
 100 105 110  
 Val Asn Ala Glu Leu Thr Gly Leu Gln Ala Asn Val Glu Glu Phe Asn  
 115 120 125  
 Arg Gln Val Asp Asn Phe Leu Asn Pro Asn Arg Asn Ala Val Pro Leu  
 130 135 140  
 Ser Ile Thr Ser Ser Val Asn Thr Met Gln Gln Leu Phe Leu Asn Arg  
 145 150 155 160  
 Leu Pro Gln Phe Gln Met Gln Gly Tyr Gln Leu Leu Leu Leu Pro Leu  
 165 170 175  
 Phe Ala Gln Ala Ala Asn Leu His Leu Ser Phe Ile Arg Asp Val Ile  
 180 185 190  
 Leu Asn Ala Asp Glu Trp Gly Ile Ser Ala Ala Thr Leu Arg Thr Tyr  
 195 200 205  
 Arg Asp Tyr Leu Lys Asn Tyr Thr Arg Asp Tyr Ser Asn Tyr Cys Ile  
 210 215 220  
 Asn Thr Tyr Gln Ser Ala Phe Lys Gly Leu Asn Thr Arg Leu His Asp  
 225 230 235 240  
 Met Leu Glu Phe Arg Thr Tyr Met Phe Leu Asn Val Phe Glu Tyr Val  
 245 250 255  
 Ser Ile Trp Ser Leu Phe Lys Tyr Gln Ser Leu Leu Val Ser Ser Gly  
 260 265 270  
 Ala Asn Leu Tyr Ala Ser Gly Ser Gly Pro Gln Gln Thr Gln Ser Phe  
 275 280 285  
 Thr Ser Gln Asp Trp Pro Phe Leu Tyr Ser Leu Phe Gln Val Asn Ser  
 290 295 300  
 Asn Tyr Val Leu Asn Gly Phe Ser Gly Ala Arg Leu Ser Asn Thr Phe  
 305 310 315 320  
 Pro Asn Ile Val Gly Leu Pro Gly Ser Thr Thr Thr His Ala Leu Leu  
 325 330 335  
 Ala Ala Arg Val Asn Tyr Ser Gly Gly Ile Ser Ser Gly Asp Ile Gly  
 340 345 350  
 Ala Ser Pro Phe Asn Gln Asn Phe Asn Cys Ser Thr Phe Leu Pro Pro  
 355 360 365  
 Leu Leu Thr Pro Phe Val Arg Ser Trp Leu Asp Ser Gly Ser Asp Arg  
 370 375 380  
 Glu Gly Val Ala Thr Val Thr Asn Trp Gln Thr Glu Ser Phe Glu Thr  
 385 390 395 400  
 Thr Leu Gly Leu Arg Ser Gly Ala Phe Thr Ala Arg Gly Asn Ser Asn  
 405 410 415  
 Tyr Phe Pro Asp Tyr Phe Ile Arg Asn Ile Ser Gly Val Pro Leu Val  
 420 425 430  
 Val Arg Asn Glu Asp Leu Arg Arg Pro Leu His Tyr Asn Glu Ile Arg  
 435 440 445  
 Asn Ile Ala Ser Pro Ser Gly Thr Pro Gly Gly Ala Arg Ala Tyr Met  
 450 455 460  
 Val Ser Val His Asn Arg Lys Asn Asn Ile His Ala Val His Glu Asn  
 465 470 475 480  
 Gly Ser Met Ile His Leu Ala Pro Asn Asp Tyr Thr Gly Phe Thr Ile

															485										490										495				
Ser	Pro	Ile	His	Ala	Thr	Gln	Val	Asn	Asn	Gln	Thr	Arg	Thr	Phe	Ile																								
															500					505					510														
Ser	Glu	Lys	Phe	Gly	Asn	Gln	Gly	Asp	Ser	Leu	Arg	Phe	Glu	Gln	Asn																								
															515					520					525														
Asn	Thr	Thr	Ala	Arg	Tyr	Thr	Leu	Arg	Gly	Asn	Gly	Asn	Ser	Tyr	Asn																								
															530					535					540														
Leu	Tyr	Leu	Arg	Val	Ser	Ser	Ile	Gly	Asn	Ser	Thr	Ile	Arg	Val	Thr																								
															545					550					555					560									
Ile	Asn	Gly	Arg	Val	Tyr	Thr	Ala	Thr	Asn	Val	Asn	Thr	Thr	Thr	Asn																								
															565					570					575														
Asn	Asp	Gly	Val	Asn	Asp	Asn	Gly	Ala	Arg	Phe	Ser	Asp	Ile	Asn	Ile																								
															580					585					590														
Gly	Asn	Val	Val	Ala	Ser	Ser	Asn	Ser	Asp	Val	Pro	Leu	Asp	Ile	Asn																								
															595					600					605														
Val	Thr	Leu	Asn	Ser	Gly	Thr	Gln	Phe	Asp	Leu	Met	Asn	Ile	Met	Leu																								
															610					615					620														
Val	Pro	Thr	Asn	Ile	Ser	Pro	Leu	Tyr																															
															625					630																			

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/US 99/26086A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 C12N15/12 C12N15/82 C12N5/10 A01H5/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A01H C07K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WONG E Y ET AL: "ARABIDOPSIS THALIANA SMALL SUBUNIT LEADER AND TRANSIT PEPTIDE ENHANCE THE EXPRESSION OF BACILLUS THURINGIENSIS PROTEINS IN TRANSGENIC PLANTS" PLANT MOLECULAR BIOLOGY, NL, NIJHOFF PUBLISHERS, DORDRECHT, vol. 20, no. 1, 1 October 1992 (1992-10-01), pages 81-93, XP000570449 ISSN: 0167-4412 the whole document — —/—	2,4,6,9, 11,13, 15-26, 28,30, 32,34, 36,38, 40-56

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"S" document member of the same patent family

Date of the actual completion of the international search

7 March 2000

Date of mailing of the international search report

24/03/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Burkhardt, P

Form PCT/ISA/210 (second sheet) (July 1992)

page 1 of 2

## INTERNATIONAL SEARCH REPORT

Int. Patent Application No.  
PCT/US 99/26086

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>WIDNER W R ET AL: "TWO HIGHLY RELATED INSECTICIDAL CRYSTAL PROTEINS OF BACILLUS THURINGIENSIS SUBSP. KURSTAKI POSSESS DIFFERENT HOST RANGE SPECIFICITIES" JOURNAL OF BACTERIOLOGY, US, WASHINGTON, DC, vol. 171, no. 2, 1 February 1989 (1989-02-01), pages 965-974, XP002071367 ISSN: 0021-9193 figure 2</p>	1-56
Y	<p>WO 95 24492 A (CALGENE INC ; STATE UNIVERSITY OF NEW JERSEY (US)) 14 September 1995 (1995-09-14)</p> <p>page 13, lines 30-32 page 15, line 7 -page 16, line 9; examples 4,5</p>	1,3,7,8, 10,12, 14,27, 29,31, 33,35, 37,39
Y	<p>US 5 689 052 A (SANDERS PATRICIA RIGDEN ET AL) 18 November 1997 (1997-11-18) SEQ ID NO:2</p>	1-56
Y	<p>EP 0 385 962 A (MONSANTO CO) 5 September 1990 (1990-09-05) claim 2; figure 13</p>	1-56
A	<p>CRICKMORE N ET AL: "Revision of the nomenclature for the Bacillus thuringiensis pesticidal crystal proteins." MICROBIOLOGY AND MOLECULAR BIOLOGY REVIEWS, (1998 SEP) 62 (3) 807-13. REF: 126, XP002132141 page 809 -page 810</p>	1-56
P,X	<p>KOTA M ET AL: "Overexpression of the Bacillus thuringiensis (Bt) Cry2Aa2 protein in chloroplasts confers resistance to plants against susceptible and Bt-resistant insects." PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF THE UNITED STATES OF AMERICA, (1999 MAR 2) 96 (5) 1840-5. , XP002132142 the whole document</p>	1,3,7,8, 10,12, 14,27, 29,31, 33,35, 37,39

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/US 99/26086

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9524492 A	14-09-1995	US 5545818 A CA 2185340 A EP 0749490 A JP 10511261 T	13-08-1996 14-09-1995 27-12-1996 04-11-1998
US 5689052 A	18-11-1997	NONE	
EP 0385962 A	05-09-1990	AU 638438 B AU 5163090 A CA 2024811 A EP 0413019 A IL 93513 A JP 3504333 T NO 303546 B NZ 232654 A WO 9010076 A US 5880275 A US 5500365 A ZA 9001417 A TR 24354 A	01-07-1993 26-09-1990 25-08-1990 20-02-1991 14-11-1995 26-09-1991 27-07-1998 23-12-1991 07-09-1990 09-03-1999 19-03-1996 30-01-1991 01-09-1991

Form PCT/ISA/210 (patent family annex) (July 1992)